IOH. CHRIST. FABRICII

HIST. NAT. OECONOM. ET CAMERAL. P. P. O.
CIET. HAFN. NORW. NAT. CVRIOS. BEROL. LIPS. LVND.
PETR. OECON. TAVR. LOND. HIST. NAT. IENENS.
BOVRD. NEC NON PARIS. AGRICVLT. HIST. NAT.
ET PHILOM. MEMBR.

# SYSTEMA ELEVTHERATORVM

SECVNDVM

ORDINES, GENERA, SPECIES:

ADIECTIS

YNONYMIS, LOCIS, OBSERVATIONIBVS,
DESCRIPTIONIBVS.

TOMVS II.

KILIAE

IMPENSIS BIBLIOPOLII ACADEMICI NOVI.

1801.

Digitized by Google

Original from
NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google



Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google
Digitized by Google
Original from NEW YORK PUBLIC LIBRARY

3. EROTYLVS. *Palpi* quatuor inaequales, *anteriores* malleiformes; *posteriores* subfiliformes. *Antennae* claua compressa perfoliata.

1. E. niger, coleoptris punctis fuluis numerosissimis. Ent. syst. 2. 35. 1. *gigantеus.*
*Chrysomela gigantea.* Linn. syst. Nat. 2. 586. 1.
Oliv. Inf. 89. tab. 1. fig. 6.
Degeer Inf. 5. 349. 1. tab. 16. fig. 8.
Habitat in India.

2. E. ater, elytris flauis, nigro-reticulatis. Ent. syst. 2. 36. 2. * *reticulatus.*
Habitat in Brasilia. Mus. Dr. Hunter.

3. E. ater, elytris fasciis sex vndatis, fuluis: anteriore interrupta. Ent. syst. 2. 36. 3. * *6fasciatus.*
Habitat in America meridionali.

4. E.

---

Erotyli corpus magnum, ouatum, gibbum, immarginatum, agile; *capite* paruo, ouato, inserto; *oculis* rotundatis, parum prominulis, marginalibus; *antennis* thorace longioribus, ante oculos insertis; *thorace* transuerso, plano: margine vtrinque antice prominulo, postice sinuato; *scutello* paruo, rotundato; *elytris* rigidis, gibbis, fornicatis, longitudine elytrorum; *pedibus* longiusculis, tenuibus, cursoriis; *tarsis* quadriarticulatis; *colore* vario.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*hiſtrio.* 4. E. ater, elytris nigro-flauoque faſciatis: macu baſeos apicisque coccinea. Ent. ſyſt. 2. 36. 4
Habitat in Cayenna. Muſ. Dr. Hunter.

*teſtaceus.* 5. E. teſtaceus, antennis tibiisque nigris. Ent. ſy 2. 36. 5. *
Habitat in America. D. Drury.

*ſphacelatus* 6. E. ater, elytris faſciis duabus flauis: punctis gris impreſſis.
Habitat in America meridionali. Muſ. D. de Seb ſtedt.
Corpus magnum, atrum. Elytra baſi faſciaq pone medium flauis: punctis plurimis impr ſis atris.

*gibboſus.* 7. E. niger, elytris flaueſcentibus nigro-punctati faſcia media poſticaque nigris. Ent. ſyſt. 36. 6.
*Chryſomela gibboſa.* Linn. ſyſt. Nat. 2. 586.
Oliv. Inſ. 89. tab. 1. fig. 4.
Herbſt. Arch. tab. 23. fig. 3.
Linn. Amoen. acad. 6. 393. 13.
Gronov. Zooph. 606. tab. 14. fig. 5.
Habitat in America meridionali.

*eleuatus.* 8. E. ater, elytris gibbis, flauis: maculis nigris i preſſo-punctatis.
Habitat - - Muſ. D. Lund.
Minor E. gibboſo. Thorax depreſſus, inaequal ater. Elytra dorſo eleuata, laeuia, flaua, maculis plurimis, irregularibus, atris, et in puncta impreſſa. Corpus atrum.

*notatus.* 9. E. ater, elytris faſcia media flaua, nigro-mac lata, baſi rubro-notatis. Ent. ſyſt. 2. 37. 7.
*Erotylus notatus.* Oliv. Inſ. 89. tab. 1. fig. 1
Habitat in Cayenna. Muſ. D. Olivier.

10.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

0. E. ater, elytris flauo-nigroque reticulatis: fa- *concatena-*
ſciis duabus atris. Ent. ſyſt. 2. 37. 8. * *tus.*
Habitat - - Dr. Schulz.

1. E. elytris nigris: punctis quinque rubris. Ent. *5punctatus.*
ſyſt. 2. 37. 9.
*Erotylus 5punctatus.* Oliv. Inſ. 89. tab. 1. fig. 5.
*Chryſomela 5punctata.* Linn. ſyſt. Nat. 2. 586. 3.
Petiv. Gazoph. 613. tab. 16. fig. 7.
Habitat in America meridionali.

2. E. niger, elytris flauis: punctis numeroſiſſimis *punctatiſſi-*
nigris. Ent. ſyſt. 2. 37. 10. * *mus.*
Habitat in America meridionali. Muſ. Dr. Hunter.

3. E. ater, elytris punctatis medio fuluo-macula- *variegatus.*
tis. Ent. ſyſt. 2. 37. 11. *
*Erotylus variegatus.* Oliv. Inſ. 89. tab. 1. fig. 7.
Habitat Surinami. Muſ. D. Banks.

4. E. flauus, elytris laeuibus teſtaceis: maculis *8maculatus*
quatuor nigris.
Habitat - - Muſ. D. Lund.
Paullo minor E. vnifaſciato. Antennae parum compreſſae, nigrae, baſi apiceque albae. Caput nigrum, ore flauo. Thorax planus, flauus, immaculatus. Elytra laeuia, teſtacea, maculis quatuor nigris: prima humerali, tertia transuerſa. Corpus flaueſcens.

5. E. ater, elytris punctis fuluis, numeroſiſſimis, *gemmatus.*
thorace varioloſo. Ent. ſyſt. 2. 38. 12. *
Habitat in Cayenna. Muſ. D. Boſc.

6. E. ater, elytris faſciis atris flauisque alternis, *longimanus*
pedibus anticis elongatis. Ent. ſyſt. 2. 38. 13. *
Habitat in India. D. Daldorff.

7. E. ater, elytris flauis: faſciis quatuor, vndatis *abdomina-*
punctoque diſtincto nigris. Ent. ſyſt. 2. 38. 14. * *lis.*
Habitat in Braſilia. Muſ. D. Banks.

18. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*vnifaſciatus.* 18. E. ater, elytris flauis: faſcia media lata, atra
Ent. ſyſt. ſuppl. 101. 13. *
Habitat - - Muſ. D. Lund.

*faſciatus.* 19. E. ater, elytris faſciis tribus flaueſcentibus. Ent
ſyſt. 2. 38. 15. *
Habitat in America. D. Drury.
*Os* maxillis palpisque. *Labrum* elongatum, cor
neum, truncatum. *Palpi* quatuor inaequales
*anteriores* longiores, malleiformes, quadria
ticulati: articulo primo breuiſſimo, ſecund
longiori, obconico, tertio breui, rotundat
quarto maximo, ouato, transuerſo, concauo, i
medio articuli tertii inſerto, adhaerentes ma
xillae dorſo, *poſteriores* breuiores, filiforme
triarticulati: articulo primo breuiſſimo, ſecun
do longiori, incuruo, cylindrico, tertio aequal
vix craſſiori, adnati ligulae ante apicem. *Ma
dibula* breuis, cornea, incurua, apice bifid
*Maxilla* breuis, membranacea, cylindrica, ob
tuſiſſima, integra. *Ligula* porrecta, membra
nacea, apice cylindrica, integra, obtuſa, ant
apicem palpigera. *Labium* breue, transue
ſum, corneum, tridentatum.

*4puſtulatus* 20. E. ater, elytris maculis duabus fuluis.
Habitat in Sumatra. D. Daldorff.
Statura et magnitudo E. Zebrae. Antennae nigrae
claua compreſſa: articulis magis approximatis
Caput et thorax atra, immaculata. Elytra lae
uia, atra, maculis duabus magnis, ferrugineis
Corpus atrum.

*Zebra.* 21. E. flaueſcens, capite, thoracis baſi, faſciis tribu
elytrorum, pedibusque atris. Ent. ſyſt. 2. 3
16. *
Habitat in Cayenna. D. v. Rohr.

22. I

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

12. E. niger, coleoptris faſciis duabus flauis: anteriore nigro-punctata. Ent. ſyſt. 2. 39. 17. * *alternans.*

*Erotylus alternans.* Oliv. Inſ. 89. tab. 1. fig. 10.

*Chryſomela Gronouii.* Herbſt. Arch. tab. 23. f. 4.

Habitat in America meridionali.

Nimis affinis E. faſciato, et forte mera eius varietas.

13. E. obſcurus, elytris punctatis, aeneo-nitidulis: ſtrigis tribus dentatis ferrugineis. *dentatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Magnitudo E. Zebrae. Antennae nigrae. Caput et thorax decliuia, obſcura, thoracis margine ferrugineo. Elytra punctata, viridi-aenea, margine ſtrigisque tribus valde dentatis, rufis. Interdum ſtrigae per dentes coëunt, et maculas nigro-aeneas in elytris rufis formant. Corpus obſcurum, pedibus ferrugineis.

14. E. ater, thoracis margine elytrorumque limbo flauo. Ent. ſyſt. 2. 39. 20. * *limbatus.*

*Erotylus limbatus.* Oliv. Inſ. 89. tab. 2.

Habitat in Cayenna. Muſ. D. Olivier.

15. E. brunneus, antennis, ſcutello pectoreque nigris. *brunneus.*

*Erotylus brunneus.* Web. Inſ. 59. 1.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Minor E. alternante. Antennae nigrae, baſi ferrugineae. Caput et thorax brunnea, fere immaculata, interdum linea dorſali fuſca. Scutellum nigrum. Elytra ſublaeuia, brunnea, margine vix pallidiore. Corpus brunneum, pectore tibiisque nigris.

*Variat* corpore pallidiore.

26. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*atratus.* 26. E. ſupra ater, ſubtus pedibusque brunneis.
Habitat in America meridionali. D. Smidt. Mu
D. de Seheſtedt.
Statura omnino praecedentis, at paullo maio
Antennae nigrae, articulis vltimis tribus pa
rum craſſioribus, compreſſis. Thorax decliu
planus, laeuis, niger, obſcurus. Elytra vi
ſtriata, atra, immaculata. Corpus cum pedibu
brunneum.

*rufipennis.* 27. E. ater, nitidus, elytris rufis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Statura et magnitudo praecedentium. Antenna
nigrae. Caput et thorax decliuia, laeuia, atra
immaculata. Scutellum nigrum. Elytra ſubti
liſſime punctato-ſtriata, obſcure rufa. Corpu
atrum.

*tricolor.* 28. E. ferrugineus, elytris flaueſcentibus: diſco ob
ſcuriore.
Habitat in America meridionali. D. Smidt. Mu
D. de Seheſtedt.
Statura omnino E. brunnei, at paullo minor. Ca
put et thorax decliuia, nunc obſcure ferrugi
nea, nunc magis flaueſcentia. Scutellum ni
grum. Elytra punctato-ſtriata, flaueſcentia
futura margineque parum nigricantibus et di
ſco fuſco. Corpus ferrugineum.

*undatus.* 29. E. ferrugineus, elytris nigris, ſtrigis duabus un
datis parique punctorum poſticorum ferru
gineis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Antennae nigrae. Caput et thorax decliuia, fer
ruginea, thorace maculis quatuor nigris. Ely
tri

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tra ſtriata, nigra, baſi dentata: ſtrigis duabus vndatis, punctorum pari apiceque rufis. Corpus ferrugineum.

38. AEGITHVS. *Palpi* quatuor inaequales, *anteriores* malleiformes; *poſteriores* capitati. *Antennae* claua compreſſa, perfoliata.

1. A. niger, elytris rubris immaculatis. *Surinamenſis.*
*Erotylus Surinamenſis.* Ent. ſyſt. 2. 39. 18. *
*Coccinella Surinamenſis.* Linn. ſyſt. Nat. 2. 579. 2. Amoen. acad. 6. 393. 12.
Oliv. Inſ. 89. tab. 1. fig. 9.
Habitat in America meridionali.
Genus affine praecedenti, attamen diſtinctum.
*Os* maxillis palpisque. *Labrum* breue, corneum, rotundatum, integrum. *Palpi* quatuor inaequales, *anteriores* longiores, quadriarticulati, malleiformes: articulo primo breui, ſecundo longiori, cylindrico, tertio breui, rotundato, quarto maximo, transuerſo, concauo, in medio articuli tertii inſerto, adhaerentes maxillae dorſo, *poſteriores* breuiores, capitati, triarticulati: articulo primo ſecundoque breuiſſimis, aequa-

---

Aegithi corpus ſubrotundum, ſupra gibbum, ſubtus planum, glabrum, immarginatum, tardum; *capite* paruo, ouato, inſerto; *oculis* paruis, rotundatis, marginalibus; *antennis* diſtantibus, longitudine thoracis, ante oculos inſertis; *thorace* breui, transuerſo, poſtice latiore; *ſcutello* paruo, rotundato; *elytris* fornicatis, rigidis, longitudine abdominis; *pedibus* breuibus, validis, curſoriis; *tarſis* quadriarticulatis; *colore* obſcuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

aequalibus, rotundatis, tertio magno, craſſo, capitato, adnati ligulae apici. *Mandibula* breuis, cornea, incurua, fiſſa. *Maxilla* breuis, cylindrica, ſubmembranacea, obtuſa. *Ligula* breuis, cornea, plana, integra. *Labium* transuerſum, corneum, tridentatum, dentibus lateralibus maioribus, rotundatis.

*marginatus* 2. A. flaueſcens, elytris nigris: margine omni flaueſcente. Ent. ſyſt. ſuppl. 101. 18. *
Habitat in Cayenna. D. Richard.

*cinctus.* 3. A. ater, elytrorum limbo faſciaque media ferrugineis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura rotundata et magnitudo praecedentium. Antennae nigrae, claua parua, compreſſa. Caput et thorax nigra. Elytra ſubpunctata, nigra, margine omni tenui faſciaque media rufis. Corpus nigrum, pedibus obſcure rufis.

*diſcoideus.* 4. A. ferrugineus, elytrorum diſco cyaneo-nigro.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Summa affinitas A. marginati. Antennae ferrugineae, claua compreſſa, nigra. Caput et thorax ferruginea, immaculata. Elytra laeuia, cyaneo-nigra, margine exteriori, et antico, ferrugineis. Corpus ferrugineum.

*Guadeloupenſis.* 5. A. atra, elytris fuſcis: limbo flaueſcente.
*Galleruca Guadeloupenſis.* Ent. ſyſt. 2. 16. 17. *
Habitat in Inſula Guadeloupe. D. Badier. Muſ. D. de Seheſtedt.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

89. EVMORPHVS. *Palpi* quatuor inaequales, *anteriores* ſubfiliformes; *poſteriores* breues, capitati. *Antennae* claua compreſſa, perfoliata.

1. E. niger, elytris immarginatis: punctis duobus flauis. *immarginatus.*

*Eumorphus Sumatrae.* Web. Inſ. 59. 1.

*Erotylus quadriguttatus.* Illig. Widem. Arch. 1. 2. 124. 18. tab. 1. fig. 4.

Habitat in Sumatra. D. Daldorff.

Statura omnino ſequentis, at elytra haud marginata nec violacea.

*Os* maxillis palpisque. *Labrum* porrectum, corneum, quadratum, vix emarginatum. *Palpi* quatuor inaequales, *anteriores* longiores, extrorſum vix craſſiores, quadriarticulati: articulo primo breuiſſimo, orbiculato, ſecundo longiori, obconico, baſi tenuiſſimo, tertio quartoque aequalibus, breuibus, ſubcraſſioribus, obtuſis, adhaerentes maxillae dorſo, *poſteriores* breues, capitati, triarticulati: articulo primo ſecundoque breuiſſimis, rotundatis, tertio incraſſato,

---

Eumorphi corpus oblongum, glabrum, ſubmarginatum, tardum; *capite* paruo, inſerto, thorace multo anguſtiore; *oculis* magnis, oblongis, transuerſis, marginalibus, thoracis angulo antico, ſubtectis; *antennis* thorace longioribus, ante oculos inſertis; *thorace* plano, ſubquadrato, angulis anticis poſticisque prominulis; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* tenuibus, longiuſculis, curſoriis; *tarſis* triarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

craſſato, obtuſo, truncato. adnati ligulae baſi exteriori. *Mandibula* craſſa, cornea, compreſſa, edentula, apice acutiſſima. *Maxilla* breuis, membranacea, bifida: laciniis aequalibus, rotundatis. *Ligula* breuis, membranacea, late emarginata: laciniis diſtantibus, tenuibus, acutiuſculis. *Labium* breue, corneum, transuerſum, truncatum.

*marginatus* 2. E. ater, elytris marginatis, violaceis: punctis duobus flauis.

Habitat in noua Cambria. Muſ. D. Billardiere.

Praecedente paullo maior. Antennae porrectae, nigrae, claua perfoliata, compreſſa. Thorax planus, marginatus, ater, nitidus. Elytra margine dilatato, plano, futura in medio eleuata, gibba, violaceo-nitida, punctis duobus magnis, flauis. Corpus atrum.

## 90. CNODVLON.

*Palpi* quatuor inaequales, *anteriores* ſecuriformes; *poſteriores* breues, capitati.

*Ligula* apice rotundata, integra.

*Antennae* filiformes.

*cupreum.* 1. C. atrum, thorace elytrisque cupreis.

*Ero-*

---

Cnoduli corpus oblongum, glabrum, immarginatum, tardum; *capite* paruo, ouato, inſerto; *oculis* magnis, approximatis, oblongis, transuerſalibus, marginalibus, bamo medio capitis firmatis; *antennis* diſtantibus, thorace longioribus, cantho oculorum inſertis; *thorace* breui, gibbo; *ſcutello* paruo, rotundato; *elytris* fornicatis, rigidis, longitudine abdominis; *pedibus* longiuſculis, tenuibus, curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Erotylus cupreus.* Ent. ſyſt. 2. 40. 21. *
Habitat in noua Hollandia. Muſ. D. Banks.

*Os* maxillis palpisque. *Palpi* quatuor inaequales, *anteriores* longiores, ſecuriformes, quadriarticulati: articulo primo breui, ſecundo longiori, obconico, tertio breui, quarto maiori, altero latere dilatato, adhaerentes maxillae dorſo, *poſteriores* breuiores, triarticulati: articulo primo ſecundoque breuibus, rotundatis, aequalibus, tertio craſſiori, clauato, compreſſo, adhaerentes ligulae medio. *Mandibula* craſſa, difformis, incurua, fornicata, intus repanda. *Maxilla* magna, compreſſa, medio vnidentata, apice rotundata. *Ligula* porrecta, cylindrica, membranacea, apice rotundata, integra. *Labium* breue, corneum, transuerſum.

2. C. atrum, thorace elytrisque ferrugineo variis. *nebuloſum.*
*Erotylus nebuloſus.* Ent. ſyſt. 2. 40. 22. *
Habitat in America meridionali. Muſ. Dr. Hunter.

3. C. atrum, thorace elytrisque ferrugineis. *dilatatum.*
*Erotylus dilatatus.* Ent. ſyſt. 2. 40. 23. *
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

4. C. nigrum, elytris cyaneis. *triſte.*
*Erotylus triſtis.* Ent. ſyſt. ſuppl. 102. 23. *
Habitat - -

5. C. atrum, elytris ſtriatis viridibus. *ſmaragdulum.*
*Erotylus ſmaragdulus.* Ent. ſyſt. 2. 40. 24. *
Habitat in noua Hollandia. Muſ. D. Banks.

6. C. atrum, thorace elytrisque cyaneis. *amethyſtinum.*
*Erotylus amethyſtinus.* Ent. ſyſt. 2. 40. 25. *
*Cnodulon.* Latreille Inſ. 54. 26.
Habitat in noua Hollandia. Muſ. D. Banks.

7. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bicolor.* 7. C. atrum, ſupra aeneum.
*Erotylus bicolor.* Ent. ſyſt. 2. 40. 26. *
Habitat in noua Hollandia. Muſ. D. Banks.
Caput aeneum, antennis nigris. Elytra punctat
ſtriata.

## 91. CEBRIO. *Palpi* quatuor filiformes.
*Maxilla* membranacea, vix vn
dentata.
*Ligula* apice palpigera.
*Antennae* filiformes.

*gigas.* 1. C. villoſus, fuſcus, elytris, abdomine femoribu
que teſtaceis. Ent. ſyſt. 2. 41. 1.
*Cebrio gigas.* Oliv. Inſ. 31. tab. 1. fig. 1.
I.37 Roſſ. Fn. Etr. 1. 100. 256. tab. 7. fig. 9.
Panz. Fn. Germ. 5. tab. 10.
Habitat in Gallia auſtraliori. D. Brouſſonet.

*fuſcus.* 2. C. fuſcus, elytris ſtriatis. Ent. ſyſt. 2. 41. 2. *
Habitat ad Cap. Bon. Spei. Dom. de Paykull.

*bicolor.* 3. C. ſupra griſeus, ſubtus ferrugineus.
Habitat in Carolina. Muſ. D. Boſc.
Statura praecedentium, at paullo minor. Capu
nigrum, antennis palpisque ferrugineis. Tho
rax parum tomentoſus, ſubpunctatus, poſtic
vtrin

---

Cebrionis corpus elongatum, immarginatum, pubeſcens, agi
le; *capite* ouato, thorace ſublatiore, mandibulis extertis
inſerto; *oculis* magnis, rotundatis, prominulis, margina
libus; *antennis* longitudine fere corporis, ante oculos in
ſertis; *thorace* transuerſo, poſtice latiore, angulo vtrinqu
porrecto, acuto; *ſcutello* paruo, rotundato; *elytris* rigidis
fornicatis, longitudine abdominis; *pedibus* validis; *femo
ribus* craſſiuſculis, compreſſis; *tarſis* anticis quinque-po
ſticis quadriarticulatis; *colore* obſcuro.

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google
Digitized by Google
Original from NEW YORK PUBLIC LIBRARY

vtrinque dente prominulo. Elytra ſtriata, fuſca. Corpus cum pedibus ferrugineum.

4 C. fuſcus, thoràce femoribusque rufis, elytris teſtaceis. Ent. ſyſt. ſuppl. 102. 3. * *ruficollis*
Habitat in Marocco. D. Schousboe. Muſ. D. de Seheſtedt.

92. ATOPA. *Palpi* aequales, filiformes.
*Maxilla* vnidentata.
*Ligula* membranacea, bifida: laciniis linearibus, diſtantibus.
*Antennae* filiformes.

1. A. liuida, pedibus fuſcis. *ceruina.*
*Ciſtela ceruina.* Ent. ſyſt. 2. 42. 1.
*Chryſomela ceruina.* Linn. ſyſt. Nat. 2. 602. 115. Fn. Sv. 575.
*Atopa ceruina.* Payk. Fn. Sv. 2. 116. 1.
*Daſcillus.* Latreille Inſ. 56. 24.
Degeer Inſ. 4. 235. tab. 9. fig. 8.
Oliv. Inſ. 54. tab. 1. fig. 2.
Habitat in Europae plantis.

2. A. liuida, elytris pedibusque fuſcis. *cinerea.*
*Ciſtela cinerea.* Ent. ſyſt. 2. 42. 2.
Habitat in Germania et Italia. *Variet. praecedentis*

Vix

---

Atopae corpus oblongum, immarginatum, pubeſcens, agile; *capite* paruo, ouato, inſerto; *oculis* rotundatis, vix prominulis, marginalibus; *antennis* thorace longioribus, ante oculos inſertis; *thorace* planiuſculo, ſubmarginato, angulo poſtico haud prominulo; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* mediocribus, compreſſis; *tarſis* quinquearticulatis; *colore* cinereo, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Vix a praecedente ſatis diſtincta.

*Os* maxillis palpisque. *Palpi* quatuor aequale filiformes: *anteriores* quadriarticulati: art culis ſubaequalibus; vltimo obtuſo, adhaeren tes maxillae dorſo, *poſteriores* triarticulati articulis aequalibus, cylindricis, adnati ligula baſi ſuperiori. *Mandibula* cornea, arcuata ſubintegra, apice obtuſa. *Maxilla* porrecta membranacea, vnidentata. *Ligula* porrecta membranacea, cylindrica, apice bifida: lacinii aequalibus, linearibus, diſtantibus. *Labium* breue, corneum, transuerſum, immarginatum

*liuida.* 3. A. liuida, antennis fuſcis.
Habitat in Terra del Fuego. Muſ. D. Banks.

## 93. CISTELA. *Palpi* inaequales, filiformes. *Maxilla* bifida. *Ligula* membranacea, rotun data, integra. *Antennae* filiformes.

*ceramboides.* 1. C. atra, thorace antice anguſtato, elytris ſtriati teſtaceis. Ent. ſyſt. 2. 42. 4.
*Chryſomela ceramboides.* Linn. ſyſt. Nat. 2. 602 117. Fn. Sv. 576.
*Ciſtela ceramboides.* Payk. Fn. Sv. 2. 121. 1. Oliv

---

Ciſtelae corpus oblongum, immarginatum, glabrum, agile *capite* ouato, exſerto; *oculis* magnis, oblongis, transuer ſis, reniformibus, marginalibus; *antennis* thorace longio ribus, oculis ipſis inſertis; *thorace* poſtice latiore, angul poſtico parum prominulo; *ſcutello* paruo aucto; *elytri* rigidis, fornicatis, longitudine abdominis; *pedibus* me diocribus, validis, curſoriis; *tarſis* anticis quinque poſti cis quadriarticulatis.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 54. tab. 1. fig. 4.
Herbſt. Arch. tab. 23. fig. 27.
Geoff. Inſ 1. 354. 1.
Habitat in Europa boreali.
Os maxillis palpisque. *Palpi* quatuor inaequales, ſubfiliformes, *anteriores* longiores, quadriarticulati: articulo primo breuiſſimo, ſecundo longiori, obconico, tertio breui, quarto paullo longiori, cylindrico, obtuſo, adhaerentes maxillae dorſo: *poſteriores* breues, triarticulati: articulis ſubaequalibus; vltimo obtuſo, adnati ligulae medio. *Mandibula* breuis, cornea, arcuata, edentula, acuta. *Maxilla* membranacea, bifida, lacinia exteriore paullo longiore, rotundata. *Ligula* breuis, membranacea, rotundata, integra. *Labium* breue, corneum, transuerſum, ſubemarginatum.

C. atra, thorace quadrato, elytris ſtriatis, teſtaceis. *lepturoides.* Ent. ſyſt. 2. 43. 5. *
*Ciſtela ruſtarſis.* Lesk. Iter. 1. 15. 5. tab. A. fig. 4.
Oliv. Inſ. 54. tab 1. fig. 3. I. 33.
Panz. Fn. Germ. 5. tab. 11.
Habitat in Europae auſtralioris ſpicis ſecalinis. Muſ. D. de Seheſtedt.

C. nigra, thorace, elytris abdomineque teſtaceis. *teſtacea.* Ent. ſyſt. 2. 43. 6. *
Habitat in Chryſanthemo, Sambuco aliisque Barbariae plantis. D. Vahl. Hungaria, Lusitania nec non Sueciae

C. nigra, thorace quadrato, elytris teſtaceis, antennarum baſi tibiisque piceis. Ent. ſyſt. 2. 43. 7. * *picipes.*
Habitat in Daniae plantis. Muſ. D. de Seheſtedt.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bicolor.* 5. C. nigra, elytris pedibusque ſulphureis. Ent. ſyſt. 4. 447. 8. *

Panz. Fn. Germ. 34. tab. 8.

Habitat in Germania. D. Schneider.

*ſulphurea.* 6. C. flaua, elytris ſulphureis. Ent. ſyſt. 2. 43. 8. *

*Chryſomela ſulphurea.* Linn. ſyſt. Nat. 2. 602. 114.

*Ciſtela ſulphurea.* Payk. Fn. Sv. 2. 125. 7.

*Tenebrio lutea.* Geoff. Inſ. 1. 351. 11.

Oliv. Inſ. 54. tab. 1. fig. 6.

Herbſt. Arch. tab. 23. fig. 28.

Habitat in Europae floribus vmbellatis.

*ruficollis.* 7. C. atra, thorace ferrugineo, elytris ſtriatis. Ent. ſyſt. 2. 43. 9. *

Oliv. Inſ. 54. tab. 1. fig. 5.

Habitat in Europae auſtralioris Santolina.

*coerulea.* 8. C. atra, elytris ſtriatis, coeruleſcentibus. Ent. ſyſt. 2. 43. 10. *

Oliv. Inſ. 54. tab 2. fig. 18.

Habitat in Barbariae plantis. D. Prof. Vahl.

*nigripennis* 9. C. ferruginea, capite elytrisque ſtriatis nigris. Ent. ſyſt. 2. 44. 11. *

Habitat in Gallia meridionali.

*analis.* 10. C. rufeſcens, antennis, macula, elytrorum anoque nigris. Ent ſyſt. 2. 44. 12. *

Habitat Tranquebariae. D. Koenig.

*melanocephala.* 11. C. teſtacea, capite anoque nigris.

Habitat Tranquebariae. Muſ. D. Lund.

Statura et ſumma affinitas C. analis, at diſtincta videtur, paullo minor. Caput nigrum. Thorax, elytra, corpus teſtacea, immaculata, ano ſolo nigro. Pedes teſtacei.

12. C.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

12. C. nigra, elytris laeuibus, antennis pedibusque ferrugineis. Ent. syst. 2. 44. 13. * *rufipes.*
Habitat in Germania. D. Smidt.

13. C. nigra, elytris striatis, pedibus ferrugineis. Ent. syst. 2. 44. 14. * *fuluipes.*
*Cistela fuluipes.* Payk. Fn. Sv. 2. 122. 2.
Habitat in Germania. D. Smidt.

14. C. atra, immaculata. *Morio.*
*Erotylus Morio.* Ent. syst. 2. 39. 19. *
Habitat in noua Hollandia. Mus. D. Banks.

15. C. nigra, elytris striatis pedibusque flauis. Ent. syst. 2. 44. 16. * *murina.*
*Cistela murina.* Payk. Fn. Sv. 2. 126. 8.
*Chrysomela murina.* Linn. syst. Nat. 2. 602. 118. Fn. Sv. 577.
*Mordella.* Geoff. Inf. 1. 355. 4.
Oliv. Inf. 54. tab. 1. fig. 13.
~~Herbst Arch. tab. 23. fig. 29.~~
Habitat in Europae plantis.

16. C. fusca, thorace pedibusque ferrugineis, elytris laeuibus. Ent. syst. 2. 45. 18. * *thoracica.*
Habitat Halae Saxonum. Dom. Hybner.

17. C. atra, elytris nigris: macula baseos flaua, abdominis basi pedibusque flauis. Ent. syst. 2. 45. 19. * *flauipes.*
*Cistela flauipes.* Payk. Fn. Sv. 2. 125. 6.
Habitat in Europae plantis. Mus. D. Banks.

18. C. nigra, elytris substriatis: puncto humerali tibiisque flauis. Ent. syst. 2. 45. 21. * *humeralis.*
*Cistela bipustulata.* Payk. Fn. Sv. 2. 124. 5.
~~*Cistela axillaris.* Payk. Fn. Sv. 2. 123. 4.~~
Habitat in Europa boreali.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

C. flauipede minor. Antennae nigrae, basi flauae Elytra substriata.

*Euonymi.* 19. C. testacea, abdomine griseo, elytris laeuibus Ent. syst. 2. 45. 20. *

III. 36 Panz. Fn. Germ. 34. tab. 7.

Habitat in Germaniae Euonymo D. Schneider.

*nigrita.* 20. C. atra, elytris substriatis, antennis pedibusqu concoloribus. Ent. syst. 4. App. 447. 21. *

Habitat in Germania. D. Schneider.

*maura.* 21. C. atra, elytris substriatis, antennarum basi pedi busque ferrugineis. Ent. syst. 2. 45. 22. *

II. 9. III. 58 Habitat in Germania. D. Smidt.

*pallipes.* 22. C. nigra, nitidula, antennarum basi pedibusqu pallidis. Ent. syst. 2. 46. 23. *

Habitat in Germania. D. Smidt.

*angustata.* 23. C. thorace elytrisque obscure rufis, medio nigri Ent. syst. 2. 46. 25. *

Habitat in Anglia. Mus. D. Banks.

*festiua.* 24. C. ferruginea, elytris cyaneo-viridibus: mar gine ferrugineo. Ent. syst. 2. 47. 28. *

Habitat ad Cap. Bon. Spei. Mus. D. Banks.

*aulica.* 25. C. atra, thorace rufo, elytris cyaneis. Ent. syst 2. 47. 29. *

Habitat ad Cap. Bon. Spei. Mus. D. Banks.

*hirta.* 26. C. hirta, capite thoraceque rufis, elytris cyaneis Ent. syst. 2. 47. 31. *

Habitat ad Cap. Bon. Spei. Mus. D. Banks.

De genere haereo, species mihi haud rite nota.

*agilis.* 27. C. atra, elytris pedibusque ferrugineis.

Habitat in Germania.

Parua. Caput nigrum, antennis palpisque ferru gineis. Thorax laeuis, niger, margine ferrugi

nec

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

neo. Elytra vix ſtriata, ferruginea. Corpus nigrum, pedibus ferrugineis.

28. C. ferruginea, elytris flauis: lineolis tribus nigris. Ent. ſyſt. 2. 47. 33. * *6lineata.*
Habitat in - - Muſ. D. Banks.

94. ALLECVLA. *Palpi* quatuor, inaequales, *anteriores* ſecuriformes; *poſteriores* clauati.
*Labium* emarginatum.
*Antennae* ſetaceae.

1. A. nigra, pedibus piceis. *Morio.*
*Erotylus rufipes.* Ent. ſyſt. 2. 40. 27. *
*Ciſtela Morio.* Ent. ſyſt. 2. 46. 24. *
*Ciſtela Morio.* Payk. Fn. Sv. 2. 123. 3. III. 38
Oliv. Inſ. 54. tab. 1. fig. 7. VIII. 93.
Habitat Kiliae.
*Os* maxillis palpisque. *Palpi* quatuor inaequales, *anteriores* longiores, ſecuriformes, quadriarticulati: articulo tertio breuiſſimo, vltimo craſſiori, altero latere dilatato, adhaerentes maxillae dorſo, *poſteriores* breuiſſimi, ſubclauati: articulo ſecundo breuiſſimo, vltimo craſſiori, truncato, inſerti ligulae medio laterali. *Mandibula* cornea, concaua, acuta, edentula. *Maxilla*

---

Alleculae corpus oblongum, glabrum, immarginatum, agile; *capite* ouato, exſerto; *oculis* magnis, oblongis, transuerſis, marginalibus; *antennis* longitudine fere corporis, diſtantibus, ante oculos inſertis; *thorace* antice anguſtiore, margine laterali rotundato, poſtico ſinuato; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, tenuibus, curſoriis; *tarſis* quadriarticulatis; *colore* nigro, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*xilla* cornea, bifida: lacinia exteriore, maiore. clauata, concaua, rotundata. *Ligula* - - - - *Labium* breue, corneum, basi cylindricum, apice dilatatum, truncatum, subemarginatum.

*varians.* 2. A. grisea, oculis nigris, elytris subtriatis.
*Cistela varians.* Ent. syst. 2. 44. 15. *
Oliv. Inf. 54. tab 2. fig. 17.
Habitat Halae Saxonum. Dom. Hybner.

*pallicornis.* 3. A. fusca, elytris striatis ferrugineis, antennis pedibusque testaceis.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura et magnitudo praecedentis. Caput fuscum, ore antennisque testaceis. Elytra striata, ferruginea. Corpus fuscum, pedibus flauis.

*ferruginea.* 4. A. testacea, capite thoraceque fuscis, elytris striatis.
*Cistela ferruginea.* Ent. syst. 2. 45. 17. *
Habitat in Germania. D. Smidt.

*fuscata.* 5. A. nigra, elytris striatis, antennis pedibusque testaceis.
*Cistela fuscata.* Ent. syst. suppl. 102. 22. *
Habitat Kiliae.

## 95. ZYGIA.

*Palpi* inaequales, filiformes.
*Maxilla* vnidentata.
*Ligula* elongata, membranacea.
*Antennae* moniliformes.

*oblonga.* 1. ZYGIA.
*Zygia oblonga.* Ent. syst. 2. 48. 1.
Habitat in Oriente. D. Prof. Forskahl.

96. ZO-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

16. ZONITIS. *Palpi* filiformes.
*Maxilla* palpis longior, integra.
*Antennae* ſetaceae.

1. Z. thorace mutico, pallide teſtacea, antennis pedumque apicibus nigris. Ent. ſyſt. 4. App. 447. 1. * ***pallida.***
Habitat in Americae meridionalis Inſulis. Dom. v. Rohr.

2. Z. thorace mutico, teſtacea, antennis elytrorumque apicibus nigris. Ent. ſyſt. 2. 48. 1. * ***praeuſta.***
Panz. Fn. Germ. 36. tab. 6. 7.
Habitat in Italia. Dr. Allioni.

3. Z. thorace mutico, ruſo, elytris pectoreque nigris. Ent. ſyſt. ſuppl. 103. 2. * ***nigripennis***
Habitat in Marocco. Muſ. D. Lund.

4. Z. thorace angulato, ſpinoſo, teſtacea, elytris apice cyaneis, antennis pedibusque nigris. Ent. ſyſt. 2. 49. 2. * ***angulata.***
Hybn. Naturf. 24. 45. 11. tab. 2. fig. 12.
Habitat in Inſula Amſterdam. Muſ. D. Banks.

5. Z. thorace mutico, nigra, thorace elytrisque teſtaceis. Ent. ſyſt. 2. 49. 3. * ***mutica.***
*Zonitis mutica* Hybn. Naturf. 24. 44. 10. tab. 2. fig. 11.
Habitat in Hungaria. Dom. Hybner.

6. Z.

---

Zonitidis corpus oblongum, glabrum, immarginatum, agile; *capite* ouato, inflexo, exſerto; *oculis* magnis, reniformibus, transuerſis, marginalibus, vix prominulis; *antennis* diſtantibus, ſub oculis inſertis, longitudine corporis; *thorace* ſubquadrato, plano, antice parum anguſtiore; *ſcutello* paruo, triangulari; *elytris* molliuſculis, fornicatis, longitudine abdominis; *pedibus* tenuibus, longiuſculis; *tarſis* anticis quinque - poſticis quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fuluipennis.* 6. Z. thorace mutico, atra, elytris teſtaceis. Ent. ſyſt. 2. 49. 4. *
Habitat in Hungaria

*chryſomelina.* 7. Z. flaua, elytris puncto medio apiceque nigris. Ent. ſyſt. 2. 49. 5. *
Habitat in Oriente. D. Prof. Forſkahl.

*flaua.* 8. Z. ruſeſcens, elytris flauis apice nigris. Ent. ſyſt. 2. 49. 6.
Habitat in Oriente. D. Prof. Forſkahl.

*viridipennis.* 9. Z. thorace rufo, elytris viridibus. Ent. ſyſt. ſuppl. 103. 7. *
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

*roſtrata.* 10. Z. teſtacea, pectore elytrorumque punctis tribus nigris. Ent. ſyſt. 2. 50. 7. *
Habitat in Barbaria. Muſ. Dom. Desfontaines.

*vittata.* 11. Z. ferruginea, elytris teſtaceis: vitta lata atra. Ent. ſyſt. ſuppl. 104. 7. *
Habitat in Carolina. Muſ. D. Boſc.

## 97. APALVS.

*Palpi* aequales, filiformes.
*Maxilla* cornea, vnidentata.
*Ligula* membranacea, truncata, integra.
*Antennae* filiformes.

*amaculatus* 1. A. niger, elytris teſtaceis: puncto nigro. Ent. ſyſt. 2. 50. 1.
*Meloë bimaculatus.* Linn. ſyſt. Nat. 2. 680. 9. Fn. Sv. p. 228. n. 828. Vdm. Diſſ. 82. Oliv.

IX.

---

Apali corpus oblongum, glabrum, immarginatum, agile; *capite* ouato, exſerto; *oculis* oblongis, transuerſis, marginalibus; *antennis* thorace longioribus, ante oculos inter tis

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 52. tab. 1. fig. 2.
Degeer Inf. 5. tab. 1. fig. 18.
Payk. Fn. Sv. 2. 127. 1.
Habitat primo vere in Europae borealis arenofis.

2. A. rufus, capite elytrorumque maculis duabus nigris. Ent. fyſt. 2. 50. 2. * *4maculatus*
Habitat in America boreali.

98. ALVRNVS. *Palpi* ſex breuiſſimi.
*Maxilla* cornea, forricata.
*Antennae* cylindricae.

1. A. thorace coccineo, elytris flauis. Ent. fyſt. 2. 51. 1. * *groſſus.*
Voet. Col. 2. tab. 29. fig. 9.
Habitat in Cayenna.
Antennae nigrae, corpore breuiores.

2. A. teſtaceus, antennis elytrisque ante apicem nigris. *teſtaceus.*
Habitat in noua Cambria. Muſ. D. Billardiere.

Huius

---

tis, diſtantibus; *thorace* rotundato, anguſtiore, linea media impreſſa; *ſcutello* breui, rotundato; *elytris* molliuſculis, incumbentibus, longitudine abdominis; *pedibus* tenuibus, longiuſculis, curſoriis; *tarſis* anticis quinque-poricis quadriarticulatis; *colore* nigro flauoque, obſcuro.

---

Alurni corpus magnum, oblongum, depreſſum, glabrum, immarginatum, tardum; *capite* decliui, rotundato, inſerto; *oculis* magnis, ſubrotundatis, marginalibus; *antennis* thorace longioribus, approximatis, inter oculos inſertis; *thorace* rotundato, plano, vix marginato; *ſcutello* maiuſculo, ouato, obtuſo; *elytris* rigidis, baſi gibbere notatis, fornicatis, abdomine latioribus longioribusque; *pedibus* validis, longiuſculis, curſoriis; *tarſis* triarticulatis: articulis latis, cordatis, ſubtus valde villoſis; *colore* vario, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Huius generis videtur, praecedente triplo minor. Antennae filiformes, corpore breuiores, articulis breuibus, cylindricis. Caput et thorax laeuia, teſtacea, immaculata. Elytra punctato-ſtriata, teſtacea, ante apicem late atra. Corpus et pedes teſtacea.

*ferrugineus* 3. A. teſtaceus, thorace punctato, elytris ſtriato-punctatis.

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Statura omnino A. groſſi, at duplo fere minor. Antennae cylindricae, fuſcae: articulo primo breui, craſſiori, rotundato. Caput et thorax punctata, obſcure teſtacea, immaculata. Elytra obſcure teſtacea, ſtriata et inter ſtrias linea geminata e punctis impreſſis. Corpus et pedes ferruginea.

## 99. SAGRA. *Palpi* inaequales, filiformes. *Maxilla* cornea, vnidentata. *Labium* breue, corneum, rotundatum, ciliatum, fiſſum.

*femorata.* 1. S. viridi-aenea, femoribus tibiisque poſticis dentatis. Ent. ſyſt. 2. 51. 1. * Web. Inſ. 60. 1.

*Tenebrio femoratus.* Drury Inſ. 2. tab. 34. f. 5.

*Tenebrio viridis.* Sulz. Hiſt. Inſ. tab. 7. fig. 8.

Habitat in India. 2. S.

---

Sagrae corpus magnum, craſſum, ouatum, glabrum, immarginatum, tardum; *capite* magno, cylindrico, inſerto; *oculis* oblongis, transuerſis, medio margini infertis; *antennis* diſtantibus, corpore breuioribus, ante oculos inſertis; *thorace* rotundato, anguſto, ſubcylindrico, immarginato; *ſcutello* minuto, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* curſoriis; *femoribus* poſticis incraſſatis, dentatis; *tibiis* incuruis, ſaepius dentatis; *tarſis* triarticulatis: articulis cordatis; *colore* nitido.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

2. S. cyaneo-purpurea, femoribus posticis dentatis, tibiis apice sinuatis. *splendida.*

*Sagra splendida.* Web. Inf. 61. 2.

Habitat in China.

Statura omnino praecedentis, at color omnino distinctus, totus purpureus, cyaneo-nitidus. Antennae nigrae. Tibiae apice sinu magno excauatae

3. S. purpurea, nitida, femoribus posticis vnidentatis, tibiis integris. *purpurea.*

*Sagra purpurea.* Web. Inf. 61. 3.

Habitat in China.

Diu marem praecedentis credidi, qua duplo fere minor. Differt tamen femoribus posticis tantum vnidentatis tibiisque haud sinuatis.

4. S. cyanea, elytris aureis, tibiis posticis medio vnidentatis. *fulgida.*

*Sagra fulgida.* Web. Inf. 62. 5.

Habitat in China. D. Weber.

Statura omnino praecedentium. Corpus, caput, thorax cyaneo-nitida, immaculata. Tibiae quatuor anticae incuruae. Femora postica incrassata, tridentata: dente medio maiori, tibiae incuruae, ante apicem vnidentatae.

5. S. obscure viridis, femoribus posticis bidentatis, tibiisque vnidentatis. *tristis.*

Habitat in Cayenna.

Statura et summa affinitas praecedentium. Antennae corpore breuiores, cyaneae. Caput, thorax, elytra obscure viridia, minime nitida. Abdomen subtus basi macula media villosa, aurea. Femora postica incrassata, basi interiori macula magna aurea, apice bidentata: dente posteriori longiore. Tibiae incuruae, apice vnidentatae.

6. S.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*dentipes.* 6. S. nigra, femoribus tibiisque poſticis dentati
Ent. ſyſt. 2. 52. 2. *
*Sagra dentipes.* Web. Inſ. 63. 6.
Habitat ad Cap. Bon. Spei. Muſ. D. Schulz.
Tibiae poſticae apice ſinuatae.
Species Sagrae omnes valde affines et forte mera varietates, quamuis colore pedumque dentibu differunt.

## 100. CLYTHRA. *Palpi* quatuor inaequales filiformes.
*Ligula* breuis, membra nacea, emarginata.
*Antennae* breues, ſerratae

*longipes.* 1. C. nigra, obſcura, elytris pallidis: maculis tri bus nigris, pedibus anticis elongatis. Ent. ſyſt ſuppl. 109. 1.
*Clythra longipes.* Laichart. 1. 166. 1.
*Cryptocephalus longipes.* Ent. ſyſt. 2. 52. 1.
Schaeff. Icon. tab. 6. fig. 3.
Habitat in Germaniae Corylo.

*3punctata.* 2. C. nigro-cyanea, elytris teſtaceis: punctis tri bus nigris, pedibus anticis elongatis. Ent. ſyſt ſuppl. 110. 2.

*Cry*

---

Clythrae corpus paruum, immarginatum, cylindricum, ſa pius glabrum, tardum; *capite* paruo, retracto, inſerto, ob tuſo; *oculis* oblongis, marginalibus, vix prominulis; *an tennis* thoracis longitudine, ſerratis, diſtantibus, ante oc los inſertis; *thorace* transuerſo, rotundato: margine d flexo; *ſcutello* minuto, rotundato; *elytris* rigidis, forn catis, abdominis longitudine; *pedibus* validis, anticis mi ſculorum ſaepius elongatis; *tarſis* quadriarticulatis: articul penultimo bifido; *colore* vario.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cryptocephalus 3punctatus.* Ent. ſyſt. 2. 53. 2. ♀.

Habitat in Italia. Dr. Allioni.

3. C. ferruginea, thorace macula, elytris flaueſcentibus: faſcia apiceque atris. ***curuata.***

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Antennae breues, craſſae, atrae. Caput flauum, macula inter oculos transuerſa, atra. Thorax laeuis, rufeſcens, macula magna, atra. Elytra punctata, flaueſcentia, faſcia extus ad baſin dilatata, apiceque late atris. Corpus ferrugineum: tibiis incuruis nigris.

4. C. flaueſcens, thoracis macula elytrisque late atris. ***curuipes.***

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Media. Antennae nigrae. Caput flauum, vertice nigro. Thorax laeuis, flauus: macula magna dorſali nigra. Elytra punctata, baſi oblique flaua, apice vltra medium atra. Corpus flauum, femoribus incraſſatis, compreſſis, tibiisque incuruis, nigris.

5. C. flaueſcens, thoracis macula, elytris macula baſeos faſciaque atris. ***tibialis.***

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et ſumma affinitas praecedentis. Differt tantum elytris macula magna baſeos, faſciaque media, lata, atris.

6. C. flaueſcens, thorace punctis quatuor, elytris faſciis quatuor nigris: anticis duabus obliquis. ***obliqua.***

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Sta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura praecedentis, paullo minor. Antennae nigrae, ferraturis flauis. Caput flauum, puncto verticali atro. Thorax obſcure flaueſcens, puntis quatuor nigris. Elytra punctata, flaua: faſciis duabus, baſeos obliquis duabusque pone medium nigris. Corpus flaueſcens.

*palliata.* 7. C. atra, elytris dilatatis, teſtaceis: punctis duobus faſciaque media atris.

Habitat in India orientali. D. Daldorff. Muſ. D. Lund.

Magna. Antennae valde ſerratae, nigrae, baſi ferrugineae. Caput atrum. Thorax ater, margine laterali interdum rufo. Elytra laeuia, teſtacea, puncto baſeos, faſcia media punctoque ante apicem atris. Elytra valde dilatata, abdomine multo latiora, margine parum angulato. Corpus atrum.

*hirta.* 8. C. villoſa, cyanea, elytris rufis: punctis tribus nigris.

Habitat in Marocco. Muſ. D. de Seheſtedt.

Paullo maior C. tripunctata, ei affinis, at magis hirta et elytra rufa. Puncta elytrorum 1. 2.

*faſciata.* 9. C. thorace ferrugineo. nigro, bipunctato, elytris teſtaceis: macula baſeos faſcia margineque poſtico atris.

Habitat in Guinea. Muſ. D. Lund.

Magnitudo et ſtatura omnino C. tripunctatae. Caput nigrum. Antennae breues, ſerratae, nigrae, baſi rufae. Thorax glaber, rufus, nitidus: punctis duobus nigris. Elytra laeuia, teſtacea: macula magna baſeos, faſcia pone medium lata, margineque poſtico atris. Corpus nigrum, argenteo-tomentoſum.

10. C.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

10. C nigra, thorace rubro immaculato, elytris rubris: punctis tribus nigris. Ent. syst. suppl. 110. 3. *6 maculata*

*Cryptocephalus 6maculatus.* Ent. syst. 2. 53. 5.*

Habitat in Italia. Dr. Allioni.

11. C. flaua, thoracis basi et elytrorum fasciis tribus atris. *argentata*

Habitat in Guinea. Mus. D. de Seheſtedt.

Magna. Caput nigrum, albo-villosum. Thorax niger, margine antico et laterali lineaque media flauis. Elytra flaua, fasciis tribus latis, subdentatis, obliquis atris: tertia terminali. Macula vna alteraue coccinea, marginalis in colore flauo. Corpus argenteo-villosum.

12. C. cinereo-fericea, thorace rubro immaculato, elytris rubris: punctis tribus nigris; medio maiori transuerso. *6notata.*

Habitat in noua Cambria. D. Billardiere.

C. 6maculata duplo minor. Antennae breues, serratae, nigrae, basi ferrugineae. Caput nigrum, orbita oculorum cinereo-fericea. Thorax rufus, immaculatus. Elytra laeuia, rubra, puncto paruo nigro, ad marginem exteriorem baseos, maiori transuerso in medio et tertio ad futuram versus apicem. Corpus nigrum, cinereo-fericeum.

13. C. nigra, elytris rubris: punctis duobus nigris. Ent. syst. suppl. 110. 4. *4punctata.*

*Cryptocephalus 4punctatus.* Ent. syst. 2. 54. 6. *

*Chrysomela 4punctata.* Linn. syst. Nat. 2. 596. 76. Fn. Sv. 547.

Geoff. Inf. 1. 195. tab. 3. fig. 4.

Degeer Inf. 5. 392. 32. tab. 10. fig. 7.

Schaeff.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Schaeff. Icon. tab. 6. fig. 1. 2.
Payk. Fn. Sv. 2. 128. 1.
Habitat in Europae Corylo frequens.
Differt interdum puncto posteriori transuerso, fe fasciam constituente.
*Larua* saccata: sacco ouato scabro, antice obliq truncato. Schall. Act. Hall. 1.

*picta.* 14. C. flaua, thorace maculato, elytris vittis duab postice coëuntibus nigris.
Habitat in Guinea. Mus. D. Lund.
Magna. Antennae breues serratae, nigrae, b pallidae. Caput nigrum. Thorax glaber, ni dus, flauus, maculis duabus dorsalibus, margi postico lunato nigro connexis. Elytra nitid flaua, vittis duabus, quarum interior. bas haud attingit, pone medium fere in fasciam d latatis, coëuntibus, et postice in vnicum te minantibus. Corpus nigrum, cinereo-nitid lum, pedibus anticis elongatis.

*4notata.* 15. C. cyaneo-nigra, elytris rubris: maculis du bus cyaneis. Rub. Suppl. no 5
*Cryptocephalus 4notatus.* Ent. syst. 2. 54. 7.
Habitat in Barbaria. D. Prof. Vahl.

*3maculata.* 16. C. nigra, cinereo-villosa, elytris rufis: macul tribus cyaneis: posteriore maiore transuersi Ent. syst. suppl. 110. 6. *
Habitat in India orientali. D. Daldorff. Mus. I de Seheſtedt.

*rugosa.* 17. C. atra, elytris testaceis: fasciis duabus atri Ent. syst. suppl. 110. 6. *
Habitat ad Cap. Bon. Spei. Mus. D. Lund.

*Atraphaxidis.* 18. C. nigra, thorace rubro trimaculato, elytris te staceis: maculis tribus nigris, tibiis rufis. En syst. suppl. 110. 8.

Cr

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cryptocephalus Atraphaxidis.* Ent. ſyſt. 2. 54. 9. *

*Chryſomela Atraphaxidis.* Pall. Iter. 2. 725. 65.

Habitat in Sibiriae Atraphaxide. Muſ. D. Banks.

). C. viridi-cyanea, thorace el trisque eleuato-tuberculatis. Ent. ſyſt. ſuppl. 111. 9. * *monſtroſa.*

*Clythra monſtroſa.* Oliv. Inſ.

Habitat in Cayenna. Muſ. D. Boſc.

). C. obſcura, thorace elytrisque eleuato-tuberculatis. Ent. ſyſt. ſuppl. 111. 10. * *plicata.*

Habitat in Carolina. Muſ. D. Boſc.

:. C. nigra, thorace gibbo, elytris tuberculatis. Ent. ſyſt. ſuppl. 112. 11. * *gibber.*

Habitat in Carolina. Muſ. D. Boſc.

). C. atra, thorace elytrisque rufis nigro-punctatis. *impreſſa.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentium, at parua. Corpus atrum. Thorax ruber, maculis quatuor impreſſis, nigris; dorſalibus duabus maioribus, ſinuatis. Elytra inaequalia, rufa, punctis quatuor aut quinque nigris impreſſis. Anus obtuſus.

). C. atra, thorace gibbo, ſtriato-criſtato, elytris rugoſis, tuberculatis. *criſtata.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Minuta. Statura praecedentium tota atra, immaculata. Thorax ſtriatus, gibbus, in medio carinatus. Elytra rugoſa, tuberculis plurimis eleuatis.

). C. atra, nitida, elytris flauis: lunula, faſcia punctoque apicis nigris. Ent. ſyſt. ſuppl. 112. 12. *lunulata.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cryptocephalus lunulatus.* Ent. syst. 2. 55. 10.
Habitat - - Mus. D. Banks.

*12maculata* 25. C. nigra, thorace elytrisque rubris: punctis quatuor nigris. Ent. syst. suppl. 112. 13.
*Cryptocephalus 12maculatus.* Ent. syst. 2. 55. 11. *
Habitat ad Cap. Bon. Spei. Mus. D. Banks.

*obsita.* 26. C. testacea, antennis elytrorumque margine nigris. Ent. syst. suppl. 112. 14. *
*Cryptocephalus obsitus.* Ent. syst. 2. 55. 12.
Habitat in America. Dom. Drury.

*tridentata.* 27. C. coerulescens, elytris testaceis: puncto humerali nigro. Ent. syst. suppl. 112. 15.
*Cryptocephalus tridentatus.* Ent. syst. 2. 55. 14. *
*Chrysomela tridentata.* Linn. syst. Nat. 2. 595. 73. Fn. Sv. 546.
*N. 51.* Geoff. Inf. 1. 196. 3.
~~Degeer Inf. 5. 333. 36. tab. 10. fig. 19.~~
Schaeff. Icon. tab. 77. fig. 5.
Payk. Fn. Sv. 2. 129. 2.
Habitat in Lonicera Xylosteo.
Pedes anteriores marium elongati.

*dominicana* 28. C. nigra, elytris punctatis basi gibbere notatis.
Habitat in Carolina. Mus. D. Bosc.
Media. Corpus totum nigrum, obscure nitidum subtus cinereo-holosericeum. Elytra punctata gibbere ad marginem exteriorem notata.

*taxicornis.* 29. C. obscure cyanea, elytris testaceis immaculatis antennis elongatis serratis. Ent. syst. suppl. 112. 16.

C.

---

Clythrae genus Cryptocephalis valde affine. Characterem generis naturalem in Eoromologiae systematicae supplemento dedi.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cryptocephalus taxicornis.* Ent. ſyſt. 2. 56. 15. ♂
Payk. Fn. Sv. 130. 3.
Habitat in Italia. Dr. Allioni.

*decora.*

o. C. cyanea nitida, elytris rufis, immaculatis.
Habitat in India orientali. Muſ. D. Lund.
Statura C. taxicornis, at omnino diſtincta. Caput et thorax cyanea, nitida. Elytra ſubſtriata, rufa. Corpus cyaneum.

*bifaſciata.*

1. C. thorace teſtaceo, baſi cyaneo, elytris teſtaceis: puncto faſciaque media cyaneis.
*Clythra bifaſciata.* Illig. Wied. Arch. 1. 2. 130. 24.
Habitat in Sumatra. D. Daldorff.
Magnitudo et ſtatura C. taxicornis. Antennae breues, valde ſerratae, obſcurae. Caput cyaneum, nitidum. Thorax laeuis, teſtaceus, baſi cyaneus, nitidus, qui color marginem tamen haud attingit. Scutellum cyaneum, apice recuruum. Elytra laeuia, puncto paruo baſeos, quod interdum deeſt, faſciaque media, margines tamen haud attingente, cyaneis, nitidis. Corpus cinereo-pubeſcens. Pedes teſtacei, femoribus cyaneis.

*bisti-*
*notata.*

2. C. nigra, thorace rubro, immaculato, elytris maculis tribus cyaneis: poſtica maiori transuerſa.
Habitat in Tanger. D. Schousboe. Muſ. D. de Seheſtedt.
Media. Antennae breues, ſerratae Caput nigrum. Thorax laeuis, rufus, immaculatus. Elytra vix punctata, teſtacea, nitida, punctis tribus: 2 baſeos, tertio in medio, maiori transuerſo, fere quadrato. Corpus nigrum.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*8ſignata.* 33. C. atra, elytris ferrugineis: punctis quatuor nigris.

Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

Magna. Antennae breues, valde ſerratae. Caput et thorax laeuia, atra, nitida, immaculata. Elytra vix punctata, rufa, punctis quatuor nigris 2. 2. oblique poſitis. Corpus nigrum.

*8punctata.* 34. C. nigra, thorace rufo, elytris teſtaceis: punctis quatuor nigris. Ent. ſyſt. ſuppl. 112. 17.

*Cryptocephalus 8punctatus.* Ent. ſyſt. 2. 56. 18. *

~~Naturf. 24. 17. 22. tab. 1. fig. 22.~~

Habitat in Barbariae plantis. D. Prof. Vahl.

*maxilloſa.* 35. C. capite thoraceque fuluis, elytris flauis: puncto baſeos ſcutelloque nigris. Ent. ſyſt. ſuppl. 112. 18.

*Cryptocephalus maxilloſus.* Ent. ſyſt. 2. 56. 19. *

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*aurita.* 36. C. atra, thorace vtrinque macula lutea, tibiis flauis. Ent. ſyſt. ſuppl. 112. 19.

*Cryptocephalus auritus.* Ent. ſyſt. 2. 57. 20. *

*Chryſomela aurita.* Linn. ſyſt. Nat. 2. 256. 75.

Herbſt. Arch. tab. 44. fig. 5.

Habitat in Saxoniae Corylo.

*Lentiſci.* 37. C. coerulea, elytris ſanguineis: maculis coeruleis, cicatriſantibus. Ent. ſyſt. ſuppl. 113. 20.

*Cryptocephalus Lentiſci.* Ent. ſyſt. 2. 57. 21. *

*Chryſomela varioloſa.* Linn ſyſt. Nat. 2 591. 33.

Habitat in Africae Lentiſco. Muſ. D. Desfontaines.

*4maculata.* 38. C. rufa, capitis baſi elytrorumque maculis duabus cyaneis. Ent. ſyſt. ſuppl. 113. 21.

*Cryptocephalus 4maculatus.* Ent. ſyſt. 2. 57. 23. *

Chry-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Chryſomela 4 maculata.* Linn. ſyſt. Nat. 2. 596. 77.
Schaeff. Icon. tab. 6. fig. 67.
Habitat in Saxoniae Corylo.

39. C. nigra, thorace rufo: maculis duabus, elytris flauis, macula, faſcia media punctoque apicis nigris. *turcica.*
Habitat in India orientali. D. Daldorff. Muſ. D. de Seheſtedt.
Paullo maior C. longimana. Caput nigrum, antennis breuibus, ſerratis. Thorax rotundatus, rufus: lunula magna in ipſa baſi, punctisque duobus baſi connexis, nigris. Elytra punctata, teſtacea, macula magna baſeos, faſcia lata, media punctoque apicis nigris. Corpus nigrum, cinereo-holoſericeum.

40. C. obſcure aenea, elytris teſtaceis: puncto baſeos nigro. Ent. ſyſt. ſuppl. 113. 22. *longimana.*
*Cryptocephalus longimanus* Ent. ſyſt. 2. 58. 25.
*Chryſomela longimana.* Linn. ſyſt. Nat. 2. 599. 95.
Payk. Fn. Sv. 2. 131. 4. *IV. 51.*
Habitat in Sueciae Trifolio montano, Euphorbia cypariſſia.
Pedes in altero ſexu elongati.
*Larua* glabra, ſaccata: ſacco pyriformi, piloſo, fuſco.

41. C. rufa, elytris fuſcis, cyaneo-nitidulis, pedibus anticis elongatis. Ent. ſyſt. ſuppl. 113. 23. *Lar.*
*Cryptocephalus Lar.* Ent. ſyſt. 2. 58. 26. *
Habitat in America meridionali. Muſ. Dr. Hunter.

42. C. nigra, obſcura, elytris macula humerali rufa. *humeralis.*
Habitat in Carolina. Muſ. D. Boſc.

Media.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Media. Caput et thorax ſubpunctata, nigra, aene parum nitidula. Elytra ſubſtriata, nigra, pun cto magno humerali, rufo.

*plagiocephala.* 43. C. nigra, thorace elytrisque teſtaceis, capit craſſiuſculo. Ent. ſyſt. ſuppl. 113. 24. *

*Cryptocephalus plagiocephalus.* Ent. ſyſt. 1 60. 37. *

Habitat in Gallia meridionali. Muſ. D. de Seb ſtedt.

*pallens.* 44. C. nigra, cinereo-villoſa, thorace elytrisqu pallidis. Ent. ſyſt. ſuppl. 114. 25.

*Cryptocephalus pallens* Ent. ſyſt. 2. 61. 39. *

Habitat in China. Dr. Pflug.

*ruficollis.* 45. C. nigra. thorace rufo, elytris teſtaceis, pedibu anticis elongatis, femoribus apice vnidentati Ent. ſyſt. ſuppl. 114. 26.

*Cryptocephalus ruficollis.* Ent. ſyſt. 2. 61. 42.

Habitat in Europa auſtraliori. D. Prof. Helwig

*nigripes.* 46. C. capite thoraceque rufis, elytris pallidis, ped bus apice nigris. Ent. ſyſt. ſuppl. 114. 27.

*Cryptocephalus nigripes.* Ent. ſyſt. 2. 61. 43.

Habitat in America.

*brunnea.* 47. C. brunnea, antennis, corpore pedibusque n gris.

Habitat in America meridionali. D. Smidt. Mu D. Lund.

Parua. Antennae craſſae, nigrae, baſi ferruginea Caput et thorax ſublaeuia, brunnea, obſcur Elytra punctato-ſtriata, brunnea, immaculat

*Variat* corpore pedibusque rufis.

*coerulans.* 48. C. cyanea, nitida, antennis ſuſcis.

*Cryptocephalus coerulans.* Ent. ſyſt 2. 62. 45.

Habitat in Africa aequinoctiali. Muſ. D. Banks

49.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

49. C. rubro-cuprea, nitida, ſubtus cyanea. Ent. ſyſt. ſuppl. 114. 28. *cuprea.*
*Cryptocephalus cupreus.* Ent. ſyſt. 2. 62. 47.*
Habit t in Cayenna. D. v. Rohr.

50. C. cyanea, thorace pedibusque rufis. Ent. ſyſt. ſuppl. 114. 29. *cyanea.*
*Cryptocephalus cyaneus.* Ent. ſyſt. 2. 63. 52.* IV. 52
Geoff. Inſ. 1. 197. 4.
Habitat in Gallia. D. Mallet. Brunswigera in corylo

51. C. cyanea, ore thoracis marginibus pedibusque rubris. Ent. ſyſt. ſuppl. 114. 30. *bucephala.*
*Cryptocephalus bucephalus.* Ent. ſyſt. 2. 63. 54.* IV. 52
*Chryſomela bucephalu.* Schall. Act. Hall. 1. 276.
Habitat in Anthyllide vulneraria Saxoniae.

52. C. nigra, thorace rufo, elytris rufis: faſciis duabus cyaneis, pedibus nigris. Ent. ſyſt. ſuppl. 114. 31. *Scopolina.*
*Cryptocephalus Scopolinus.* Ent. ſyſt. 2. 68. 79.* IV. 52
*Chryſomela Scopolina.* Linn. ſyſt. Nat. 2. 597. 81.
~~Panz. Naturf. 24. 17. 23. tab. 1. fig. 23.~~ ref. Illig Mag. III. 165
Habitat in America et Europa auſtrali.

53. C. rufa, elytris punctis duobus coeruleſcentibus. Ent. ſyſt. ſuppl. 114. 32. *Koenigii.*
*Cryptocephalus Koenigii.* Ent. ſyſt. 2. 68. 80.*
Habitat Tranquebariae. Dr. Koenig.

54. C. elytris flauis: lineola punctisque tribus nigris. Ent. ſyſt. ſuppl. 115. 33. *paracentheſis.*
*Cryptocephalus paracentheſis.* Ent. ſyſt. 2. 68. 81.*

*Chry-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Chryſomela paracentheſis.* Linn. ſyſt. Nat. 2 1066. 1.

Habitat in Europae Salice, Betula.

*bicolor.* 55. C. atra, capite thoraceque flauis.

Habitat in America meridionali. D. Smidt. Muſ D. de Seheſtedt.

Antennae breues, craſſae, ſerratae, nigrae, baſ ferrugineae. Caput et thorax laeuia, fulua, ni tida. Elytra ſtriata, nigra. Corpus nigrum, fe moribus, inprimis anticis, fuluis.

*2puſtulata.* 56. C. atra, capite, thorace elytrorumque macula diſci rufis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et ſumma affinitas praecedentis, et forte mera eius varietas. Differt inprimis puncto elytrorum ferrugineo. Femora antica rufa.

*punctulata.* 56. C. atra, nitida, elytris punctatis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et ſumma affinitas C. nitidulae, at antennae et pedes concolores, et elytra magis punctata.

*nitidula.* 57. C. atra, aeneo-nitida, antennis flaueſcentibus.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Antennae flaueſcentes, apice parum fuſcae. Caput, thorax et elytra laeuia, glabra, atra, aeneo-nitida. Corpus cum pedibus atrum.

*concolor.* 58. C. viridi-aenea, nitida, antennis baſi flaueſcentibus, fronte impreſſa. Ent. ſyſt. ſuppl. 115. 35.

*Cryptocephalus concolor.* Ent. ſyſt. 2. 69. 83.

Habitat in Europa.

59. C.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

59. C. aenea, nitida, fronte cuprea. Ent. syst. suppl. 115. 34. *Hordei.*

*Cryptocephalus aeneus.* Ent. syst. 2. 69. 82. *

Habitat in Barbariae Hordeo murino. D. Prof. Vahl.

101. CRYPTOCEPHALVS. *Palpi* quatuor filiformes.
*Maxilla* vnidentata.
*Labium* corneum, integrum.
*Antennae* filiformes.

1. C. niger, thorace elytrisque rufis, elytris punctis duobus nigris. *elatus.*

Habitat in Tanger. D. Schousboe. Mus. D. de Seheſtedt.

Gigas in hoc genere. Antennae nigrae, basi rufae. Caput nigrum, immaculatum. Thorax rotundatus, laeuis, rufus, nitidus. Scutellum atrum. Elytra subpunctata, rufa, punctis duobus nigris, baseos humerali, posteriore pone medium. Corpus nigrum, cinereo-villosum.

2. C. rufus, elytris punctato-striatis: punctis tribus nigris. *6signatus.*

Habi-

---

Cryptocephali corpus parum, cylindricum, antice posticeque obtusum, glabrum, immarginatum, tardum; *capite* plano, inserto; *oculis* rotundatis, marginalibus; *antennis* thorace longioribus, distantibus, ante oculos insertis; *thorace* brevi, rotundato, transuerso; *scutello* paruo, rotundato; *elytris* rigidis, fornicatis, abdomine vix longioribus; *pedibus* validis, longiusculis, cursoriis; *tarsis* quadriarticulatis; *colore* vario, saepius nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat Tranquebariae. D. Daldorff. Muſ. D. Lun
Corpus magnum, rufum. Thorax gibbus, nitidu
Elytra punctato-ſtriata, punctis tribus nigris
1. baſeos 2. in medio. Pedes rufi.

*Salicis.* 3. C. ater, nitidus, elytris rufis: punctis tribus ni
gris. Ent. ſyſt. 2. 53. 3. *
Habitat in Saxonia. Dom. Hybner.

*6notatus.* 4. C. ater, elytris teſtaceis: punctis duobus linea
laque poſtica nigris. Ent. ſyſt. 2. 53. 4. *
Habitat in Barbaria. Muſ. D. Desfontaines.

Ill. III. 166

*laetus.* 5. C. viridi-aeneus, elytris teſtaceis: maculis du
bus nigris. Ent. ſyſt. 2. 54. 8. *
IX. Habitat in Saxonia. Dom. Hybner.

*lineatus.* 6. C. ruſeſcens, thorace maculis, elytris lineis du
bus nigris. Ent. ſyſt. 2. 55. 13. *
Habitat in Braſilia. Muſ. D. Banks.

*venuſtus.* 7. C. griſeo-fuſcus, thorace margine maculis du
bus flauis, elytris flauis: vittis duabus nigri
Ent. ſyſt. 2. 56. 16. *
Habitat in America. Muſ. Dr. Hunter.

*6guttatus.* 8. C. ater, glaber, elytris punctis ſex albis. Ent. ſyſ
ſuppl. 105. 17. *
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

*Gorteriae.* 9. C. ater, glaber, thorace elytrisque punctis quatuo
flaueſcentibus. Ent. ſyſt. 2. 56. 17.
*Chryſomela Gorteriae.* Linn. ſyſt. Nat. 2. 596
74. Amoen. acad. 6. 394. 16.
Habitat in Gorteria Cap. Bon. Spei.

*modeſtus.* 10. C. thorace rufo, elytris flauis rufo-lineatis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Medius. Antennae elongatae, ferrugineae, apic
fuſcae. Caput et thorax glabra, rufa, nitida
Elytr

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Elytra flaua, nitida, lineis numeroſis. punctatis. rufis: intermediis duabus ante apicem coeuntibus. Corpus flaueſcens.

11. C. rufus, elytris maculis tribus nigris: anteriore poſtice didyma. Ent. ſyſt. 2. 57. 22. * *didymus.*

Habitat in noua Hollandia. Muſ. D. Banks.

12. C. niger, thorace fuluo: punctis ſex nigris, elytris pallidis: punctis duobus. Ent. ſyſt. 2. 58. 24. * *flauicollis.*

Habitat in Sibiria. Muſ. D. Banks.

13. C. nigro-aeneus, elytris flauis: margine nigro. Ent. ſyſt. 2. 58. 27. * *marginatus.*

*Cryptocephalus phaleratus.* Act. Hall. 1. 216.

Panz. Fn. Germ. 68. tab. 1.

Habitat in Germaniae plantis.

14. C. niger, elytris teſtaceis: puncto humerali nigro. *humeralis.*

Habitat in Mauritania. D. Schousboe. Muſ. D. de Seheſtedt.

Statura et magnitudo praecedentis. Caput, thorax, corpus nigra, obſcura, immaculata. Elytra vix punctata, teſtacea, puncto humerali nigro.

*Variat* puncto poſtico vno alteroue elytrorum.

15. C. thorace elytrisque obſcure aeneis, pubeſcentibus. Ent. ſyſt. 2. 58. 28. * *pubeſcens.*

Habitat in America boreali.

16. C. niger, nitidus, elytris rubris: punctis duobus nigris, antennis longitudine corporis. Ent. ſyſt. 2. 59. 29. *2punctatus*

*Chryſomela 2punctata.* Linn. ſyſt. Nat. 2. 597. 78. Fn. Sv. 548.

Geoff. Inſ. 1. 234.

Degeer Inſ. 5. 332. 35.

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 68. tab. 2.
Payk. Fn. Sv. 2. 142. 15.
Schaeff. Icon. tab. 8. fig. 8.
Petiv. Gazoph. tab. 31. fig. 4.
Habitat in Europae Corylo.

*lineola.* 17. C. ater, nitidus, elytris rubris: linea media atra, antennis longitudine corporis. Ent. syst. 2. 59. 30. *
*Chrysomela 2punctata.* Fn. Sv. 548. β.
*Cryptocephalus dispar.* Payk. Fn. Sv. 2. 143. 15. β.
Habitat in Europae Corylo.
~~Forte~~ mera praecedentis varietas.

*2maculatus.* 18. C. niger, obscurus, thorace fulvo, elytris testaceis: punctis duobus nigris. Ent. syst. 2. 59. 31. *
~~*Chrysomela melanocephala.* Act. Hall. 1. 272.~~
Habitat in Italia. Dr. Allioni.

*cordiger.* 19. C. C. thorace variegato, elytris rubris: punctis duobus nigris. Ent. syst. 2. 59. 32.
*Chrysomela cordigera.* Linn. syst. Nat. 2. 598. 79. Fn. Sv. 588.
Geoff. Inf. 1. 235. 2.
Payk. Fn. Sv. 2. 133. 7.
Panz. Fn. Germ. 13. tab. 6.
Degeer. Inf. 5. 330. 33.
Schaeff. Icon. tab. 30. fig. 1.
Habitat in Europae Corylo, Salice.
Thorax ater, lateribus albis: puncto nigro. Linea dorsalis antica alba, terminatur postice in maculam cordatam, distinctam.

*variabilis.* 20. C. thorace atro, albo-trilineato, elytris rubris: punctis tribus nigris. Ent. syst. suppl. 105. 32. *

Cry-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cryptocephalus variabilis.* Schneid. Magaz. 2. 36. 5
Panz. Fn. Germ. 13. tab. 7.
Habitat in Germaniae Salice.

21. C. niger, thorace lineis tribus flauis. Ent. ſyſt. ſuppl. 105. 32. * *trilineatus.*
Habitat in Germania.

22. C. ater, thorace linea longitudinali rubra. Ent. ſyſt. 2. 59. 33. * *Bothnicus.*
Panz. Fn. Germ. 68. tab. 3.
*Chryſomela Bothnica.* Linn. ſyſt. Nat. 2. 598. 90. Fn. Sv. 557.
*Cryptocephalus varians.* Payk. Fn. Sv. 2. 148. 21.
Herbſt. Arch. tab. 44. fig. 8.
Degeer Inſ. 5. 335. 39.
Habitat in Sueciae Salice.

23. C. ater, capite, thorace pedibúsque rufis, thorace punctis duobus ~~rufis~~ nigris. Ent. ſyſt. 2. 60. 34. * *frenatus.*
Panz. Fn. Germ. 68. tab. 4.
Habitat in Auſtria. D. Schneider.

24. C. niger, obſcurus, elytris ſubreticulatis. Ent. ſyſt. ſuppl. 106. 34. * *luridus.*
Habitat in America boreali. D. Mauerhoff.

25. C. ater, thorace punctato, elytris ſulcatis: ſulcis rugoſis. *impreſſus.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Thorax punctis numeroſiſſimis impreſſus. Elytra ſtriis eleuatis, hinc inde interruptis, ſulcata ſulcis rugoſis.

26. C. niger, thorace elytrisque teſtaceis; ſutura nigra. Ent. ſyſt. 2. 60. 38. * *Coryli.*

*Chry-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Chryſomela Coryli.* Linn. ſyſt. Nat. 2. 598. 88. Fn. Sv. 555.
Panz. Fn. Germ. 68. tab. 6.
Payk. Fn. Sv. 2. 137. 10.
Habitat in Europae Corylo.

*variegatus.* 27. C. niger, thorace linea dorſali abbreuiata, marginibusque rubris, elytris teſtaceis. Ent. ſyſt. 2. 61. 40. ♂
*Cryptocephalus variegatus.* Payk. Fn. Sv. 2. 135. 8.
Habitat in Italia. Dr. Allioni.

*trifaſciatus* 28. C. ſupra teſtaceus, thorace, faſcia, elytris tribus nigris. Ent. ſyſt. 2. 61. 41. ♂
Habitat in China. Dr. Pflug.

*imperialis.* 29. C. ater, nitidus, elytris teſtaceis: punctis quinque nigris. Ent. ſyſt. ſuppl. 106. 44. ♂
*Cryptocephalus imperialis.* Laichart. Tyrol. 49. 16.
Habitat in Gallia. Muſ. D. Boſc.

*6punctatus.* 30. C. niger, thorace variegato, elytris rubris: maculis tribus nigris. Ent. ſyſt. 2. 62. 44. ♂
*Chryſomela 6punctata.* Linn. ſyſt. Nat. 2. 599. 92. Fn. Sv. 559.
Degeer Inſ 5. 331. 34. tab. 10. fig. 9.
Sulz. Inſ. tab. 3. fig. 18.
Schaeff. Icon. tab. 30. fig. 3.
Panz. Fn. Germ. 68. tab. 7.
Payk. Fn. Sv. 2. 136. 9.
Habitat in Europae plantis.

*flaueſcens.* 31. C. ſupra flaueſcens, elytris punctis quinque nigris: marginalibus diſtinctioribus. Ent. ſyſt. 4. App. 448. 33. ♂
Habitat in Germania. D. Smidt.

32. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

32. C. rufus, elytris flauis: vittis duabus abbreuiatis atris. Ent. syst. suppl. 106. 44. * *ornatus.*
Habitat in Carolina. Mus. D. Bosc.

33. C. nigro-cyaneus, antennis pedibusque nigris, elytris substriatis. Ent. syst. 2. 62. 46. * *violaceus.*
Habitat in Germania. D. Smidt. IX.

34. C. rufus, elytris rufo nigroque variis. Ent. syst. suppl. 107. 44. * *congestus.*
Habitat in Carolina. Mus. D. Bosc.

35. C. rufus, elytris oblique striatis, pedibus flauescentibus. *obliquus*
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura omnino C. congesti. Thorax laeuis, rufus, nitidus, immaculatus. Elytra vti in C. congesto oblique striata, rufa. Pedes flauescentes.

36. C. cyaneus, nitidus, fronte, thoracis fascia femoribusque rubris. Ent. syst. 2. 62. 48. * *rubrifrons.*
Habitat - - Dom. Prof. Leske.

37. C. violaceus, capite, thorace femoribusque dentatis rufis. Ent. syst. 2. 62. 49. * *calcaratus.*
Habitat in Sierra Leon. Mus. D. Banks.

38. C. aeneus, cupro-nitidus, femoribus posticis dentatis. *dentipes.*
Habitat Tranquebariae. Mus. D. de Seheſtedt.
Statura C. calcarati, at minor. Antennae nigrae. Caput et thorax punctata, viridi-aenea. Elytra punctata, aenea, cupro-nitida. Corpus aeneum, femoribus posticis dente paruo, acuto.

39. C. obscure cyaneus, tibiis posticis apice lobatis. Ent. syst. 2. 63. 50. * *lobatus.*
Panz. Fn Germ. 13. tab. 9. II. 31.
Habitat in Hungaria. D. Hybner.

40. C.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*cinctus.* 40. C. capite thoraceque rufis, elytris nigris: margine rufo. Ent. ſyſt. 2. 63. 53. *

Habitat in America meridionali. Muſ. Dr. Hunter.

*Billardierii* 41. C. flauus, thorace punctis duobus, coleoptris vittis tribus atris.

Habitat in noua Cambria. D. Billardiere.

Statura parua C. Seheſtedtii. Antennae baſi flauae, apice fuſcae. Caput laete flauum, oculis clypeoque nigris. Thorax gibbus, flauus: punctis duobus atris. Elytra glabra, laeuia, flaua: vitta media ſuturaque late atris, apicem tamen haud attingentibus. Scutellum flauum. Abdomen atrum, pedibus atris.

*Billardiere*, Profeſſor Pariſienſis, Botanicus eximius, qui Inſecta Syriae et Auſtralaſiae collegit, communicauit.

*Lundii.* 42. C. ſupra teſtaceus, oculis atris. Ent. ſyſt. ſuppl. 107. 54.

Habitat Tranquebariae. Muſ. D. Lund.

Dom. Lund, amiciſſimus Norwagus, miram Inſectorum copiam collegit, beneuole communicauit.

*Seheſtedtii.* 43. C. capite thoraceque rufis, nitidis, elytris flauis: vitta lata atra. Ent. ſyſt. ſuppl. 107. 54. *

Habitat in India orientali. D. Daldorff. Muſ. D. de Seheſtedt.

Dom. de Seheſtedt, Norwagus, cubicularius Regis, Lundio mihique amicitia iunctus, inſecta vndiquaque coërcet, in Muſeo diſponit.

*Vahlii.* 44. C. rufus, thorace maculis duabus atris, elytris atris: faſcia punctoque rufis. Ent. ſyſt. ſuppl. 108. 54 *

Habitat in Iudia orientali. Muſ. D. Lund.

Dom.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Dom. Vahl, Profeſſor Haſnienſis, Botanicorum princeps, Barbariae, Hiſpaniae Entomologiam caſtigauit.

5. C. cyaneus, nitidus, thoracis lateribus elytrorum apicibus femoribusque rubris. Ent. ſyſt. 2. 63. 55. * *collaris.*
Habitat in Sibiria. Muſ. D. Banks.

6. C. viridi-coeruleus, antennis nigris. Ent. ſyſt. 2. 63. 56. *fericeus.*
*Chryſomela fericea.* Linn. ſyſt. Nat. 2. 598. 86.
Fn. Sv. 554.
Geoff. Inſ. 1. 233.
Oliv. Inſ. 96. tab. 1. fig. 5. IX.
Payk. Fn. Sv. 2. 139. 12.
Degeer Inſ. 5. 334. 37.
Schaeff. Icon. tab. 65. fig. 2. 3. 5.
Sulz. Hiſt. Inſ. tab. 3. fig. 13.
Habitat in Europae Salice.

7. C. viridis, nitens, ore pedibusque teſtaceis. Ent. ſyſt. 2. 64. 57. Payk. Fn. Sv. 2. 140. 13. *nitens.*
*Chryſomela nitens.* Linn. ſyſt. Nat. 2. 598. 84.
Fn. Sv. 551.
Panz. Fn. Germ. 68. tab. 8. VI. 13.
Degeer Inſ. 5. 334. 38.
Habitat in Europae Corylo.

8. C. violaceus, thorace elytrisque aeneis, antennis fuſcis. Ent. ſyſt. 2. 64. 58. * *glabratus.*
Geoff. Inſ. 1. 233. 4.
Schaeff. Icon. tab. 77. fig. 8. Habit. in America.
Herbſt. Arch. tab. 23. fig. 23. Mus. D. Banks.
Habitat in Europae auſtralioris gramineis.

9. C. niger, capite thoraceque flauo-maculatis, elytris flauis: vittis duabus abbreuiatis, nigris. *viduatus.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Carolina. Muſ. D. Boſc.

Statura et magnitudo C. vittati. Antennae nur flauae, nunc fuſcae. Caput nigrum, macul tribus frontalibus, flauis. 1. 2. Thorax fcabe niger, margine omni lineolisque tribus flaui media cum margine antico, lateralibus cu margine poſtico coëuntibus. Elytra punctat vittis duabus latis, nigris, quae tamen apice haud attingunt. Corpus nigrum, pedibus v riegatis.

*lituratus.* 50. C. niger, thoracis margine elytrisque vittis du bus flauis.

Habitat in Carolina. Muſ. D. Boſc.

Statura C. vittati, at duplo fere minor. Cap cum antennis nigrum, immaculatum. Thor niger, margine late flauo, et in hoc margi punctum magnum atrum. Elytra punctat ſtriata, atra, vittis duabus poſtice coëuntibu flauis. Corpus atrum.

*Variat* rarius rudimento vittae mediae, nunc baſin, nunc verſus apicem.

*puſtulatus.* 51. C. ater, thoracis margine antico elytrorumq baſi maculisque tribus rubris. Ent. ſyſt. ſupp 108. 58. *

Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

*vittatus.* 52. C. niger, elytris, margine ſtriaque abbreuia flauis. Ent. ſyſt. 2. 64. 59. *

Geoff. Inſ 1. 233. 4.

Schaeff. Icon. tab. 77. fig. 8.

Herbſt. Arch. tab. 23. fig. 23.

Habitat in Europae auſtralioris graminoſis.

*limbatus.* 53. C. ferrugineus, thorace rufo, elytrorum ſutu cyaneo-nitidis. Ent. ſyſt. 2. 64. 60. *

Habitat in Cayenna. Dr. Schulz.

54.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

54. C. thorace aeneo, nitido, elytris cyaneis, capite, antennarum basi pedibusque flauis. Ent. syst. 2. 64. 61. * *nitidulus.*

Habitat in Russia meridionali. D. Boeber.

55. C. rufus, elytris viridibus. *tricolor.*

Habitat in noua Cambria. D. Billardiere.

Paruus. Antennae nigrae. Caput nigrum, fronte rufa. Thorax laeuis, rufus, nitidus. Elytra punctata, nitida, immaculata. Abdomen albidum, pedibus rufis.

56. C. ater, nitidus, ore, pedibus basique antennarum lutescentibus. Ent. syst. 2. 65. 62. * *labiatus.*

*Chrysomela lubrata.* Linn. syst. Nat. 2. 598. 87. Fn. Sv. 553.

Panz. Fn. Germ. 68. tab. 9. VI. 13

Payk. Fn. Sv. 2. 147. 20.

Habitat in Europae rosis.

57. C. violaceus nitidus, ore ~~pube~~lescente, antennis pedibusque nigris. Ent. syst. 2. 65. 63. * *flauilabris.*

*Cryptocephalus puncticulus.* Mysl. Prodr. 519.

Payk. Fn. Sv. 2. 146. 18. ~~tab. mas 6 marginatus~~ cf. Rossi Hellen. 211 C. marginatus

Habitat in Saxoniae Betula. D. Schaller.

*Larua* saccata: sacco antice oblique truncato.

58. C. ater, nitidus, capite pedibusque luteis. Ent. syst. 2. 65. 64. * *flauipes.*

Panz. Fn. Germ. 68. tab. 10. VI. 14

Payk. Fn. Sv. 2. 144. 16.

Habitat in Italia et Germania. Dr. Allioni.

59. C. coeruleo-ater, nitidus, fronte pedibusque flauis. Ent. syst. 2. 65. 65. * *flauifrons.*

Habitat Kiliae.

60. C. niger, capite, elytrorum apicibus pedibusque flauis. Ent. syst. 2. 65. 66. * *Hybneri.*

 Panz. IV. 49

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 39. tab. 16.
Habitat in Germania. Dom. Hybner.

*Moraei.* 61. C. ater, elytris maculis duabus flauis marginalibus. Ent. ſyſt. 2. 66. 67.
*Chryſomela Moraei.* Linn. ſyſt. Nat. 2. 597. 82
Fn. Sv. 550.
Geoff. Inſ. 1. 235. 5.
Payk. Fn. Sv. 2. 145. 17.
Panz. Fn Germ. 68. tab. 11.
Schaeff. Icon. tab. 30. fig. 7.
tab. 238. fig. 5-2
Habitat in Europae Hypericis.

*canclius.* 62. C. teſtaceus, elytris punctis duobus ſuturaque nigris.
Habitat in Carolina. Muſ. D. Boſc.
Paruus. Antennae fuſcae, baſi teſtaceae. Caput et thorax laeuia, teſtacea, immaculata. Elytra punctato-ſtriata, punctis duobus: altero ad baſin, altero in medio ſuturaque tenuiſſima nigris. Corpus teſtaceum.

*8guttatus.* 63. C. ater, elytris maculis quatuor flauis. Ent. ſyſt. 2. 66. 68. *
Habitat in Saxoniae Fraxino. D. Hybner.

*15guttatus* 64. C. rufus, thorace maculis tribus, elytris ſex flauis. Ent. ſyſt. 2. 66. 69. *
Habitat in Braſilia. Muſ. D. Banks.

*guttula.* 65. C. thorace elytrisque rufis, albo-punctatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura parua praecedentis. Caput nigrum lineolis duabus obliquis, flauis. Thorax laeuis, rufus, nitidus, punctis ſeptem albis, 3 marginali antico connexis, 4. Elytra laeuia, rufa, punc-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

etis plurimis ſparſis, albis. Corpus rufum, pectore nigro. Anus prominet punctis duobus albis.

*Variat* varius colore viridi-aeneo, loco rufi.

6. C. thorace teſtaceo: vitta marginali atra, coleoptris flauis: punctis decem nigris. Ent. ſyſt. 2. 66. 70. * *10punctatus.*

*Cryſomela 10maculata* Linn. ſyſt. Nat. 2. 599. 93. Fn. Sv. 560.

*Cryptocephalus hieroglyphicus.* Herbſt. Arch. tab. 23. fig. 26. VI. 14

*Cryptocephalus varians.* Payk. Fn. Sv. 2. 148. 21. β.

Panz. Fn. Germ. 68. tab. 12.

Habitat in Europae Salice. *Femati. Ill.*

*Varietatem* C. Bothnici credit D. Paykull.

7. C. thorace flauo: punctis nigris quinque. coleoptris flauis: punctis nigris decem. Ent. ſyſt. 2. 66. 71. * *hieroglyphicus.*

Panz. Fn. Germ. 68. tab. 13. VI. 15

Habitat in Italia. Dr. Allioni.

8. C. niger, thorace fuluo: punctis quatuor nigris, elytris teſtaceis, lineola baſeos punctisque duobus. Ent. ſyſt. ſuppl. 108. 71. * *coloratus.*

*Cryptocephalus 4maculatus.* Schneid. Mag. IX. 195. 2.

Habitat in Germania. D. Mauerhoff.

9. C. obſcure griſeus, femoribus poſticis vnidentatis. *dubius.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Huius generis videtur. Antennae ſiliformes, fuſcae, articulo primo maiori, incuruo. Caput, thorax, elytra ſubpunctata, obſcure griſea, minime

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nime nitida. Thorax vtrinque in angulum par uum, obtuſum excurrit. Pedes obſcuri, femo ribus poſticis ſubdentatis.

*12punctatus.* 70. C. thorace fuluo: punctis duobus, elytris te ſtaceis: punctis quinque nigris. Ent. ſyſt. 2 67. 72. *

Habitat in Germania. D. de Loewenſkiold.

*haemorrhoidalis.* 71. C. cyaneus, elytris apice pedibusque fuluis. Ent ſyſt. 2. 67. 73. *

Panz. Fn. Germ. 68. tab. 14.

~~Roſſ. Fn. Etr. 1. 97. 247. β.~~

Schaeff. Icon. tab. 77. fig. 7.

Habitat Pariſiis. Muſ. D. Boſc.

*ſcutellaris.* 72. C. ater, thoracis margine, ſcutello elytrorumqu maculis duabus rufis.

Habitat in Carolina. Muſ. D. Boſc.

Statura et magnitudo C. bipuſtulati. Antenn flauae, apice nigrae. Caput nigrum, orbita oc lorum flaua. Thorax ſubpunctatus, ater ma gine flauo. Scutellum flauum. Elytra punct to-ſtriata: maculis duabus flauis; anterio maiore, transuerſa. Corpus nigrum, ano pr minulo, nunc toto flauo, nunc maculis duab flauis. Pedes flaui.

*4puſtulatus* 73. C. ater, elytris macula apicis rufa. Ent. ſyſt. 67. 74. *

*Cryptocephalus Podae.* Laichart. Tyrol. 197.

*Cryptocephalus diſpar.* Payk. Fn. Sv. 2. 14 15. Z.

Panz. Fn. Germ. 68. tab. 15.

Herbſt. Arch. tab. 44. fig. 6.

Habitat in Chryſanthemo coronario.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

74. C. rufus, thorace maculis duabus, elytris faſciis duabus nigris. Ent. ſyſt. 2. 67. 75. * *2faſciatus.*

Habitat in Africa aequinoctiali. Muſ. D. Banks.

75. C niger, elytris laeuibus, rufo-bimaculatis. Ent. ſyſt. 2. 68. 76. * *4puſtulatus*

*Chryſomela 4puſtulata.* Linn. ſyſt. Nat. 2. 597. 80. Fn. Sv. 549.

Habitat in Suecia.

76. C. niger, elytris punctato-ſtriatis: faſcia punctoque apicis teſtaceis. Ent. ſyſt. 2. 68. 77. * *notatus.*

Habitat in America. Muſ. Dr. Hunter.

77. C. niger, thorace elytrisque ferrugineo-variis, pedibusque ferrugineis: geniculis nigris. Ent. ſyſt. 2 68. 78. * *hiſtrio.*

Roſſ. Fn Etr. 1. 98. 249.

Habitat in Italia. Dr. Allioni.

78. C. nigro-aeneus, antennis pedibusque teſtaceis. *ſmaragdulus.*

Habitat in Guinea. Muſ. D. de Seheſtedt.

Paruus. Caput, thorax, elytra nigro-aenea, nitida. Elytra ſtriata. Antennae elongatae pedesque teſtacei.

79. C. ater, thoracis margine elytrisque punctis tribus albis. *3guttatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paruus. Caput nigrum, puncto frontali et verticali albis. Thorax laeuis, ater, nitidus, margine exteriori, linea dorſali, abbreuiata albis. Elytra ſubpunctata, atra: punctis tribus albis; vno pone alterum. Abdomen atrum, margine albo. Femora baſi alba.

80. C. cyaneus, nitidus, antennarum articulo primo flaueſcente, elytris punctato-ſtriatis. *ſaponatus.*

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Carolina. Muſ. D. Boſc.

Corpus paruum praecedentium cyaneum, nitidum, ſolo articulo antennarum primo flauo. Thorax laeuis. Elytra punctato-ſtriata.

*Spengleri.* 81. C. thorace rufo. nitido: punctis duobus nigris, elytris teſtaceis: macula poſtice cyanea. Ent. ſyſt. ſuppl. 109. 81. *

Habitat in India. D. Daldorff. Muſ. D. de Seheſtedt.

*hecticus.* 82. C. pallidus, immaculatus, elytris ſtriatis, antennis apice fuſcis.

Habitat Tranquebariae. Muſ. D. Lund.

Corpus pallidum C. Pini. Antennae filiformes apice fuſcae. Caput et thorax laeuia, pallida, nitida. Elytra punctato-ſubſtriata, pallida, immaculata. Corpus pallidum.

*Pini.* 83. C. teſtaceus, elytris vage punctatis, pallidis, antennis fuſcis. Ent. ſyſt. 2. 69. 84.

*Chryſomela Pini.* Linn. ſyſt. Nat. 2. 598. 89. Fn. Sv. 556.

*V. 50.* Panz. Fn. Germ. 39. tab. 17.

Payk. Fn. Sv. 2. 138. 11.

Habitat in Sueciae Pino.

*reticulatus.* 84. C. thorace elytrisque albis, teſtaceo-reticulatis. Ent. ſyſt. 2. 69. 85. *

Habitat in Cayenna. D. v. Rohr.

*puſillus.* 85. C. thorace fuluo, elytris ſtriatis, teſtaceis, nigro-maculatis. Ent. ſyſt. 2. 69. 86. *

Roſſ. Fn. Etr. 1. 100. 255.

Payk. Fn. Sv. 2. 149. 22. β.

Habitat Hamburgi. Dr. Schulz.

*Varietatem* C. minuti credit D. de Paykull.

*nanus.* 86. C. rufus, elytris ſtriatis atris.

Habitat in Carolina. Muſ. D. Boſc.

Paruus.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paruus. Antennae nigrae, baſi rufae. Caput et thorax laeuia, rufa, nitida. Elytra ſtriata, atra, immaculata. Corpus rufum.

87. C. thorace fuluo, elytris ſtriatis, teſtaceis, immaculatis. Ent. ſyſt. 2. 70. 87. * *minutus.*
Panz. Fn. Germ. 39. tab. 18.
Payk. Fn. Sv. 2. 149. 22.
Habitat Kiliae.

88. C. ater, nitidus, capite thoraceque rufis, pedibus flauis. Ent. ſyſt. ſuppl. 109 87. * *parnulus.*
Habitat in America boreali. D. Mauerhoff.

89. C. ater, capite thoraceque fuluis, elytris linea marginali baſeos alba. Ent. ſyſt. 2. 70. 88. * *gracilis.*
Geoff. Inſ. 1. 237. 11.
Habitat in Gallia et Dania. Muſ. D. Boſc.
β Caput et thorax laeuia, aenea, obſcure nitida: margine thoracis flauo. Elytra ſubtiliſſime ſtriata, aenea. Pedes flaui.

91. C. ater, nitidus, elytris teſtaceis: ſutura nigra. Ent. ſyſt. 2. 70. 89. * ~~*murinus.*~~ *pygmaeus*
Habitat in Gallia. Muſ. D. Boſc.
*Variat* rarius, elytris vitta media abbreuiata, atra.

*femoribus ſaltatoriis.*

92. C. ſaltatorius, ater, elytris rubris apice atris. ~~*pygmaeus.*~~ *ſaliens*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura parua praecedentium. Antennae nigrae, baſi ferrugineae. Caput et thorax laeuia, ferruginea, apice late atra. Corpus et pedes ferruginea, femoribus poſticis valde incraſſatis, atris.

93. C. ſaltatorius, ferrugineus: capite, elytrorum apicibus femoribusque poſticis atris. *ſaltator.* ~~*ſaliens.*~~

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentis, at paullo minor et diſtinctus. Antennae nigrae, baſi ferrugineae. Caput nigrum. Thorax laeuis, ferrugineus immaculatus. Elytra laeuia, rufa, apice atra. Corpus ferrugineum, femoribus poſticis valde incraſſatis, atris.

## 102. HISPA. *Palpi* aequales, medio craſſiores
*Maxilla* biſida.
*Labium* corneum, integrum.
*Antennae* cylindricae.

~~ſolitaria~~ Spinoſa 1. H. ſpinoſiſſima, rufa, elytris nigris.
*Hiſpa ſpinoſa.* Web. Inſ. 65. 2.
Habitat in Sumatra. D. Daldorff.
Statura et magnitudo H. atrae. Caput rufum, in erme, antennis nigris. Thorax rufus, ſpini quatuor nigris lateralibus trifidis, intermedii biſidis. Elytra nigra, ſpinis plurimis, porrecti validis, acutis. Corpus rufum.

~~ſpinoſa~~ atra 2. H. atra, thorace elytrisque ſpinoſis. Ent. ſyſt. 2 70. 1. Linn. ſyſt. Nat. 2. 603. 1.
Geoff. Inſ. 1. 243. 7.
Act. Soc. Berol. Nat. Cur. 4. tab. 7. fig. 6.
Roſſ. Fn. Etr. 52. 129.

VIII. 23

Hab

---

Hiſpae corpus paruum, oblongum, immarginatum, tardum *capite* ouato, inferto; *antennis* approximatis et fere co natis, longitudine thoracis, inter oculos infertis; *oc* magnis, oblongis, marginalibus; *thorace* breui, rotund to; *ſcutello* triangulo; *elytris* rigidis, longitudine abdom nis, ſaepius, aculeatis, aut apice dentatis; *pedibus* bre bus, validis, rotundatis, curſoriis; *tarſis* quadriarticulati *colore* vario, minus nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Europa auſtraliori ad radices graminum.
Duplo maior e Guinea in Muſ. D. de Seheſtedt.
vix diſtincta.

3. H. thorace vtrinque palmato, ſexdentato, flauo; lineis duabus ſanguineis dorſalibus.
Habitat Tranquebariae. D. Daldorff. Muſ. D. Lund.
Statura omnino et ſumma affinitas H. Hyſtricis. Antennae fuſiformes, ſanguineae. Thorax vtrinque palmato-ſexdentatus, flauus, lineis duabus dorſalibus, ſanguineis. Elytra punctato-ſtriata, obſcura, ſpinis duabus erectis, acutis, validis, rufis et aliquot minoribus, atris. Margo ſpina valida baſeos atra, tunc palmato-ſexdentatus, flauus, denique ſolitaria, flaua, et tandem palmato-triſpinoſus: ſpina vnica flaua, duabus rufis, demum apice ſerratus, dentibus anticis flauis, poſticis rufis.

4. H. teſtaceo-nigroque-variegata, thorace vtrinque dilatato, quinqueſpinoſo. Ent. ſyſt. ſuppl. 116. 2. *
Habitat in Indiae Myricetis. D. Daldorff.

5. H. ſpinoſa, corpore teſtaceo, antennis oculisque. Ent. ſyſt. 2. 71. 2. *
Linn. ſyſt. Nat. 2. 603. 2.
Roſſ. Fn. Etr. 52. 128.
Habitat in Africa.

6. H. antennis ſerratis, hirta atra, elytris macula baſeos rufa. Ent. ſyſt. 2. 71. 3. *
Panz. Fn. Germ. 34. tab. 17.
Habitat in Italia. Dr. Allioni.

7. H. antennis ſerratis, hirta atra, elytris maculis duabus rufis. Ent. ſyſt. ſuppl. 116. 3. *
Habitat in Gallia. D. Brogniard.

8. H.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſpinipes.* 8. H. rufa, elytris nigris, rufo-punctatis, tibiis anticis vniſpinoſis. Ent. ſyſt. 4. App. 448.
Habitat Surinami. Muſ. D. Holthuyſen.

*ſanguinicollis.* 9. H. thorace elytrorumque baſi ſanguineis, elytris ſerratis. Ent. ſyſt. 2. 71. 5. *
Linn. Mant. 1. 530.
Habitat in America. D. v. Rohr.

*Morio.* 10. H. atra, immaculata, elytris ſulcatis, apice rotundatis, ſerratis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura omnino H. ſerratae, at tota atra, immaculata.

*haemorrhoidalis.* 11. H. rufa, elytris nigris, apice integris, teſtaceis.
Web. Inſ. 64. 1.
Habitat in Sumatra. D. Daldorff.
Statura praecedentium. Antennae nigrae, apice parum rufeſcentes. Caput et thorax rufa, immaculata. Elytra ſulcata: ſulcis transuerſim ſtriatis, nigra, apice integra, teſtacea. Corpus rufum.

*quadrata.* 12. H. elytris ſerratis, baſi flauis, apice purpuraſcentibus.
Habitat in Carolina. Muſ. D. Boſc.
Magnitudo H. ſanguinicollis. Caput flauum, antennis purpuraſcentibus. Thorax ſupra flauus, ſubtus purpuraſcens. Elytra ſerrata, ſulcata, baſi flaua, margine tamen purpuraſcente, apice purpurea. Corpus purpuraſcens, pedibus flauis.

*ſerrata.* 13. H. nigra, elytris rufeſcentibus, apice ſerratis nigris. Ent. ſyſt. 2. 72. 6. *
Habitat Surinami. Muſ. Dr. Hunter.

14. H.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

14. H. atra, thorace abdominisque margine sanguineis, elytris apice ferratis. *thoracica.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo praecedentis. Thorax laeuis, ſanguineus, immaculatus. Elytra ſulcata, atra, apice valde ferrata, at haud truncata. Corpus nigrum, abdominis margine ferrato.

15. H. atra, thoracis margine elytrorumque macula, baſeos rufis, elytris apice ferratis. *humeralis.*
Habitat in America meridionali. Muſ. D. de Seheſtedt.
Statura H. ferratae. Caput atrum, macula frontali rufa. Thorax laeuis, ater, linea marginali rufa, quae poſtice cum macula elytrorum coit. Elytra ſulcata, atra, macula oblonga, baſeos rufa, apice rotundata, ferrata.

16. H. nigra, thoracis margine, elytris ferratis: macula baſeos faſciaque media flauis. Ent. ſyſt. 2. 72. 7. ♀ *dentata.*
Voet. Col. 2. tab. 44. fig. 1. β.
Habitat in Cayenna. D. v. Rohr.

17. H. atra, elytris truncatis, angulato-ferratis. *maura.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura omnino H. angulatae, at tota atra, pedibus ſolis piceis. Elytra ſulcata, apice truncata, dentata: dente exteriori maiori, angulato.

18. H. thorace coleoptrisque atris: vittis duabus ferrugineis, apice truncatis, ferratis. *truncata.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo praecedentis. Caput cum antennis atrum. Thorax punctatus, ater: vittis duabus

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

duabus latis, ferrugineis cum vittis coleoptr rum coëuntibus. Elytra ſulcata, atra, vitta la ferruginea ante ſuturam, quae tamen apice haud attingit, et in hac vitta faſcia pone m dium atra. Apex truncatus, acute ſerratu Corpus rufum, immaculatum.

*angulata.* 19. H. flaua, capite, thoracis linea dorſali elytr rumque margine ſericeato nigris. Ent. ſyſt. 72. 8. *

Habitat in Cayenna. D. v. Rohr.

*haſtata.* 20. H. ſanguinea, elytris ſulcatis: vitta interrup coerulea baſi apiceque vniſpinoſis.

Habitat in America meridionali. D. Smidt. Mu D. de Seheſtedt.

Statura praecedentium, at diſtincta. Antennae r fae, baſi fuſcae. Caput et thorax punctat ſanguinea, immaculata. Elytra ſulcata, ruf vitta media hinc inde interrupta, violacea, ba apiceque ſpina valida, acuta, incurua. Ape alias integer. Corpus rufum.

*emargina-ta.* 21. H. nigra, elytris baſi macula rufa, apice bider tatis. Ent. ſyſt. 2. 72. 9. *

Habitat Surinami. Muſ. D. Boſc.

*eleuata.* 22. H. nigra, thoraci margine elytrorumque baſi ru fis, elytris linea eleuata vnica apice quadriden tatis.

Habitat in America meridionali. D. Smidt. Mu D. Lund.

Statura et ſumma affinitas H. carinatae. Capu rufum, ore antennisque nigris. Thorax rufus macula magna dorſali maculaque vtrinque ſub lateribus rufis. Elytra punctato-ſtriata, atra macula oblonga baſeos rufa, linea vnica ele uata, carinata, apice ſpinoſa. Praeterea dente

quatuo

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717 Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

quatuor validi in apice elytrorum: intermediis minoribus. Corpus nigrum, abdominis macula baſeos margineque rufis. Pedes nigri, femoribus rufis.

3. H. nigra, thoracis margine elytrorumque baſi rufis, elytris vniſtriatis, apice bidentatis. *carinata.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Caput rufum, margine poſtico antennisque nigris. Thorax niger, margine antico et laterali rufo, antico, dentem in medio emittente. Elytra punctato-ſtriata, nigra, macula baſeos rufa lineaque media eleuata, carinata, apice bidentata. Pedes nigri, femoribus baſi rufis.

4. H. nigra, thorace ferrugineo, elytris ſtriatis, apice ſubtiliſſime ſerratis. *ruficollis.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Parua. Caput nigrum. Thorax rotundatus, rufus, immaculatus. Elytra ſulcata, ſulcis punctato-ſtriatis, apice parum ſerratis. Corpus nigrum.

5. H. atra, thorace, antice ſcutelloque fuluis, elytris ſerratis, angulatis, plicisque eleuatis. *plicatula.*

Habitat in Carolina. Muſ. D. Boſc.

Media. Caput nigrum, antennis baſi flauis. Thorax ater, antice fuluus: punctis aliquot atris. Elytra ſerrata, apice angulata, et in diſco plicae plurimae eleuatae, compreſſae, acutae. Pedes nigri, femoribus annulo fuluo.

6. H. flaua, antennis elytrorumque ſutura nigris. *futuralis.*

*Hiſpa roſea.* Web. Inſ. 66. 5.

Habitat in Carolina. Muſ. D. Boſc.

Reliquis minor. Caput flaueſcens, antennis nigris. Thorax ſcaber, teſtaceus, margine elytrorum tenuiſ-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tenuiſſime punctisque aliquot nigris. Elytra ſulcata, integra, flaua, ſutura nigra. Puncta interdum ſparſa, nigra in diſco.

*Variat* elytris corporeque toto teſtaceis, antennis ſolis fuſcis.

*ſerraticornis.* 27. H. antennis baſi compreſſis, ſerratis. nigra, thoracis margine elytrisque macula baſeos flauis. Ent. ſyſt. 2. 73. 10. *

Habitat Surinami. Muſ. D. Boſc.

*compreſſicornis.* 28. H. antennis compreſſis, ſerratis, nigra, thoracis margine elytrorumque macula baſeos rufis, elytris ſulcatis, apice truncatis, angulato-ſerratis.

Habitat in America meridionali. Muſ D. Lund.

Nimis affinis praecedenti. Caput atrum, macula frontali rufa. Thorax laeuis, ater, margine ſanguineo. Elytra ſulcata, atra, macula baſeos ſanguinea, apice truncata, ſerrata, dente exteriori maiori, angulato.

*bidens.* 29. H. ferruginea, thoracis linea laterali nigra, elytris apice vnidentatis. Ent. ſyſt. 2. 73. 11. *

Habitat Surinami. Muſ. D. Boſc.

*nigricornis.* 30. H. glabra, coccinea, antennis nigris. Ent. ſyſt. 2. 73. 12. *

Habitat ad Cap. Bon. Spei. Muſ. D. de Paykull.

*vittata.* 31. H. nigra, elytris obſcure aeneis: vitta rufa.

Habitat in Carolina. Muſ. D. Boſc.

Statura praecedentium. Caput obſcure rufum, antennis atris. Thorax obſcure aeneus, nitens, lateribus rufeſcentibus. Elytra ſtriata punctataque, obſcure aenea, nitidula: vitta lata, fuſca. Corpus nigrum.

32. H.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

32. H. coccinea, elytris nigris: lineola baſeos faſcia- *ſulcata.*
que poſtica coccineis. Ent. ſyſt. 4. App. 448. 13. *

Habitat Surinami. Muſ. D. Holthuyſen.

33. H. atra, thoracis margine pallido, elytris rotun- *rotundata.*
datis, integris.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura reliquis longior. Antennae compreſſae, vix tamen ſerratae. Thorax planus, laeuis, ater, margine pallido. Elytra laeuia, nigra, margine parum pallido, apice rotundata, integra.

34. H. ferruginea, elytris baſi vtrinque gibbis, apice *gibba.*
angulato-dentatis, coeruleis. Ent. ſyſt. ſuppl. 116. 13. *

*Hiſpa gibba.* Oliv. Inſ.

Habitat in St. Domingo. Muſ. D. Boſc.

35. H. atra, thorace maculaque elytrorum baſeos *integra.*
rufis, elytris integris.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Corpus paruum, atrum. Thorax rotundatus, rufus, immaculatus. Elytra ſulcata: ſulcis punctatis, macula magna, oblonga, baſeos rufa, apice omnino integra. Corpus nigrum, femoribus baſi rufis.

6. H. glabra, laeuis, ferruginea, elytris cyaneis. *cyanipennis*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo H. coccineae. Caput coccineum, antennis nigris. Thorax glaber, laeuis, ferrugineus, nitidus. Elytra punctato-ſtriata,



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

cyanea, immaculata. Corpus rufum, pedi geniculis nigris.

*metallica.* 37. H. glabra, nigra, aeneo-nitidula, elytris int gris, punctato-ſtriatis.

Habitat in Carolina. Muſ. D. Boſc.

Reliquis longior et anguſtior. Antennae nigra Caput et thorax vix punctata, nigra, aeneo p rum nitidula. Elytra omnino integra, punct to-ſtriata.

103. CVPES. *Palpi* quatuor aequales: art culo vltimo craſſiori, tru cato.

*Ligula* breuis, membranace bifida.

*Antennae* cylindricae.

*capitata.* 1. CVPES.

Habitat in Carolina. Muſ. D. Boſc.

Characterem generis Dom Latreille debeo. 6 maxillis palpisque. *Palpi* quatuor aequale ſubfiliformes: *anteriores* quadriarticulati: a ticulis ſubaequalibus; vltimo parum craſſio truncato, adhaerentes maxillae dorſo, *poſteri res* triarticulati: articulo ſecundo longiori. nati ligulae baſi. *Mandibula* breuis, cra apice bifida: lacinia exteriore tenui, filiform interiore latâ, rotundata. *Ligula* breuis, me branacea, bifida: laciniis ouatis, rotundatis, ſtantibus. Antennae cylindricae. Genus Hiſp affine.

Corpus elongatum, glabrum, immarginatum, a le, nigrum; *capite* paruo, ouato, exſerto, aequali, rufo; *antennis* thorace multo lo gioribus, approximatis, cylindricis: artic pri

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

primo craſſiori, nigrae, inter oculos inſertae. *Oculi* parui, globoſi, prominuli, marginales. *Thorax* planus, breuis, transuerſus, inaequalis, niger, margine parum prominulo. *Elytra* rigida, fornicata, ſulcata, ſulcis punctis eleuatis, ſcabris, longitudine abdominis, atra. *Pedes* breues, compreſſi, curſorii, nigri, *tarſis* quadriarticulatis, ruſeſcentibus.

## 104. DRYOPS. *Palpi* quatuor inaequales; *anteriores* ſecuriformes, *poſteriores* filiformes. *Ligula* membranacea, late emarginata. *Antennae* filiformes.

1. D. liuida, fronte maculisque duabus thoracis atris, femoribus poſticis incraſſatis. Ent. ſyſt. 2. 75. 1. * *femorata.*

*Oedemera femoralis* Oliv. Inſ. 50. tab. 1. f. 1.
Habitat in Heluetia. Muſ. D. Olivier.

2. D. obſcura, aenea, immaculata. Ent. ſyſt. 2. 75. 2. * *aenea.*

*Oedemera aenea.* Oliv. Inſ. 50. tab. 1. fig. 3.
Habitat in Americae Inſulis. Muſ. D. Banks.



---

Dryopis corpus elongatum, cylindricum, glabrum, immarginatum, laeue, agile; *capite* ouato, infetto, ore palpisque prominulis; *oculis* magnis, globoſis, lateralibus, antennis diſtantibus ſub oculo inſertis, corpore breuioribus; *thorace* ſubcylindrico, breui; *ſcutello* paruo, rotundato; *elytris* rigidiuſculis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, tenuibus, curſorius; *tarſis* quadriarticulatis; *colore* vario, ſubnitidulo.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*liuida.* 3. D. liuida, oculis nigris. Ent. syst. 2. 75. 3.
*Oedemera liuida.* Oliv. Inf. 50. tab. 1. fig. 2
Habitat in Insulis maris australis. Mus. D. Bank

*lineata.* 4. D. liuida, thorace lineis, elytris vitta fusca. En
syst. 2. 75 4. *
*Oedemera lineata* Oliv. Inf. 50. tab. 1. fig. 4
Habitat in noua Zelandia. Mus. D. Banks.

*cyanea.* 5 D. coerulea, antennis fuscis. Ent. syst. 2. 75. 5.
*Oedemera cyanea.* Oliv. Inf. 50. tab. 1. fig.
Habitat in noua Hollandia. Mus. D. Banks.

*marginata.* 6. D. rufa, elytris fuscis: marginibus lineaque m
dia rufis. Ent. syst. 2. 76. 6. ♀
*Oedemera marginata.* Oliv. Inf. 50. tab. 1. f.
Habitat in America meridionali. Mus. Dr. Hunte

*vittata.* 7. D. rufa, elytris vitta abdominisque apice fuscis.
*Oedemera vittata.* Oliv. Inf. 50. tab. 1. fig. 6.
Habitat in Americae Insulis. D. v. Rohr.

*striata.* 8. D. cyanea, thorace rufo, elytris margine striaqu
albidis.
Habitat in Americae Insulis. Mus. D. Lund.
Statura omnino D. rufifrontis, at paullo maior
Caput cyaneum, antennis palpisque fusci
Thorax laeuis, rotundatus, rufus, immacula
tus. Elytra cyanea, margine, sutura striaqu
media, tenuissima, albidis. Corpus et pedes cya
nea.

*rufifrons.* 9. D. cyanea, capite thoraceque rufis. Ent. syst.
76. 8. ♀
Habitat in America boreali.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

105. LAGRIA. *Palpi* inaequales;
*anteriores* ſecuriformes,
*poſteriores* extrorſum craſſiores.
*Ligula* membranacea, integra.
*Antennae* filiformes.

. L. ouata, glabra, atra, elytris tuberculatis. Ent. ſyſt. 2. 78. 1. * — *tuberculata*

Habitat in Cayenna. D. v. Rohr.

. L. villoſa, atra, elytris viridibus. Ent. ſyſt. ſuppl. 108. 1. * — *viridipennis.*

Habitat in Tanger. D. Schousboe.

. L. villoſa, atra, thorace elytrisque viridibus. Ent. ſyſt. 2. 79. 2. * — *villoſa.*

Oliv. Inſ. 48. tab. 1. fig. 1.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

. L. villoſa, fuſca, abdominis margine rubro. — *marginata.*

Habitat in noua Cambria. D. Billardiere.

Statura et magnitudo omnino praecedentium. Corpus totum villoſum, fuſcum, abdominis margine ſolo rufeſcente.

. L. villoſa, nigra, thorace rugoſo. — *rugoſa.*

Habitat in Sumatra. D. Daldorff.

Praecedentibus duplo minor. Caput nigrum. Thorax

---

Lagriae corpus oblongum, hirtum, immarginatum, tardum; *capite* ouato inſerto; *oculis* oblongis, transuerſis, marginalibus; *antennis* diſtantibus, ſub oculos inſertis, longitudine thoracis; *thorace* breui, anguſto, rotundato; *ſcutello* paruo, orbiculato; *elytris* molliuſculis, fornicatis, longitudine abdominis; *pedibus* tenuibus, curſoriis; *tarſis* quadriarticulatis; *colore* vario, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

rax rotundatus, villoſus, rugoſus, niger. Ely-
tra nigra, coeruleo parum micantia.

*pubeſcens.* 6. L. villoſa, nigra, thorace tereti: puncto medio impreſſo, elytris teſtaceis. Ent. ſyſt. 2. 79 3
*Chryſomela pubeſcens.* Linn. ſyſt. Nat. 2. 603 122. *Mus L. hirtae*
Habitat in Europae plantis.

*lata.* 7. L. villoſa, atra, elytris rugoſis, teſtaceis.
Habitat in Mauritania. D. Schousboe. Muſ. D Lund. *Lusitania*
Maior L. pubeſcente et alia. Caput et thorax villoſa, nigra. Elytra valde rugoſa, pallida, teſtacea, immaculata. Corpus atrum.

*hirta.* 8. L. villoſa, nigra, thorace tereti: elytris pallidis Ent. ſyſt. 2. 79. 4.
*Chryſomela hirta.* Linn. ſyſt. Nat. 2. 602. 119 Fn. Sv. 578.
*Tenebrio villoſus.* Degeer Inſ. 5. 44. 6. tab. 2 fig. 23. 24.
Payk. Fn. Sv. 2. 155. 1. *foem. L. pubeſc.*
Oliv. Inſ. 48. tab. 1. fig. 2. *ventr.*
Geoff. Inſ. 1. 344. 6.
Habitat in Europae Saponariis. Muſ. D. Lund.

*tomentoſa.* 9. L. villoſa, cyanea, abdomine femoribusque rufis Ent. ſyſt. 2. 79. 6. *
Habitat in noua Hollandia. Muſ. D. Banks.
Forte potius Melyris, mihi haud rite nota.

*obſcura.* 10. L villoſa, cuprea, elytris nigris. Ent. ſyſt. 2 - 79. 7. *
Habitat in Guinea. D. Iſert.

*triſtis.* 11. L. ſubuilloſa, obſcura, elytris punctatis, antennis nigris.
Habitat Tranquebariae. D. Daldorff. Muſ. D. d Seheſtedt.

Sti

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo L. lugubris, tota obſcura, minime nitida. Antennae nigrae. Tibiae teſtaceae.

12. L. glabra, aenea, elytris ſtriatis, nitidis, cupro-ſtriatis. *nitidula.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Media. Caput et thorax obſcure aenea, vix pubeſcentia, antennis nigris. Elytra nitida, aenea, cupro-lineata. Corpus aeneum, pedibus cupreis.

13. L. glabra, ferruginea, capite thoraceque obſcurioribus, antennis apice nigris. Ent. ſyſt. 2. 80. 8. * *lugubris.*

Habitat in Guinea. D. Iſert.

14. L. liuida, pedibus flaueſcentibus. Ent. ſyſt. ſuppl. 108. 15.* *liuida.*

Habitat in Germania.

106. DASYTES. *Palpi* inaequales, filiformes.
*Maxilla* vnidentata.
*Ligula* membranacea, ſubemarginata.
*Antennae* ſerratae.

1. D. hirtus, ater. *ater.*

La-

---

Daſytae corpus paruum, elongatum, immarginatum, ſaepius hirtum, tardum; *capite* paruo, ouato, inſerto; *oculis* rotundatis, marginalibus, prominulis: *antennis* diſtantibus, thorace longioribus, ante oculos inſertis; *thorace* conuexo, vix marginato, fere latitudine elytrorum; *ſcutello* rotundato, breui; *elytris* molliuſculis, fornicatis, longitudine abdominis; *pedibus* tenuibus, longiuſculis, curſoriis; *tarſis* quinquearticulatis; *colore* vario, interdum nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Lagria atra.* Ent. syst. 2. 80. 10.
*Dermestes hirtus.* Linn. syst. Nat. 2. 563. 14.
*Melyris atra.* Oliv. Inf. 21. tab. 2. fig. 8.
~~Panz. Fn. Germ. 9. tab. 9.~~
Habitat in Europae australioris Tritico.

*Os* maxillis palpisque. *Palpi* quatuor, inaequales, filiformes: articulo vltimo obtuso, subtruncato, *anteriores* paullo longiores, quadriarticulati, articulo secundo longiore, adhaerentes maxillae dorso, *posteriores* breuiores, triarticulati: articulis subaequalibus, adnati ligulae basi laterali. *Mandibula* cornea, arcuata, edentula, acuta. *Maxilla* membranacea, rotundata, vnidentata. *Ligula* membranacea, plana, subemarginata. *Labium* breue, corneum, transuersum, integrum.

*scutellaris.* 2. D. niger, cinereo-hirtus, scutello albo.
*Lagria scutellaris.* Ent. syst. 2. 80. 9. *
Habitat in Hispania. D. Prof. Vahl.

*haemorrhoidalis.* 3. D. hirtus, ater, elytris macula apicis rufa.
*Lagria haemorrhoidalis.* Ent. syst. suppl. 118. 10. *
Habitat in Tanger, D. Schousboe.

*niger.* 4. D. pilosus, niger, elytris laeuiusculis.
*Lagria nigra.* Ent. syst. 2. 80. 11.
*Dasytes niger.* Payk. Fn. Sv. 2. 156. 1.
VIII. 24. *Melyris nigra.* Illig. Col. Bor. 1. 309. 2.
*Melyris villosa.* Oliv. Inf. 21. tab. 2. fig. 10.
Habitat in Europae floribus.

*lineatus.* 5. D. pilosus, niger, elytris lineis sesquitertiis rufis.
*Lagria lineata.* Ent. syst. 2. 80. 12. *
*Melyris lineata.* Oliv. Inf. 21. tab. 1. fig. 6.
Habitat in Brasilia. Mus. D. Banks.

6. D.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. D. villoſus, nigricans, antennarum baſi pedibusque teſtaceis. *flauipes.*
*Lagria flauipes.* Ent. ſyſt. 2. 80. 13. *
*Daſytes flauipes.* Payk. Fn. Sv. 2. 158. 4.
~~Panz. Fn. Germ. 6. tab. 11.~~ *Melyris plumbea*
Habitat in Europa boreali. Dr. Schulz. *Illig. K. Pr.*

. D. ſubuilloſus, cyaneus, antennis nigris. *coeruleus*
*Lagria coerulea.* Ent. ſyſt. 2. 81. 14.
~~*Melyris coerulea.* Oliv. Inſ. 21. tab. 2. fig. 9.~~
*Melyris coerulea.* Illig. Col. Bor. 1. 308. 1.
*Daſytes coeruleus.* Payk. Fn. Sv. 2. 157. 2. *VIII. 24.*
~~*Cicindela.* Geoff. Inſ. 1. 177. 14.~~
Habitat in Europa auſtraliori.

. D. hirtus, obſcure aeneus. *metallicus.*
*Lagria metallica.* Ent. ſyſt. 2. 81. 15. *
Habitat in Barbariae floribus compoſitis. D. Vahl.

. D. villoſus, aeneus, antennis pedibusque nigris. *aeneus.*
*Lagria aenea.* Ent. ſyſt. ſuppl. 419. 15. *
Habitat in Tanger. D. Schousboe.

o. D. hirtus, aeneus, nitidus, antennis nigris, tibiis teſtaceis. *nigricornis.*
*Lagria nigricornis.* Ent. ſyſt. 2. 81. 16. *
*Daſytes nigricornis.* Payk. Fn. Sv. 2. 158. 3.
Habitat in Suecia. Dom. de Paykull.

1. D. obſcurus, thorace rotundato, canaliculato. *linearis.*
*Daſytes linearis.* Payk. Fn. Sv. 3. App. 454. 2.
*Ti[illegible]us filiformis.* Creutz. Inſ. 121. tab. 3. f. 25.
Habitat in Suecia. Muſ. D. Lund. *VI. 16*
Corpus paruum, elongatum, filiforme, totum obſcurum, antennis pedibusque nigris. Thorax magis rotundatus, quam in reliquis, linea dorſali impreſſa.

12. D.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*minutus.* 12. D. ſubuilloſus, niger, thorace elytrisque cyanei
*Lagria minuta.* Ent. ſyſt. 2. 81. 17. *
Habitat in noua Zelandia. Muſ. D. Banks.

107. CEROCOMA. *Palpi* aequales, filiformes
*Maxilla* linearis, integra
*Ligula* membranacea, bi fida.
*Antennae* moniliformes irregulares.

*Schaefferi.* 1. C. viridis, antennis pedibusque luteis. Ent. ſyſt 2. 81. 1.
*Meloe Schaefferi.* Linn. ſyſt. Nat. 2. 681. 12
*Cerocoma* Geoff. Inſ. 1. 358. 1.
Schaeff. Elem. tab. 37.
VIII. 24. - - Icon. tab. 53. fig. 8. 9.
Oliv. Inſ. 48. tab. 1. 1. fig. 1.
Sulz. Hiſt. Inſ. tab. 7. fig. 13.
Habitat in Europa auſtrali.

*Vahlii.* 2. C. viridis. antennis pedibusque nigris. Ent. ſyſt 2. 82. 2. * Est facies C. Schreberi
Habitat in Barbariae vmbellatis. D. Vahl.

*Schreberi.* 3. C. viridis, antennis, pedibus abdominisque ſegmentis tribus flauis. Ent. ſyſt. 2. 82. 3. *
Oliv. Inſ. 48. tab. 1. fig. 2.
Habitat in Europa auſtrali.

10

Cerocomae corpus oblongum, laeue, pubeſcens, immargin tum, tardum; *capite* ouato, promiuulo, inflexo; *oc* ouatis, marginalibus; *antennis* approximatis, thorace br uioribus, inter oculos inſertis; *thorace* breui, ouato, ſ pra plano; *ſcutello* breuiſſimo, rotundato; *elytris* incu bentibus, molliuſculis, longitudine abdominis; *pedi* tenuibus, curſoriis; *tarſis* anticis quinque-, poſticis quad articulatis; *colore* aeneo.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

## 108. PAVSSVS. *Palpi* quatuor, inaequales. *Antennae* biarticulatae: articulo vltimo maximo.

1. P. antennis biarticulatis: claua irregulari dentata maxima, corpore fuſco. ***microcephalus.***

*Cerocoma microcephala.* Ent. ſyſt. 2. 82. 4.
***Pauſſus microcephalus.*** Linn. Diſſ. Big. Inſ. tab. 1. fig. 6-10.
***Pauſſus microcephalus.*** Thunb. Act. Sv. 1781. 170. 1.
***Pauſſus microcephalus.*** Afzel. Act. Soc. Linn. 4. 18. tab 22.
Habitat in Africa. Muſ. D. Banks.
Nouum genus et diſtinctum, at mihi haud rite examinatum, nec mihi ſatisfacit character generis a D. Afzelio in Act. Soc. Linn. datus, ſed e nouo conficiendus.

2. P. antennis biarticulatis: claua elongata integra, fuſcus, elytris linea nigra. ***lineatus.***

*Cerocoma lineata.* Ent. ſyſt. 2. 82. 5. *
***Pauſſus lineatus.*** Thunb. Act. Soc. Suec. 1781. 170. 3. tab. 1. fig. 4. 5.
Habitat ad Cap. Bon. Spei.

3. P. antennis biarticulatis: claua elongata integra, niger, thorace elytrorumque ſtriga ferrugineus. ***ruficollis.***

*Cerocoma ruficollis.* Ent. ſyſt. 2. 83. 6. *
Habitat - - Muſ. D. Lund.

4. P. antennarum articulo primo incuruo, ſecundo rotundato, compreſſo, flauis; corpore nigro. ***flauicornis.***

Habitat in Iaua. Muſ. D. de Seheſtedt.
Animalculum ſingulare vix huius generis. Corpus paruum, nigrum. Caput et thorax cinereo-villoſa. Antennae articulo primo, incuruo, elongato,

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

gato, secundo magno, compresso, flauis, reliquis breuibus, filiformibus, nigris. Elytra cyanea, nitida.

109. LYTTA. *Palpi* quatuor, inaequales; *posteriores* clauati.
*Maxilla* bifida.
*Labium* truncatum.
*Antennae* filiformes.

*vesicatoria.* 1. L. viridis, antennis nigris. Ent. syst. 2. 83. 1.
*Meloë vesicatorius.* Linn. syst. Nat. 2. 679. 3. Fn. Sv. 827.
*Cantharis vesicatoria.* Oliv. Inf. 46. tab. 1. f. 1.
Geoff. Inf. 1. 341. 1. tab. 6. fig. 5.
Degeer. Inf. 4. 12. 2. tab. 1. fig. 9.
Schaeff. Icon. tab. 47. fig. 1.
*IV. 54.* Panz. Fn. Germ. 41. tab. 4.
Sulz. Hist. Inf. tab. 7. fig. 55.
Habitat in Europae Ligustro, Fraxino, Sambuco.
Officinalis pro vesicatoriis.

*segetum.* 2. L. aurea, elytris viridibus. Ent. syst. 2. 84. 2. *
Habitat in Barbariae segete. Mus. D. Desfontaines.

*nitidula.* 3. L. virídi-aenea, elytris testaceis. Ent. syst. 2. 84. 3. *
Oliv. Inf. 46. tab. 2. fig. 15.
Habitat ad Cap. Bon. Spei.

4. L.

---

Lyttae corpus oblongum, laeue, glabriusculum, immarginatum, tardum; *capite* magno, exserto, ouato, thorace latiore; *oculis* rotundatis, lateralibus; *antennis* distantibus, cantho interiori oculorum insertis; *thorace* subcylindrico; *scutello* minuto; *elytris* molliusculis, fornicatis, longitudine abdominis; *pedibus* elongatis, tenuibus, cursoriis; *tarsis* anticis quinque-, posticis quadriarticulatis; *colore* vario, nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. L. atra, vertice, thorace pedibusque ferrugineis, elytris cyaneis. Ent. ſyſt. 2. 84. 4. * *collaris.*
Oliv. Inſ. 46. tab. 2. fig. 12.
Pall. Inſ. Sib. 96. tab. E. fig. 27.
Habitat in Roſſia meridionali. D. Boeber.

. L. cyanea, pectore ferrugineo. Ent. ſyſt. 2. 84. 5. * *Gigas.*
Oliv. Inſ. 46. tab. 1. fig. 9.
Habitat in Guinea. D. Iſert.

. L. aeneo-villoſa, elytrorum marginibus albis. Ent. ſyſt. ſuppl. 119. 15. * *villoſa.*
Habitat in Guinea. Muſ. D. Lund.

. L. teſtacea, elytris coccineis. *coccinea.*
Habitat in Guinea. Muſ. D. de Seheſtedt.
Statura magna praecedentis. Caput teſtaceum, antennis nigris: articulo primo teſtaceo. Thorax teſtaceus. Elytra laeuia, coccinea, immaculata. Corpus teſtaceum, pedibus apice nigris.

. L. villoſa, viridi-coerulea, thorace rotundato, ferrugineo. Ent. ſyſt. 2. 84. 6. * *Syriaca.*
*Meloë Syriacus.* Linn. ſyſt. Nat. 2. 680. 4.
Muſ. Lud. Vlr. 102.
*Cantharis Syriaca.* Oliv. Inſ. 46. tab. 1. fig. 5.
Panz. Fn. Germ. 41. tab. 5.
~~Herbſt. Arch. tab. 30. fig. 1.~~
Habitat in Europa auſtraliori frequens.

. L. cinereo-villoſa, capite maximo atro, thorace rufo. *melanocephala.*
Habitat in Guinea. D. Meier. Muſ. D. Lund.
Maior L. ruficolli. Caput magnum, laeue, glabrum, atrum. Thorax antice attenuatus, laeuis, rufus. Elytra et corpus villis paruis cinereis tecta. Pedes obſcuri.

10. L.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*2pustulata.* 10. L. atra, capitis vertice thoracisque maculis duabus rufis.

Habitat in Guinea. Muſ. D. de Seheſtedt.

Statura praecedentium. Antennae breues, craſſae, ſerratae. Caput atrum, ore verticeque late rufis. Thorax ſubrotundus, in medio parum canalieulatus, ater, vtrinque macula magna, rufa. Elytra ſubrugoſa, atra. Corpus nigrum.

*rufa.* 11. L. nigra, capitis vertice, thoracis maculis duabus elytrisque rufis: vitta abbreuiata nigra.

Habitat in Guinea. Muſ. D. Lund.

Statura omnino praecedentis. Antennae breues, craſſae, atrae. Caput nigrum, vertice late rufo. Thorax ſubrotundatus, niger, maculis duabus rufis. Elytra ſubrugoſa, rufa: vitta nigra, quae tamen nec baſin nec apicem attingit. Corpus nigrum.

*Afzeliana.* 12. L. thorace teſtaceo: maculis quatuor, elytris macula, vitta abbreuiata apiceque nigris.

Habitat in Carolina. Muſ. D. Boſc.

Magna. Caput teſtaceum, ore antennisque nigris. Thorax laeuis, teſtaceus, vtrinque lunula antica punctisque duobus dorſalibus nigris. Elytra laeuia, teſtacea; macula magna baſeos ad ſuturam, vitta magna poſtica dilatata, quae cum apice nigro coit, nigris. Corpus nigrum, marginibus ſegmentorum abdominis teſtaceis.

*ruficollis.* 13. L. glabra, viridi-aurea, thorace antice attenuato rufo, Ent. ſyſt. 2. 85. 7. ♀

*Cantharis ruficollis.* Oliv. Inſ. 46. tab. 1. f. 6.

~~Pall. Inſ. 3. tab. E. fig. 85.~~

~~Herbſt. Arch. tab. 48. fig. 4.~~

Habitat in India orientali. D. Olivier.

14. L

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

14. L. ſupra teſtacea, elytris macula magna, apicis oblonga, nigra. Ent. ſyſt. 2. 85. 8. * *teſtacea.*
Oliv. Inſ. 46. tab. 2. fig. 13.
Habitat Tranquebariae. Muſ. D. de Seheſtedt.

15. L. viridi-aenea, nitida, elytris teſtaceis: maculis viridi-aeneis. Ent. ſyſt. 2. 85. 9. * *feſtiua.*
Oliv. Inſ. 46. tab. 2. fig. ~~14~~ 15
Pall. Iter. 2. 721. 54.
Habitat in Sibiria. Muſ. D. Banks.

16. L. atra, elytrorum marginibus cineraſcentibus. Ent. ſyſt. 2. 85. 10. * *marginata.*
*Meloë cinereus.* Forſt. nou. ſp. Cent. 1. 62. 62.
*Cantharis marginata.* Oliv. Inſ. 46. tab. 1. f. 2.
Habitat ad Cap. Bon. Spei.

17. L. teſtacea, thorace vittis duabus, elytris tribus atris. *lemniſcata.*
Habitat in Carolina. Muſ. D. Boſc.
Media. Antennae obſcurae. Caput teſtaceum, punctis duobus nigris. Thorax rotundatus. teſtaceus, vittis duabus nigris. Elytra villoſa, teſtacea, vittis tribus atris, quae tamen apicem haud attingunt. Corpus aureo-villoſum, pedibus teſtaceis.

18. L. elytris nigris: vitta marginibusque flauis. Ent. ſyſt. 2. 86. 11. * *vittata.*
*Cantharis vittata.* Oliv. Inſ. 46. tab. 1. fig. 3.
~~Pall. Inſ. tab. E. fig.~~ 53.
Habitat in Americae Solano tuberoſo, folia depaſcens, deſtruens.

19. L. corpore atro, immaculato. Ent. ſyſt. 2. 86. 12. * *atrata.*
*Meloë Penſylvanicus.* Degeer Inſ. 5. 16. 1. tab. 13. fig. 1.

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf 46. tab. 2. fig. 19.
Habitat in Barbaria. D. Prof. Vahl.

*cinerea.* 20. L. corpore nigro, cinereo, tomentoſo. Ent. ſyſt. ſuppl. 119. 15. *
*Cantharis ſericea.* Oliv. Inf. 46. tab. 1. fig. 8.
Habitat in America. D. Hybner.

*erythrocephala.* 21. L. atra, capite teſtaceo, thorace elytrisque cinereo-lineatis. Ent. ſyſt. 2. 86. 13. *
*Meloë erythrocephalus.* Pall. Iter. 1. 466. 46
Oliv. Inf. 46. tab. 2. fig. 16.
~~Panz. Fn. Germ. 51. tab. 6.~~
Herbſt. Arch. tab. 30. fig. 2.
Scopol. Ann. Hiſt. natur. 5. 103. 81.
Habitat in ~~Auſtria. Muſ. Dr. Schulz.~~

*oculata.* 22. L. nigra, puncto calloſo, flauo pone oculos. Ent. ſyſt. 2. 86. 14. *
Habitat in Guinea. D. Iſert.

*dubia.* 23. L. atra, capitis vertice fuluo, thorace elytrisque immaculatis. Ent. ſyſt. 2. 86. 15. *
~~*Meloë Algiricus* Sulz. Hiſt.~~ Inſ. tab. 7. fig. 12.
~~Oliv. Inf. 46. tab. 1. fig. 7.~~
Pall. Iter. tab. E. fig. 29.
Habitat in Sibiria. Muſ. D. Banks.

*Afra.* 24. L. nigra, thorace rufo. Ent. ſyſt. 2. 87. 16. *
*Meloë Afer* Linn. ſyſt. Nat. 2. 680. 10.
*Cantharis Afra.* Oliv. Inf. 46. tab. 1. fig. 4.
~~Roſſ. Fn. Etr. 240. 594. tab. 3. fig. 1. 3.~~
Habitat in Africa.

*craſſicornis.* 25. L. atra, thorace elytrisque teſtaceis, antennis incraſſatis.
Habitat in Auſtria. Muſ D. Lund.
Statura omnino praecedentium. Antennae breues. craſſae, nigrae. Thorax et elytra laeuia, teſtacea, immaculata. Corpus atrum.

26. L.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

6. L. coerulèo-nigricans, abdomine poſtice rufo. Ent. ſyſt. 2. 87. 17. * *haemorrhoidalis.*
Habitat ad Cap. Bon. Spei. D. de Paykull.

110. MYLABRIS. *Palpi* filiformes.
*Maxilla* cornea, compreſſa, bifida.
*Ligula* membranacea, ſubemarginata.
*Antennae* moniliformes.

1. M. nigra, elytris faſcia media flaua. Ent. ſyſt. 2. 87. 1. *faſciata.*
Habitat in Indiis. Dr. Torſtenſon.

2. M. nigra, elytris flauis: faſciis tribus nigris. Ent. ſyſt. 2. 88. 2. *Cichorei.*
*Meloë Cichorei.* Linn. ſyſt. Nat. 2. 680. 5. *III. 41.*
Muſ. Lud. Vlr. 103.
*Mylabris Fueslini.* Panz. Fn. Germ. 31. tab. 18.
Oliv. Inſ. 47. tab. 1. fig. 1.
~~Sulz. Hiſt. Inſ. tab. 7. fig. 11.~~
Degeer Inſ. 5. 17. 2. tab. 13. fig. 2.
Pall. Icon. tab. E. fig. 7.
Habitat in totius Orientis Cichoreis, in China officinalis.

3. M.

---

Mylabridis corpus oblongum, laeue, ſubpubeſcens, immarginatum, tardum; *capite* magno, exſerto, ouato, inflexo; *oculis* rotundatis, marginalibus; *antennis* thorace breuioribus, diſtantibus, ante oculos inſertis; *thorace* rotundato; *ſcutello* paruo, quadrato; *elytris* rigidiuſculis, fornicatis, abdomine longioribus; *pedibus* longiuſculis, curſoriis; *tarſis* anticis quinque-, poſticis quadriarticulatis; *colore* obſcuro.

Antennarum apex in Mylabridibus, ſicut in Necrophoris, probabiliter pro ſexu variat.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*3faſciata.* 3. M. atra, antennis elytrisque flauis, elytris faſcii duabus apiceque nigris.
*Mylabris 3faſciata* Oliv. Inſ. 47. tab. 1. f. 8
Habitat in Guinea. Muſ. D. Lund.
Statura M. Cichorei. Antennae magnae, flauae apice nigrae. Caput et thorax atra, immacula ta. Elytra laeuia, flaua, faſciis duabus ſimpli cibus apiceque late atris.

*lunata.* 4. M. atra, antennis ferrugineis, elytris flauis: ma cula faſciisque tribus atris; anticis interrupti
*Mylabris lunata.* Oliv. Inſ. 47. tab. 1. fig. 2.
Pall. Inſ. tab. E. fig. 5.
Habitat Tranquebariae. D. Daldorff. Muſ. D. d Seheſtedt.
Maior M. Cichorei. Caput et thorax villoſa, atr immaculata. Antennae apice ferrugineae. Ely tra flaua, macula communi ſcutellari, ſcutell que atris. Praeterea faſcia baſeos futuram hau attingens, at ad marginem exteriorem vsque a baſin dilatata: macula marginali flaua. Faſc ſecunda pone medium abbreuiata, at punct futurali atro. Apex late ater.
*Variat* rarius faſciis integris.

*praeuſta.* 5. M. atra, elytris apice teſtaceis: macula atra. En ſyſt. 2. 88. 3. *
Habitat in Barbariae Segete. Muſ. D. Desfontaine

*margina-ta.* 6. M. atra, elytrorum margine ſanguineo. Ent. ſyſt 2. 88. 4. *
Habitat in Barbariae Segete. Muſ. D. Desfontaine

*Algirica.* 7. M. atra, elytris teſtaceis immaculatis. Ent. ſyſt 2. 88. 5.
*Meloë Algiricus* Linn. ſyſt. Nat. 2. 651. 11.
*Lytta Indica.* Herbſt. Arch. tab. 30. fig. 3.
Oliv. Inſ. 47. tab. 1. fig. 5.

Wu

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Wulf. Inf. Cap. tab. 1. fig 8.
Cyrill. Ent. Neap. 1. tab. 2. fig. 10.
Habitat in India.

8. M. nigra, elytris maculis ſex flauis: prima arcuata. Ent. ſyſt. 2. 88. 6. * *Capenſis.*
*Meloë Capenſis.* Linn. ſyſt. Nat. 2. 650. 7. Muſ. Lud. Vlr. 104.
Oliv. Inſ. 47. tab. 2. fig. 12.
Wulf. Inſ. Cap. tab. 1. fig. 5.
Habitat ad Cap. Bon. Spei.

9. M. villoſa, nigra, elytris macula baſeos faſciisque duabus dentatis, flauis. Ent. ſyſt. 2. 89. 7. * *Hermanniae.*
Habitat in Guineae Hermanniis. D. Iſert.

10. M. villoſa, nigra, elytris maculis duabus baſeos faſciisque duabus dentatis, ferrugineis, antennis nigris. Ent. ſyſt. ſuppl. 120. 7. * *Sidae.*
Oliv. Inſ. 47. tab. 2. fig. 11.
Thunb. nou. ſp. Inſ. 6. 114. tab. E. fig. 14.
Pall. Inſ. tab E. fig. 3.
Habitat ad Cap. Bon. Spei.
Caput et thorax ſubuilloſa, atra, immaculata. Elytra punctata, linea vna alteraue eleuata, maculis duabus baſeos faſciisque duabus ſinuatis flauis.

11. M. hirta, nigra, elytris macula baſeos faſciisque duabus ferrugineis. *Lavaterae.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.
Summa affinitas praecedentis, at corpus magis hirtum. Elytra laeuiora, nitidiora, macula tantum vnica baſeos faſciisque minus ſinuatis.

12. M. holoſericea, atra, elytris punctis ferrugineis. *atrata.*
Oliv. Inſ. 47. tab. 2. fig. 15. at puncta elytrorum plura.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. Muſ. D. de Seheſtedt.

Corpus reliquis paullo latius, parum holoſericeum, atrum, puncto vno alteroue elytrorum ferrugineis.

*punctum.* 13. M. villoſa, atra, elytris faſciis duabus: anteriore flaua; puncto nigro, poſteriore rufa. Ent. ſyſt. 2. 89. 8. *

Oliv. Inſ. 47. tab. 2. fig. 19.

Habitat Tranquebariae. Muſ. D. Lund.

*10punctata* 14. M. atra, elytris teſtaceis: punctis quinque nigris. Ent. ſyſt. 2. 89. 9. *

*Meloë 4punctatus.* Linn. ſyſt. Nat. 2. 680. 6.

Oliv. Inſ. 47. tab. 1. fig. 4.

Habitat in Italia. Dr. Allioni.

*4punctata* 15. M. atra, elytris teſtaceis: punctis quatuor nigris. Ent. ſyſt. 2. 89. 10. *

Habitat in Ruſſia. Muſ. D. Banks.

*6maculata* 16. M. atra, elytris teſtaceis: punctis tribus nigris. Ent. ſyſt. ſuppl. 120. 9. * Hungaria

Habitat in Ruſſia meridionali. D. Boeber.

*flauicornis* 17. M. antennis ferrugineis, atra, elytris faſciis ſubtribus maculisque duabus apicis flauis.

Habitat ad Cap. Bon. Spei. Muſ. D. de Seheſtedt.

Parua. Antennae fere totae ferrugineae. Caput et thorax villoſa, atra. Elytra atra, macula baſeos cum faſcia anteriore coëunte, faſciis duabus mediis dentatis maculisque duabus apicis flauis.

*geminata.* 18. M. atra, nitida, elytris pallidis: punctis duarum parium faſciaque media nigris. Ent. ſyſt. ſuppl. 120. 9. *

Habitat in Ruſſia meridionali. D. Boeber.

*ruficornis.* 19. M. atra, antennis ferrugineis, elytris pallidis: pun-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

punctis tribus faſciisque duabus atris. Ent. ſyſt. ſuppl. 121. 10. *
Habitat in Mogador Africae. D. Schousboe.

20. M. nigra, coleoptris flauis: faſcia punctoque communi fuſcis. Ent. ſyſt. 2. 89. 11. * — *3 maculata*
Cyrilli Ent. Neap. 1. tab. 3. fig. 7.
Habitat in Oriente. D. Prof. Forſkahl. *Hungaria*

21. M. hirta, nigra, elytris, faſcia poſtica punctoque apicis flauis. Ent. ſyſt. ſuppl. 121. 12. * — *minuta.*
Habitat ad Cap. Bon. Spei. D. Weber.

22. M. villoſo-argentea, elytris maculis flaneſcentibus: plerisque margine connexis. Ent. ſyſt. 2. 90. 12. * — *argentata.*
Habitat in Senegallia. D. de Paykull.
Parua, antennis ferrugineis.

## LII. HORIA.

*Palpi* filiformes.
*Maxilla* vnidentata.
*Ligula* bifida: laciniis linearibus, diſtantibus.
*Antennae* moniliformes.

1. H. flaneſcens, elytris maculis ſeptem nigris. Ent. ſyſt. 2. 90. 1. * — *maculata.*

*Horia*

---

Horiae corpus elongatum, glabrum, immarginatum, agile; *capite* magno, ouato, plano, exſerto, mandibulis palpisque prominulis; *oculis* magnis, oblongis, transuerſis, prominulis, marginalibus; *antennis* diſtantibus, thorace longioribus, ante oculos inſertis; *thorace* plano, ſubquadrato, latitudine fere elytrorum; *ſcutello* porrecto, tenui; *elytris* rigidiuſculis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, curſoriis, femoribus poſticis interdum incraſſatis; *tarſis* anticis quinque-, poſticis quadriarticulatis; *colore* nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Horia maculata.* Oliv. Inf. 53. tab. 1. fig. 1.
*Cucujus maculatus.* Swed. Act. Holm.
Habitat in Cayenna. D. Olivier.

*teſtacea.* 2. H. rufa, antennis pedibusque nigris. Ent. ſyſt. 2 91. 2. *
Oliv. Inf. 53. tab. 1. fig. 2.
Hybn. Naturf. 24. 47. 13. tab. 2. fig. 14-17.
Habitat Tranquebariae.

*maxilloſa.* 3. H. ſanguinea, antennis pedibusque atris, capite maximo: mandibulis exſertis, longitudine capitis.
*Horia cephalotes* Oliv. Inf. 53. tab. 1. fig. 3.
Habitat in Sumatra. D. Daldorff.
Statura et ſumma affinitas praecedentis. Antennae breues, atrae. Caput maximum, planum ſanguineum, mandibulis exſertis, forſicatis, interne ante apicem vnidentatis longitudine capitis. Thorax transuerſus, ſanguineus. Elytra ſanguinea. Pedes ſimplices, atri.
Aliam vidi ſimillimam, thorace itidem transuerſo pedibusque ſimplicibus, at capite mediocri mandibulisque haud exſertis, an mas?



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

112. LYMEXYLON. *Palpi anteriores* porrecti: articulo penultimo magno; appendiculo acuto, filfo, vltimo ouato, acuto. *Antennae* filiformes.

1. L. teſtaceum, oculis, alis pectoreque nigris. Ent. ſyſt. 2. 91. 1. * *dermeſtoides.*
*Cantharis dermeſtoides.* Linn. ſyſt. Nat. 2. 650. 25. Fn. Sv. 702.
*Lytta Francofurtana.* Herbſt. Arch. tab. 30. fig. 4.
Oliv. Inf. 25. tab. 1. fig. 1.
Payk. Fn. Sv. 2. 161. 1.
Habitat in Europae ligno.

2. L. nigrum, thoracis linea dorſali flaua, elytris breuiſſimis fornicatis. Ent. ſyſt. 2. 92. 2. * *abbreuiatum.*
*Necydalis breuicornis.* Elytris minutiſſimis, capite thoraceque fuſcis: linea longitudinali flaua, antennis breuiſſimis craſſiuſculis. Linn. ſyſt. Nat. 2. 643. 11.
Habitat in Sierra Leon Africae. Dr. Pflug.

3. L. nigrum, elytris teſtaceis, apice nigris. Ent. ſyſt. 2. 92. 3. *proboſcideum.*
Oliv.

---

Lym.xyli corpus elongatum, cylindricum, ſubpubeſcens, immarginatum, agile; *capite* rotundato, ſupra plano, inflexo, inſerto, latitudine thoracis; *oculis* prominulis, rotundatis, marginalibus; *antennis* diſtantibus, thorace breuioribus, ante oculos inſertis; *thorace* breui, transuerſo, vix marginato; *ſcutello* oblongo, apice rotundato; *elytris* incumbentibus, molliuſculis, longitudine abdominis; *pedibus* brenibus, curſoriis; *tarſis* quinquearticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 25. 6 5. tab. 1. fig. 5.
Habitat in Germania. D. de Hattorff.

*nauale.* 4. L. luteum, capite, item elytrorum margine apiceque nigris. Ent. ſyſt. 2. 92. 5. *
Oliv. Inſ. 25. 5. 4. tab. 1. fig. 4.
*Cantharis naualis.* Linn. ſyſt. Nat. 2. 650. 26. Fn. Sv. 718. Iter Weſtgoth. 155. tab. 2.
Friſh. Inſ. 13. 24. tab. 20.
Schaeff. Icon. tab. 59. fig. 1.
Payk. Fn. Sv. 2. 161. 2.
Habitat in ligno quercus, quod deſtruit.

*flauipes.* 5. L. nigrum, elytris baſi, abdomine apice, pedibusque flauis. Ent. ſyſt. 2. 93. 6. *
Habitat in Saxoniae ligno. Dom. Hybner.

*Morio.* 6. L. nigrum, pedibus anticis flauis. Ent. ſyſt. 2. 93. 7. *
Schaeff. Icon. tab. 279. fig. 7.
Habitat in Saxoniae ligno. D. Hybner.

## 113. DIRCAEA. *Palpi* inaequales; *anteriores* porrecti: articulo vltimo maximo, trilobo.
*Antennae* filiformes.

*barbata.* 1. D. fuſca, antennis tibiisque piceis. Ent. ſyſt. ſuppl. 121. 1. *Lyme-*

---

Dircaeae corpus elongatum, cylindricum, glabrum, immarginatum, tardum; *capite* rotundato, inferto; *palpis* prominulis; *oculis* magnis, oblongis, transuerſis, marginalibus, vix prominulis: *antennis* diſtantibus, corpore breuioribus, ante oculos inſertis: *thorace* ouato, angulo poſtico vtrinque prominulo; *ſcutello* breui, rotundato; *elytris* rigidis, ſtriatis, abdomine vix longioribus: *pedibus* curſoriis; *tarſis* anticis quadri-, poſticis quinquearticulatis; *colore* vario obſcuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Lymexylon barbatum.* Ent. syst. 2. 92. 4. *
*Mordella barbata.* Schall. Act. Hall. 1. 322. tab. 1. fig. 7.
*Serropalpus striatus.* Payk. Fn. Sv. 2. 163. 1.
Hellen. in Act. Holm. 1786. 273. tab. 8. fig. 3-5.
Illig. Col. Bor. 1. 130. 7.
Oliv. Inf. 25. 5. 3. tab. 1. fig. 3.
Schaeff. Icon. tab. 66. fig. 1.
Habitat in Germaniae lignis.
*Os* maxillis palpisque. *Palpi* quatuor, inaequales: *anteriores* porrecti, exserti, multo longiores, quadriarticulati: articulis tribus prioribus breuibus, aequalibus, vltimo elongato, compresso, trilobo: lobo inferiori conico, intermedio tenuiori, subulato, incuruo, superiori maiori acuto, adhaerentes maxillae dorso, *posteriores* breues, filiformes, triarticulati: articulis aequalibus adnati ligulae medio. *Mandibula* breuis, cornea, edentula, incurua. *Maxilla* breuissima, membranacea, bifida: laciniis aequalibus, cylindricis, obtusis. *Ligula* breuis, membranacea, rotundata, integra. *Labium* breue, transuersum, corneum, emarginatum.

2. D. nigra, elytris obscure testaceis. Ent. syst. suppl. 121. 2. * *discolor.*
*Serropalpus laeuigatus.* Illig. Col. Bor. 1. 131. 8.
*Xylita bupreſtoides.* Payk. Fn. Sv. 2. 49. 1. *II. 33.*
~~*Elater bupreſtoides.* Ent. syst. 2. 234. 54.~~ *IX.*
Hellen. Act. Sv. 7. 274.
Habitat Halae Saxonum. Dom. Hybner.

3. D. atra, elytris maculis quatuor fuluis. Ent. syst. suppl. 122. 3. * *4guttata.*
*Serropalpus 4maculatus.* Illig. Col. Bor. 1. 130. 6.

*Hy-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Hypalus 4guttatus* Payk. Fn. Sv. 1. 251. 1.
Habitat in Suecia. D. de Paykull.

*ruficollis.* 4. D. nigricans, thorace anoque fuluis. Ent. syst suppl. 122. 4. *
Habitat sub arborum corticibus Finlandiae.
De genere haereo; mihi haud rite examinata.

*2fasciata.* 5. D. ferruginea, capite elytrorumque puncto fusc apiceque nigris.
*Notoxus 2fasciatus.* Ent. syst. 1. 210. 3. *
*Serropalpus 2fasciatus.* Illig. Col. Bor. 1. 129. 5
*Hypalus quercinus.* Payk. Fn. Sv. 1. 252. 2.
I. 19. Panz. Fn. Germ. 6. tab. 3.
Habitat in Europa boreali.

*dubia.* 6. D. thorace fusco, elytris ferrugineis: lineola ba seos fascia media punctoque apicis nigris.
*Notoxus dubius.* Ent. syst. 1. 211. 4. *
*Serropalpus dubius.* Illig. Col. Bor. 1. 128. 4.
I. 19. Panz. Fn. Germ. 11. tab. 13.
Habitat in Germaniae ligno. D. Smidt.

*fuluicollis.* 7. D. thorace fuluo, elytris fuscis: maculis duabu albis.
Habitat in Germania. D. Smidt.
Statura et magnitudo omnino D. dubiae. Capu cum antennis nigrum. Thorax laeuis, fuluus immaculatus. Elytra striata, fascia maculi duabus magnis, sublunatis, albidis, et inte maculas lunula parua ad marginem exterio rem. Corpus nigrum, pedibus fuluis.

*variegata.* 8. D. fusca, elytris pallide testaceis, fusco variis.
*Serropalpus variegatus.* Act. Hist. Nat. Paris. 1. tab. 10. fig. 2.
Habitat in Gallia. Mus. D. Bosc.
Statura omnino et magnitudo D. dubiae. Anten nae testaceae. Caput et thorax fusca, immaculata

Ely-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Elytra laeuia, pallide teſtacea, fuſco variegata. Corpus teſtaceum.

9. D. cinereo-pubeſcens, nigra, antennis pedibusque flaueſcentibus. *murina.*

Habitat in Carolina Muſ. D. Boſc.

Duplo minor D. 4guttata. Antennae flaueſcentes, apice obſcuriores. Thorax gibbus, rotundatus, niger, tomento tenui, fugaci, cinereo-aureo parum nitidulo. Elytra ſubſtriata, nigra, tomento eodem fugaci tecta. Pedes flaueſcentes.

** *ſaltatoriae.*

10. D. rufeſcens, thorace maculis duabus nigris. *humeralis.*

*Hallomenus humeralis* depreſſus, rufeſcens, ſericeus, thorace maculis duabus nigris; elytris ſubſtriatis, humeris luteis. Illig. Col. Bor. 1. 134. 1.

*Hallomenus 2punctatus.* Payk. Fn. Sv. 2. 179. 1.

*Chryſomela 2notata.* Querſ. Ign. Inſ. ſp. 1. 14.

Panz. Fn. Germ. 16. tab. 17. II. 69

Habitat in Germaniae Fungis.

Proprii generis videtur, at Hallomeni character adhuc deſideratur. A Dom. de Paykull. datus vix rite elaboratus. Corpus paruum, holoſericeum, vtrinque attenuatum, rufeſcens. Thorax maculis duabus magnis nigris. Capitis vertex nigricat.

11. D. fuſco-holoſericea, antennis extrorſum craſſioribus. *micans.*

*Hallomenus micans.* Illig. Col. Bor. 1. 135. 3. II 70.

*Hallomenus miccans* Payk. Fn. Sv. 2. 181. 2.

*Megatoma picea.* Herbſt. Col. 4. 97. 5. tab. 39. fig. 5.

Panz. Fn. Germ. 17. tab. 17.

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat ad Germaniae ſepes.
Statura omnino praecedentis, at paullo minor.
*Variat* interdum colore magis teſtaceo.

114. CVCVIVS. *Palpi* aequales: articulo vltimo truncato, craſſiori.
*Ligula* breuis, bifida: laciniis linearibus, diſtantibus.
*Antennae* moniliformes.

*Gigas.* 1. C. depreſſus, ater, elytris biſtriatis: macula apicis coccinea.
Habitat in Guinea. Muſ. D. de Seheſtedt.
Corpus in hoc genere maximum, depreſſum, laeue, atrum, nitidum. Thoracis margo muticus. Elytra glabra, atra, ſtriis duabus diſtantibus, impreſſis, et inter has ſtrias, verſus apicem, macula magna rufa.

*heros.* 2. C. depreſſus, ater, femoribus rufis.
Habitat in Sumatra. D. Daldorff. Muſ. D. Lund.
Statura et magnitudo omnino praecedentis. Antennae breues, craſſae, moniliformes, in altero ſexu, articulis ſubtus prominulis, ſerratae. Caput mandibulis craſſis, porrectis fronteque biſulcata. Thorax planus, ater, ſtria vtrinque impreſſa. Elytra atra, immaculata, inaequaliter ſtriata. Corpus nigrum, femoribus rufis.

3. C.

---

Cucuii corpus oblongum, depreſſum, planum, glabrum, immarginatum, agile; *capite* magno, ouato, inſerto, angulo poſtico interdum prominulo; *antennis* diſtantibus, thorace longioribus, pone maudibulas inſertis; *oculis* paruis, rotundatis, prominulis, marginalibus; *thorace* ſubmarginato: margine ſaepius dentato; *ſcutello* breui rotundato; *elytris* rigidis, includentibus, longitudine abdominis; *pedibus* breuibus, validis, curſoriis, femoribus ſubclauatis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

3. C. ater, elytris ferrugineis. *rufipennis.*
Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.
Statura omnino praecedentis, at duplo fere minor, et totus ater, elytris ſolis ferrugineis.

4. C. thorace denticulato elytrisque rufis, pedibus ſimplicibus nigris. Ent. ſyſt. 2. 93. 1. *depreſſus.*
*Cantharis ſanguinolenta.* Linn. ſyſt. Nat. 2. 646. 19.
Friſh. Inſ. 12. tab. 7. fig. 1. VIII. 25.
Payk. Fn. Sv. 2. 165. 1.
Habitat in Germania rarius.

5. C. thorace ſulcato, niger, elytris crenato-ſtriatis. *ſulcatus.*
Habitat in Croatiae ligno putreſcente. D. Saldoner.

6. C. depreſſus, rufus, elytris ſtriatis. Ent. ſyſt. ſuppl. 123. 2. * *rufus.*
Habitat in Carolina. Muſ. D. Boſc.

7. C. thorace ſulcato, elytris punctatis. ater, antennis pedibusque ferrugineis. Ent ſyſt. 2. 94. 3. * *rufipes.*
Habitat in Barbariae ligno. Muſ. D. Desfontaines.

8. C. niger, thorace elytrisque abbreuiatis rufis, mandibulis exſertis, dentatis. *maxilloſus*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura omnino C. ſpinoſi, paullo minor. Antennae magis moniliformes, piloſae. Caput atrum, nitidum, mandibulis craſſis, exſertis, dentatis. Thorax planus, rufus, linea dorſali impreſſa. Elytra breuia, laeuiſſima, ferruginea, apice nigra. Corpus atrum, femoribus compreſſis.

9. C. depreſſus, thorace elytrisque abbreuiatis rufis, capite biſpinoſo. *ſpinoſus.*

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Medius. Antennae longitudine corporis, filiformes. piloſae: articulis cylindricis. Caput atrum, nitidum, inter antennas ſpinis duabus porrectis, validis, acutis. Thorax planus, laeuis, rufus, baſi vtrinque impreſſo. Elytra valde abbreuiata, ſtriata, rufa. Corpus atrum. Pedes breues, nigri, femoribus incraſſatis, rufis.

*oblongus.* 10. C. thorace integro, biſulcato, niger, elytris ſtriatis fuſcis.

Habitat in America boreali. Muſ. D. de Seheſtedt.

Corpus reliquis longius, anguſtius. Caput nigrum, punctis duobus impreſſis, oblongis. Thorax integer, ſulcis duobus abbreuiatis, dorſalibus, impreſſis. Elytra ſtriata, fuſca. Pedes breues, femoribus compreſſis.

*planus.* 11. C. planus, teſtaceus, thorace punctis duobus baſeos impreſſis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Corpus paruum, planum, teſtaceum, immaculatum. Thorax punctis duobus baſeos impreſſis. Elytra vix ſtriata.

*dermeſtoides.* 12. C thorace ſulcato, fuſcus, elytris laeuibus teſtaceis. Ent. ſyſt. 2. 96. 10. *

I. 35 Panz. Fn. Germ. 3. tab. 13.

Habitat in Germania. D. Smidt.

*muticus.* 13. C. thorace mutico, nigro: puncto vtrinque impreſſo, elytris ſtriatis fuſcis. Ent. ſyſt. 2. 96. 12. *

VIII. 25. Payk. Fn. Sv. 2. 169. 6.

Habitat in Germania. D. de Hattorff.

14. C.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

14. C. thorace mutico, niger, thoracis margine elytrorumque macula ferrugineis. Ent. syst. 2. 96. 13. * *monilis.*

Payk. Fn. Sv. 2. 169. 7. *I. 35.*
Panz. Fn. Germ. 4. tab. 12. *36*
Habitat in Germania.

15. C. thorace mutico, ater, nitidus, elytris laeuissimis, pallidis. *minutus.*

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Statura omnino C. monilis, at duplo minor. Caput et thorax laeuia, atra, nitida. Elytra laeuissima, pallida, immaculata.

## 115. PYTHO. *Palpi* sex extrorsum crassiores. *Ligula* breuis, membranacea, quadrata, emarginata. *Antennae* filiformes.

1. P. thorace sulcato, niger, elytris striatis, coeruleis, abdomine rufo. *coeruleus.*

*Cucuius coeruleus* Ent. syst. 2. 94. 4.
*Pytho.* Latreille Inf. 50. 75.
*Tenebrio lignarius.* Degeer Inf. 5. 38. 4. tab. 2. fig. 13. *VIII. 25.*
*Tenebrio depressus.* Linn. syst. Nat. 2. 677. 24.
Payk. Fn. Sv. 2. 166. 2.
Habitat in Germania. D. Schneider.

2. P.

---

Pythonis corpus oblongum, depressum, planum, immarginatum, glabrum, tardum; *capite* ouato, thoraci angustiore, inserto; *oculis* paruis, rotundatis, prominulis, marginalibus; *antennis* distantibus, vix longitudine thoracis, pone mandibulas insertis; *thorace* rotundato, plano; *scutello* breui, rotundato; *elytris* rigidis, tegentibus, longitudine abdominis; *pedibus* breuibus, cursoriis; *tarsis* anticis quinque-, posticis quadriarticulatis; *colore* obscuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*feſtiuus.* 2. P. thorace ſulcato, niger, elytris ſtriatis coerule abdominis margine tibiisque rufis.

*Cucuius feſtiuus* Ent. ſyſt. 2. 95. 5. *

Habitat in Germania. D. Smidt.

A praecedente diuerſus videtur.

*caſtaneus.* 3. P. thorace ſulcato, niger, elytris ſtriatis fuſc abdominis margine pedibusque teſtaceis.

*Cucuius teſtaceus.* Ent. ſyſt. 2. 95. 6. *

Payk. Fn. Sv. 2. 166. 2. β.

VIII. 26.

Habitat in Germania. D. Smidt.

*Os* maxillis palpisque. *Palpi* ſex inaequales, e trorſum craſſiores, *anteriores* breues, biartic lati: articulis aequalibus: vltimo paullo cra ſiori, adhaerentes maxillae dorſo, *intermed.* multo longiores, exſerti, quadriarticulati: a ticulo primo breui, globoſo. ſecundo longior obconico, tertio quartoque aequalibus, paull craſſioribus, apice obtuſis, ad anteriorem in ſerti, *poſteriores* breuiores, extrorſum craſſio res, triarticulati: articulo primo breui, globo ſo, ſecundo longiori, obconico, tertio breui craſſiore, obtuſo, adnati ligulae baſi laterali *Mandibula* cornea, incurua, fornicata, apic parum fiſſa. *Maxilla* breuis, cornea, cylindri ca. obtuſa. *Ligula* breuis, membranacea, qua drata. apice late emarginata. *Labium* cor neum, rotundatum, integrum.

P. feſtiuus & caſtaneus mihi
P. coerulei varietates ſunt. Ill. III.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

16. BRONTES. *Palpi* quatuor aequales, filiformes.
*Maxilla* bifida.
*Antennae* elongatae, filiformes: articulo primo longiori, incuruo.

B. thorace denticulato, rufus, elytris nigris. *dubius.*
*Cucuius dubius.* Ent. syst. 2. 95. 7. *
Habitat in America boreali.

B. thorace denticulato, nigro, pedibus flauescentibus. *flauipes.*
*Cucuius flauipes.* Ent. syst. 2. 95. 8. * *Fn. 16.*
*Cerambyx planatus.* Linn. syst. Nat. 2. 625. 15.
Payk. Fn. Sv. 2. 167. 3.
*Vleiota.* Latreille Inf. 50. 76.
Habitat in Europae borealis truncis decorticatis.
Character generis naturalis adhuc desideratur. Video maxillas palposque. *Palpi* anteriores filiformes, breues, quadriarticulati: articulo primo breui, secundo longiori, obconico, vltimo acutiusculo, adhaerentes maxillae dorso. *Mandibula* breuis, cornea, incurua, edentula, acuta. *Maxilla* breuis, membranacea, bifida: laciniis aequalibus, obtusis. Reliqua haud obseruaui.

3. B.

---

Brontae corpus paruum, oblongum, depressum, planum, glabrum, immarginatum, agile; *capite* ouato, inserto; *oculis* globosis, prominulis, marginalibus; *antennis* distantibus, longitudine corporis, pone mandibulas insertis; *thorace* subquadrato, vix marginato: angulo antico prominulo, acuto, postico rotundato; *scutello* paruo, rotundato; *elytris* rigidis, tegentibus, longitudine abdominis; *pedibus* breuibus, validis, cursoriis; *tarsis* quadriarticulatis; *colore* obscuro.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*pallens.* 3. B. thorace ſerrato, obſcuro, elytris ſtriatis, abd
mine pedibusque teſtaceis.
*Cucuius pallens.* Ent. ſyſt. 2. 96. 9. *
praecedentis varietat. Ill. 3. *Cucuius flauipes.* Payk. Fn. Sv. 2. 167. 3. β.
Habitat in Germania. D. Smidt.

*teſtaceus.* 4. B. thorace ſubquadrato mutico, teſtaceus, fem
ribus compreſſis.
*Cucuius teſtaceus.* Ent. ſyſt. 2. 96. 11. *
Habitat ſub Betulae cortice. Muſ. D. Romani.

*minimus.* 5. B. thorace mutico, ferrugineus, elytris laeuiſ
mis, pallidis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Minutus et praecedentibus triplo adhuc mino
Corpus planum, ferrugineum, elytris pallidis.

## 117. COSSYPHVS. *Palpi* inaequales,
*anteriores* ſecuriforme
*Maxilla* bifida.
*Antennae* claua perfoliat

*depreſſus.* 1. C. antennis clypeo breuioribus. Ent. ſyſt. 2. 9
1. *
= *Coſſyphus depreſſus.* Oliv. Inſ. 44. tab. 1. fig.
Herbſt. Arch. tab. 46. fig. 7. et Lusitania
ex Ill. 3. Habitat in India orientali et Mauritania.

*planus.* 2. C. antennis clypeo longioribus.
= Habitat Tranquebariae. Muſ. D. de Seheſtedt.
Praecedenti nimis affinis, et forte mera eius v
rietas maior.

† Herbst. Coleopt. 7. 229. tab. 109. fig. 13
Coll. Hoffmannseggi

= Herbst. 7. 228. tab. 109. fig. 12 11
Coll. depressus.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

118. LAMPYRIS. *Palpi* quatuor, ſubclauati.
*Maxilla* bifida.
*Labium* corneum, integrum.
*Antennae* filiformes.

* *corpore anguſto oblongo.*

1. L. oblonga, fuſca, clypeo cinereo. Ent. ſyſt. 2. 98. 1. *noctiluca.*
*Lampyris noctiluca.* Linn. ſyſt. Nat. 2. 643. 1.
Fn. Sv. 699.
Oliv. Inſ. 28. tab. 1. fig. 2.
Panz. Fn. Germ. 41. tab. 9. *IV. 55.*
Degeer Inſ. 4. 31. 1. tab. 1. fig. 19. 20.
Payk. Fn. Sv. 2. 170. 1.
Habitat in Europae borealis Iuniperetis.

2. L. oblonga. fuſca, clypeo apice hyalino. Ent. ſyſt. 2. 98. 2. *ſplendidula*
*Lampyris ſplendidula.* Linn. ſyſt. Nat. 2. 644. 3.
Oliv. Inſ. 28. 11. 1. tab. 1. fig. 1.
Geoff. Inſ. 1. 166. 1.
Bradl. nat tab. 26. fig. 3. *IV. 55*
Panz. Fn. Germ. 41. tab. 8.
Payk. Fn. Sv 2. 171. 2.
Habitat in Europae ſyluaticis, tempeſtate pluuioſa lucem viuaciſſimam ſpargens.

3. L.

---

Lampyridis corpus oblongum, glabrum, marginatum, tardum; *capite* paruo, rotundato, ſub thoracis clypeo recondito; *oculis* magnis, globoſis, approximatis, lateralibus; *antennis* approximatis, thorace breuioribus, ſupra oculos inſertis; thoracis *clypeo* marginato, dilatato, caput tegente; *ſcutello* rotundato; *elytris* molliuſculis, incumbentibus, longitudine abdominis ſaepius phoſphoreſcentis; *pedibus* breuibus, curſoriis; *tarſis* quinquearticulatis; *colore* obſcuro.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*corruſca.* 3. L. obſcure nigra, clypeo vtrinque lunula roſea. Ent. ſyſt. 2. 98. 3. *
*Lampyris corruſca.* Linn. ſyſt. Nat. 2. 644. 2.
Oliv. Inſ. 28. tab. 2. fig. 14.
Habitat in America. D. Blackburn.

*analis.* 4. L. fuſca, clypeo elytrorumque margine flauis, abdomine apice albiſſimo, antennis compreſſis.
Habitat in China. Muſ. D. Lund.
Magna. Antennae valde compreſſae, obſcurae. Thoracis clypeus rotundatus, flauus, immaculatus. Pectus et abdominis baſis fuſca. Vltima abdominis ſegmenta albiſſima. Pedes teſtacei.

*obſcura.* 5. L. nigra, clypeo vtrinque macula rufa. Ent. ſyſt. 2. 99. 4. *
Oliv. Inſ. 28. tab. 2. fig. 13.
Habitat in Terra del Fuego. Muſ. D. Banks.

*bicolor.* 6. L. ſupra teſtacea, ſubtus atra, abdominis ſegmentis duobus penultimis, ſtriga abbreuiata albiſſima.
Habitat in noua Cambria. D. Billardiere.
Magna in hoc genere. Antennae compreſſae, ſerratae. Clypeus et elytra laeuia, teſtacea, immaculata. Corpus totum atrum, ſegmentis duobus penultimis ſtriga abbreuiata pellucida.

*laticornis.* 7. L. ouata, atra, clypei margine rufo, antennis compreſſis. Ent. ſyſt. 2. 99. 5. *
*Lampyris atra.* atra, thoracis margine rufo. Ent. ſyſt. 2. 101. 17. *
Oliv. Inſ. 28. tab. 3. fig. 28.
Habitat in America boreali.

8. L.

---

Lampyrides plantis victitant, in tenebris lucem phoſphoreum ſpargunt. Feminae apterae.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

8. L. clypeo rufo, medio nigro, elytris nigris: marginibus abdomineque albis. Ent. ſyſt. 2. 99. 6. * *pyralis.*
*Lampyris pyralis.* Linn. ſyſt. Nat. 2. 644. 4.
Muſ. Lud. Vlr. 78.
Oliv. Inſ. 28. tab. 2. fig. 11.
Degeer Inſ. 4. 52. 7. tab. 17. fig. 7.
Habitat in America meridionali.

9. L. fuſca, thorace, elytrorum margine abdominisque apice fuluis. Ent. ſyſt. 2. 99. 7. * *cincta.*
*Lampyris cincta.* Oliv. Inſ. 28. tab. 3. fig. 22.
Habitat in Poulicandor. Muſ. D. Banks.

10. L. nigra, thoracis elytrorumque margine omni, ſcutello abdominisque poſtico flauis. Ent. ſyſt. 2. 100. 8. * *marginata.*
*Lampyris marginata.* Linn. ſyſt. Nat. 2. 644. 5.
Oliv. Inſ. 28. tab. 2. fig. 16.
~~Naturf. 24. 31. 43. tab. 1. fig. 43.~~
Habitat in America.

11. L. atra, clypeo, lateribus, baſi exteriore, elytrorum abdominisque apicibus albis, antennis pectinatis. Ent. ſyſt. 2. 100. 9. * *pectinata.*
*Lampyris pectinata.* Linn. ſyſt. Nat. 2. 649. 17.
Degeer. Inſ. 4. 51. 11. tab. 17. fig. 13.
Habitat in America.

12. L. elytris fuſcis: margine exteriori luteo, abdomine flauo. Ent. ſyſt. 2. 101. 13. * *lucida.*
*Lampyris lucida.* Linn. ſyſt. Nat. 2. 645. 8.
Oliv. Inſ. 28. tab. 2. fig. 19.
Degeer Inſ. 4. 51. 5. tab. 17. fig. 5.
Sulz. Hiſt. Inſ. tab. 5. fig. 22.
Habitat in America meridionali.

13. L. nigra, clypeo elytrisque baſi teſtaceis, abdomine punctis duobus albiſſimis. *guttula.*

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Media, oblonga. Caput cum antennis nigrum, ore teſtaceo. Clypeus rotundatus, integer, teſtaceus: linea media impreſſa. Elytra vix ſtriata, fuſca, baſi teſtacea. Corpus atrum, abdominis ſegmento penultimo et antepenultimo macula media albiſſima. Femora rufeſcentia.

*bettica.* 14. L. teſtacea, thoracis diſco fuſco, abdomine pallido.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura L. lucidae. Clypeus rotundatus, fuſcus: margine pallide teſtaceo. Elytra holoſericea, pallide teſtacea, margine concolore. Corpus obſcurum, abdomine flauo.

*nitidula.* 15. L. fuſca, thorace flaneſcente: macula media nigra, abdominis apice maculis duabus flauis. Ent. ſyſt. 2. 101. 14. *

Oliv. Inſ. 28. tab. 2. fig. 15.

Habitat in Africa aequinoctiali. Muſ. D. Banks.

*phoſphorea.* 16. L. ſubteſtacea, abdomine atro, poſtice flauiſſimo. Ent. ſyſt. 2. 101. 15. *

*Lampyris phoſphorea.* Linn. ſyſt. Nat. 2. 645. 9.

Degeer Inſ. 4. 51. 6. tab. 17. fig. 6.

Habitat in America meridionali.

Antennae nigrae.

*Iaponica.* 17. L. flaua, ſegmento abdominis antepenultimo nigro. Ent. ſyſt. 2. 101. 16. *

Thunb. nou. ſp. Inſ. 4. 79.

Oliv. Inſ. 28. tab. 2. fig. 18.

Habitat in Iaponia; menſe Maii et Iunii copioſiſſima, veſperi lumine phoſphoreo e veſiculis binis caudalibus aëra replens. Thunberg.

18. L.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

18. L. thorace rufo, medio nigro, elytris nigris: margine omni pallido, antennis planis compreſſis. *compreſſicornis.*

Habitat in America meridionali. Muſ. D. de Seheſtedt.

Statura et magnitudo praecedentium. Antennae longitudine corporis, compreſſae, planae, ſerratae. Clypeus rotundatus, pallidus: medio late nigro. Elytra nigra, margine ſuturaque parum pallidis. Corpus nigrum, ano pallido.

19. L. pallida, abdominis ſegmento quarto albiſſimo. Ent. ſyſt. ſuppl. 124. 15. * *pallens.*

Brown. Iamaic. tab. 44. fig. 9.

Habitat in Americae meridionalis Inſulis. Muſ. D. Lund.

20. L. pallida, capite, ſcutello elytrorumque vitta marginali nigris, abdomine apice albiſſimo. *ſtrigata.*

Habitat in Inſulis maris pacifici. D. Billardiere.

Statura omnino L. Capenſis. Caput atrum. Clypeus transuerſus, caput vix tegens, vti in L. Italica, Capenſi, veſpertina, pallidus, linea dorſali nigra. Scutellum maculaque elytrorum ad ſcutellum nigra. Elytra ſtriata, pallida, vitta marginali. nigra. Corpus pallidum, ſegmentis duobus vltimis albiſſimis. Pedes pallidi, plantis fuſcis.

21. L. liuida, thorace flaueſcente, abdomine poſtice flauiſſimo. Ent. ſyſt. 2. 102. 18. * *Capenſis.*

Oliv. Inſ. 28. tab. 2. fig. 17.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

22. L. teſtacea, elytris apice nigris, abdomine apice albiſſimo. *veſpertina.*

Habitat in India orientali. D. Daldorff.

Media. Caput nigrum, oculis magnis, globoſis. Cly-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

peus rotundatus, teſtaceus, immaculatus. Elytra laeuia, teſtacea, apice nigra. Corpus teſtaceum, abdominis apice albiſſimo.

*auſtralis.* 23. L. flauicans, capite elytrisque fuſcis, abdomine apice albiſſimo. Ent. ſyſt. 2. 102. 19. *

Habitat in noua Hollandia. Muſ. D. Banks.

*thoracica.* 24. L. clypeo rufo, elytris atris, abdomine fuſco puncto medio albiſſimo.

*Lampyris thoracica* Oliv. Inſ. 28. tab. 3. f. 16.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Media. Antennae ſerratae, nigrae. Clypeus rotundatus, integer, ferrugineus, margine palli diore. Scutellum ferrugineum. Elytra atra immaculata. Pectus flaueſcens. Abdomen fuſcum, puncto paruo, medio, albiſſimo.

*planicornis.* 25. L. clypeo flaueſcente, elytris nigris, antennis apice albis.

Habitat in America meridionali, D. Smidt. Muſ. D. de Seheſtedt.

Statura omnino praecedentis paullo minor. Antennae compreſſae, nigrae: articulis vltimis duobus albis. Thorax rotundatus, rufeſcens. Elytra laeuia, nigra. Corpus flaueſcens, abdomine fuſco; ano albiſſimo.

*Italica.* 26. L. nigra, thorace transuerſo pedibusque rufis, abdomine apice albiſſimo. Ent. ſyſt. 2. 102. 20. *

*Lampyris Italica*, Linn. ſyſt. Nat. 2. 645. 11.

Oliv. Inſ. 28. tab. 2. fig. 12.

Degeer. Inſ. 4. 53. 9. tab. 17. fig. 9. 10.

Sulz. Hiſt. Inſ. tab. 6. fig. 3.

Habitat in Italiae arboribus.

27. L.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

27. L. nigra, thoracis clypeo rufo: margine pallescente, antennis flabellatis. *Caiennensis.*
Habitat in Cayenna. D. Richard.
Magnitudo L. Italicae, at omnino distincta. Antennae flabellatae, nigrae, articulo primo pallido. Clypeus rotundatus, rufus, margine hyalino. Elytra atra. Pedes antici testacei. Abdomen deest.

28. L. rufa, elytris nigris: margine omni vittaque media abbreuiata flauis. Ent. syst. 2. 102. 21. * *vittata.*
Oliv. Inf. 28. tab. 3. fig. 20.
Habitat in Insula Guadeloupe. D. de Badier.

29. L. nigra, thorace maculato, elytris margine vittaque media flauescentibus, abdomine apice albissimo. Ent. syst. suppl. 123. 21. * *versicolor.*
Habitat in America boreali. D. Herschel.

30. L. elytris liuidis, corpore flauo. Ent. syst. 2. 103. 22. *mauritanica.*
*Lampyris mauritanica.* Linn. syst. Nat. 2. 645. 10.
Oliv. Inf. 28. tab. 1. fig. 5.
Habitat in Africa.

31. L. nigra, elytris abbreuiatis, capite rufo: fascia oculari atra. *erythrocephala.*
Habitat in Carolina. Mus. D. Bosc.
Antennae magnae, nigrae, valde serratae et fere flabellatae. Caput rufum, fascia magna inter oculos. Thorax planus, submarginatus. Elytra corpore dimidio breuiora. Abdomen planum, atrum. Pedes atri.

32. L. fulua, antennis pectinatis fuscis, elytris abbreuiatis, attenuatis. *plumosa.*

*Lam-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Lampyris plumoſa.* Oliv. Inſ. 2. 28. 26. 27. tab. 3. fig. 27.
Habitat in America boreali.
Elytra breuiſſima, attenuata, fulua, immaculata.

*hemiptera.* 33. L. nigra, elytris breuiſſimis. Ent. ſyſt. 1. 163. 23. *
Geoff. Inſ. 1. 168. 2.
Oliv. Inſ. 28. tab. 3. fig. 25.
Habitat in Gallia. Muſ. D. Boſc.
in Silesia - Gradenberg

** *corpore latiori, ouato.*

*maculata.* 34. L. ouata, pallide teſtacea, elytris macula duabus marginalibus nigris. Ent. ſyſt. ſuppl. 125. 21. *
Oliv. Inſ. 28. tab. 1. fig. 3.
Habitat Surinami.

*lateralis.* 35. L. ouata, liuida, elytris macula marginali pallida.
Voet. Col. 1. tab. 48. fig. 3.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo praecedentis. Clypeus rotundatus, liuidus medio obſcuriore. Elytra laeuia, liuida: macula magna, pallida ad marginem exteriorem. Abdomen liuidum, anobiſſimo.

*flabellata.* 36. L. ouata, nigra, clypeo apice elytrisque faſciis duabus teſtaceis, antennis flabellatis.
Voet. Col. 1. tab. 48. fig. 5.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo L. maculatae. Antennae magnae, flabellatae, atrae. Clypeus rotundatus, baſi niger, apice teſtaceus. Elytra laeuia nigra

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nigra, fascia in ipsa basi mediaque testaceis. Corpus testaceum, pedibus atris.
*Variat* elytris testaceis, macula marginali apiceque nigris.

37. L. ouata, atra, clypeo apice testaceo, antennis flabellatis. *guttata.*
Habitat in America meridionali D. Smidt. Mus. D. de Seheistedt.
Nimis affinis L. flabellatae. Antennae valde flabellatae, atrae. Clypeus valde dilatatus, rotundatus, integer, niger, apice testaceus. Elytra laeuia, nigra, basi parum punctoque vno, alteroue testaceis. Corpus atrum.

38. L. ouata, nigra, clypeo ante marginem. elytris lineola marginali baseos abdomineque apice albis. Ent. syst. 2. 100. 10. * *flabellicornis.*
Oliv. Inf. 28. tab. 3. fig. 26.
Habitat in Brasilia. Mus. D. Banks.

39. L. ouata, elytris fuscis: macula marginali triangulari flaua. Ent. syst. 2. 100. 11. * *hesperа.*
*Lampyris hesperа.* Linn. syst. Nat. 2. 644. 6.
Oliv. Inf. 28. tab. 1. fig. 6.
Degeer Inf. 4. 49. 2. tab. 17. fig. 1.
Habitat in America.

40. L. ouata, elytris fuscis: macula marginali ouata lutea. Ent. syst. 2. 100. 12. *ignitа.*
*Lampyris ignita.* Linn. syst. Nat. 2. 645. 7.
Oliv. Inf. 28. tab. 1. fig. 7.
Degeer Inf. 4 49. 2. tab. 17. fig. 2.
Habitat in America meridionali.
Nimis affinis videtur L. laterali.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

119. OMALYSVS. *Palpi* aequales, extror-
trorſum craſſiores.
*Maxilla* membranacea,
bifida.
*Ligula* membranacea, e-
marginata.
*Antennae* filiformes.

*ſuturalis.* 1. OMALYSVS.
*Omalyſus.* Geoff. Inſ. 1. 180. 1. tab. 2. fig. 9.
*Omalyſus ſuturalis.* Oliv. Inſ. 24. 14. 1. tab. 1
fig. 1.
Ill. 41. Fourcroy Ent. 1. 64. 1.
Habitat in Gallia. Muſ. D. Olivier.

120. PYROCHROA. *Palpi* ſubfiliformes.
*Maxilla* integra.
*Ligula* lata, bifida: lac
niis aequalibus, rotun
datis.
*Antennae* filiformes.

*coccinea.* 1. P. nigra, thorace elytrisque ſanguineis immacula-
tis. Ent. ſyſt. 2. 104. 1.
*Lumpyris coccinea.* Linn. ſyſt. Nat. 2. 646. 18
Fn. Sv. 705.
*Pyrochroa.* Geoff. Inſ. 1. 338. 1. tab. 6. fig. 4
Oliv.

---

Pyrochroae corpus oblongum, planum, depreſſiuſculum, in
marginatum, tardum; *capite* ouato, exſerto; *oculis* rotun
datis, prominulis, marginalibus; *antennis* ſerratis, thorac
longioribus, ſub oculos inſertis; *thorace* rotundato, pla
no; *ſcutello* minuto, rotundato; *elytris* rigidiuſculis, for
nicatis, longitudine abdominis; *pedibus* longiuſculis, te
nuibus, curſoriis; *tarſis* anticis quinque-, poſticis qua
driarticulatis; *colore* rubro nigroque, nitidiuſculo.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 53. tab. 1. fig. 1.
Schaeff. Icon. tab. 90. fig. 4.
Degeer Inf. 4. 47. 4.
Panz. Fn. Germ. 13. tab. 11. II. 34.
Payk. Fn. Sv. 2. 172. 1.
Habitat in Europae fepibus.
*Mandibula* magna, craffa, apice attenuata, incurua; fubulata, acuta. *Ligula* elongata, membranacea, lata, fere ad baſin fiſſa: laciniis aequalibus, rotundatis. *Labium* corneum, integrum.

P. nigra, capite, thorace elytrisque ſanguineis immaculatis. Ent. ſyſt. 2. 105. 2. * *rubens.*
*Lampyris rubens.* Act. Hall. 1. 301.
Oliv. Inf. 53. tab. 1. fig. 2. VIII. 26.
Habitat in truncis putreſcentibus Salicis.
~~Vix a praecedente fatis diſtincta.~~

P. coccinea, antennis elytrisque nigris. Ent. ſyſt. 2. 105. 3. * *flabellata.*
Oliv. Inf. 53. tab. 1. fig. 3.
Habitat in America. Muſ. Dr. Hunter.

P. atra, thorace elytrisque teſtaceis, antennis pectinatis. Ent. ſyſt. 2. 105. 4. * *pectinicornis.*
*Cantharis pectinicornis.* Linn. ſyſt. Nat. 2. 650. 20. Fn. Sv. 713.
Oliv. Inf. 53. tab. 1. fig. 4. II. 34.
Panz. Fn. Germ. 13. tab. 12.
Habitat in Germania. D. Hybner.

P. rufa, elytris ſtriatis nigris: maculis quinque rubris, thorace vtrinque trituberculato. Ent. ſyſt. 2. 105. 5. * *tomaculata.*
Habitat in Cayenna. Muſ. D. Gigot d'Orcy.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

121. LYCVS. *Os* roſtro cylindrico, inermi.
*Palpi* quatuor: articulo vltimo craſſiori truncato.
*Antennae* filiformes.

*latiſſimus.* 1. L. flauus, elytris macula marginali poſticeque nigris: margine laterali maximo dilatato. Ent. ſyſt. 2. 106. 1. *
*Lampyris latiſſima* Linn. ſyſt. Nat. 2. 646. 14
Oliv. Inſ. 29. tab. 1. fig. 2.
Habitat in Sierra Leon Africae. D. Drury.

*palliatus.* 2. L. elytris coriaceis latiſſimis teſtaceis, apice nigris. Ent. ſyſt. 2. 106. 2. *
Oliv. Inſ. 29. tab. 1. fig. 3.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*roſtratus.* 3. L. niger, thoracis lateribus elytrisque dilatatis flauis, antice poſticeque nigris. Ent. ſyſt. 2. 106. 3. *
*Lampyris roſtrata.* Linn. ſyſt. Nat. 2. 646. 15
Muſ. Lud. Vlr. 78.
Oliv. Inſ. 29. tab. 1. fig. 4.
Degeer Inſ. 7. 622. 19. tab. 46. fig. 11.
Habitat in Africa.

*marginatus.* 4. L. ater, thorace marginato rufo: medio nigro maculis tribus pallidis, elytris rufis, apice nigris.
Habitat in Sumatra. D. Daldorff.

Magnus

---

Lyci corpus oblongum, glabrum, tardum; *capite* paruo, rotundato, ſub thoracis clypeo recondito, *ore* porrecto, cylindrico, roſtrato; *oculis* paruis, globoſis, marginalibus; *antennis* approximatis, ſerratis, thorace longioribus, ante oculos inſertis; thoracis *clypeo* plano, marginato, dilatato, caput tegente; *ſcutello* minuto, rotundato: *elytris* flexilibus, dilatatis, abdomine latioribus: *pedibus* brevibus, curſoriis; *tarſis* quinquearticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Magnus. Antennae compreſſae, nigrae. Thorax rufus, margine valde eleuatus et poſtice fere emarginatus. Diſcus niger, maculis tribus baſeos pallidis. Elytra ſulcata, rufa; apice nigra. Corpus atrum.

5. L. niger, thoracis lateribus elytrisque flauis, poſtice nigris. Ent. ſyſt. 2. 107. 5. * *proboſcideus.*
Oliv. Inſ. 29. tab. 1. fig 6.
Habitat in Africa aequinoctiali.

6. L. niger, thorace elytrisque rufeſcentibus, elytris apice nigris. Ent. ſyſt. 2. 107. 5. * *ſerratidornis.*
Oliv. Inſ. 29. tab. 1. fig 14.
Habitat in noua Hollandia. Muſ. D. Banks.

7. L. ater, thoracis margine elytrorumque baſi ruſis, antennis flabellatis. *dimidiatus.*
Habitat in Carolina. Muſ. D. Boſc.
Medius. Antennae atrae, dentibus longioribus, compreſſis flabellatae. Thorax in medio carinatus, ater, marginatus: margine rufo. Elytra ſulcata: ſulcis rugoſis, baſi rufa, ſcutello et ad ſcutellum nigra, apice atra. Corpus atrum.

8. L. ater, thoracis margine flaueſcente, elytris faſcia lata alba. Ent. ſyſt. 2. 107. 7. * *faſciatus.*
*Cantharis tropica*, thorace marginato: lateribus rubris, elytris nigris: faſcia alba, antennis pectinatis. Linn. ſyſt. Nat. 2. 649. 19.
Oliv. Inſ. 29. tab. 1. fig. 8.
Habitat in Cayenna. D. v. Rohr.

9. L. ater, thoracis margine flauo, elytris flauis, faſcia apiceque nigris. Ent. ſyſt. 2. 107. 8. * *reticulatus*
Oliv. Inſ. 29. tab. 1. fig. 7.
Habitat in America boreali.

10. L. ater, thoracis limbo elytrorumque margine humerali rufis. *humeralis.*

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Carolina. Muſ. D. Boſc.

Medius. Caput cum antennis atrum. Thorax inaequalis, marginatus, ater, margine omni rufo. Elytra ſulcata: ſulcis rugoſis, atra, margine baſeos, ſiue humerali rufo. Corpus atrum.

*ſerratus.* 11. L. flaueſcens, elytris ſtriatis: faſciis tribus nigris.

*Cantharis ſerrata.* Linn. ſyſt. Nat. 2. 649. 18.

Oliv. Inſ. 29. tab. 1. fig. 5.

Degeer Inſ. 4. tab. 1. fig 12.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Magnitudo et ſtatura praecedentium. Antennae compreſſae, ſerratae, nigrae. Caput flaueſcens. Thorax flaueſcens, linea dorſali nigra, margine reflexo. Elytra ſtriata, flaua: faſciis tribus nigris; baſeos abbreuiata, apicis latiore. Corpus flaueſcens.

*inaequalis.* 12. L. ater, thorace immarginato, inaequali, elytrisque rufis, his apice nigris.

Habitat in Sumatra. D. Daldorff.

Praecedente paullo minor. Caput atrum, antennis compreſſis, ſerratis. Thorax haud marginatus, at lineis variis eleuatis, inaequalis, rufus, immaculatus. Elytra ſulcata, rufa, apice nigra. Corpus atrum.

*ferrugineus.* 13. L. niger, thoracis lateribus elytrisque ſtriatis rufis. Ent. ſyſt. 2. 125. 4. *

Habitat in India orientali. Muſ. D. de Seheſtedt.

*praeuſtus.* 14. L. ferrugineus, elytris apice nigris. Ent. ſyſt. 2. 107. 6. *

Habitat Tranquebariae. Muſ. D. de Seheſtedt.

*tricolor.* 15. L. teſtaceus, elytris nigris: faſcia media teſtacea.

*Ly-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Lycus tricolor*. Oliv. Inſ. 29. tab. 1. fig. 10.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo L. praeuſti. Antennae compreſſae, atrae. Thorax teſtaceus, linea media margineque parum eleuatis. Elytra ſtriata, baſi fuſca, in medio teſtacea, apice nigra. Corpus teſtaceum.
*Variat* colore baſeos elytrorum magis et minus diluto.

6. L. ſanguineus, elytris poſtice violaceis. Ent. ſyſt. 2. 108. 9. * *bicolor.*
*Lampyris bicolor*. Linn. ſyſt. Nat. 2. 646. 16.
*Cantharis bicolor*. Linn. Amoen. acad. 6. 395. 21.
Oliv. Inſ. 29. tab. 1. fig. 9.
Habitat in Africa.

7. L. ater, immaculatus, antennis flabellatis. *atratus.*
Habitat in Oceani pacifici Inſulis. D. Billardiere.
Magnitudo praecedentium, at totus ater, immaculatus.

8. L. ater, elytris apice rufis. *haemorrhoidalis.*
Habitat in Oceani pacifici Inſulis. D. Billardiere.
Statura praecedentis, at antennarum articuli haud flabellati, ſed compreſſi obconici. Os vix porrectum palpis, vti mihi videtur, ſubſecuriformibus. Thorax vix marginatus, inaequalis, niger. Elytra ſulcata, atra, apice rufa.
Vix huius generis os haud porrectum, et palpi anteriores ſubſecuriformes.

9. L. ater, elytrorum margine omni rufo. *cinctus.*
Habitat in maris pacifici Inſulis. D. Billardiere.
Statura et affinitas praecedentium, at margo elytrorum et ſutura rufa. Antennae flabellatae.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ruſipennis.* 20. L. ater, elytris ſtriatis rufis.
Habitat in Oceani pacifici Inſulis. D. Billardiere
Omnino diſtinctus. Antennae flabellatae, tota atrae.

*linearis.* 21. L. thorace inaequali, ferrugineo: macula baſeo nigra, elytris linearibus ferrugineis, apice atris
Habitat in Guinea. Muſ. D. Lund.
Reliquis longior. Antennae atrae. Thorax ſub marginatus, inaequalis, rufus, macula baſeo nigra. Elytra valde linearia, elongata, ſulcata ſulcis punctatis, apice atra. Corpus atrum.

*cruentus.* 22. L. ater, thorace marginato elytrisque ſanguineis his apice atris.
Habitat in Sumatra. Muſ. D. de Seheſtedt.
Omnino diſtinctus a L. inaequali, et duplo mino; Antennae compreſſae, atrae. Thorax valde mar ginatus, ſanguineus, immaculatus. Elytra ſtria ta, ſanguinea, apice atra. Corpus atrum.

*galeatus.* 23. L. ater, thoracis limbo, elytrorum baſi, macul criſtata faſciaque media flauis.
Habitat in America meridionali. D. Smidt. Muſ D. Lund.
Statura et magnitudo omnino L. criſtati. Anten nae compreſſae, nigrae: articulo vltimo flauo Thorax magis planus et laeuis quam in reli quis, niger, margine omni flauo. Elytra laeui atra, ad baſin macula parua, et in hac criſti parua eleuata, compreſſa. In medio faſcia lata flaua. Corpus atrum.

*criſtatus.* 24. L. elytris ſtriatis ferrugineis, apice nigris: baſ criſta eleuata compreſſa.
Habitat in Guinea. Muſ. D. de Seheſtedt.
Paullo minor L. bicolore. Antennae totae atrae Thorax marginatus, rufus, linea media lata atra

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

atra. Elytra ſtriata, rufa, apice nigra, et baſi criſta magna eleuata, compreſſa, marginem fere tegente. Corpus atrum.

25. L. ater, thoracis limbo elytrisque reticulato-ſtriatis: macula baſeos faſciaque media flauis. *limbatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
L. ſerrato duplo fere minor. Antennae compreſſae, atrae. Thorax limbo eleuato, flauo. Elytra ſulcata, ſulcis lineolis transuerſis reticulata, macula baſeos ad marginem exteriorem faſciaque media flauis. Corpus atrum, pedum geniculis flauis.

26. L. niger, thoracis margine flauo, elytris linearibus baſi macula criſtata flaua. *compreſſus*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura praecedentium. Antennae compreſſae, nigrae. Thorax ater, limbo deflexo, flauo. Elytra linearia, laeuia, atra, macula parua baſeos flaua, et in hac criſta parua, compreſſa, eleuata. Corpus atrum, pedibus fuſcis: femoribus flauis.

27. L. niger, thoracis margine flauo, elytris ſtriatis faſciis: punctis impreſſis pellucidis. *diaphanus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Diſtinctus omnino a reliquis. Antennae compreſſae, atrae. Thorax niger, limbo omni lineaque dorſali eleuatis rufis. Elytra ſtriata, fuſca: ſulcis punctis pellucidis ſtriata. Corpus nigrum, femoribus baſi flauis.

28. L. ater, thoracis limbo ferrugineo, elytris ſubtiliſſime ſtriatis. *diſcoideus.*



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Mu
D. de Seheſtedt.

Paullo minor L. Aurora. Antennae compreſſ
atrae. Thorax niger, limbo eleuato, ferrugine
Elytra ſubtiliſſime ſtriata, atra, immacula
Pedes atri, femoribus baſi rufis.

*Variat* forte ſexu antennis flabellatis.

*ſanguineus* 29. L. niger, thoracis lateribus elytrisque glabri
ſanguineis. Ent. ſyſt. 2. 108. 10.

*Lampyris ſanguinea.* Linn. ſyſt. Nat. 2. 646. 1
Fn. Sv. 704.

Geoff. Inſ. 1. 168. 3.

IV. 55 Panz. Fn. Germ. 41. tab 9.

Oliv. Inſ. 29. tab. 1. fig. 1.

Friſh. Inſ. 12. tab. 3.

Schaeff. Icon. tab. 24. fig. 1.

~~Sulz. Hiſt. Inſ.~~ tab. 6. fig. 4.

~~Degeer Inſ. 4. 45.~~ 2. tab. 2. fig. 2.

Payk. Fn. Sv. 2. 174. 1.

Habitat in Europae lapidoſis.

*aurora.* 30. L. niger, thoracis lateribus elytrisque reticul
to-ſtriatis rufis.

Panz. Fn. Germ. 41. tab. 10.

IV. 56 Herbſt. Arch. tab. 5. 105. 4.

Voet. Col. 2. tab. 47. fig. 9.

*Varietas* thorace toto atro, vix differt.

*nigripes.* 31. L. ſanguineus, antennis pedibusque nigris. En
ſyſt. 2. 108. 12. *

*Lycus nigripes.* Oliv. Inſ. 29. tab. 1. fig. 12.

Habitat in Siam. Muſ. D. Banks.

*flabellicor-nis.* 32. L. niger, thoracis margine teſtaceo, antennis fe
ratis. Ent. ſyſt. 2. 109. 14. *

Habitat in Cayenna. D. de Paykull.

33. L

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

8. L. ater, thoracis margine antennarumque vltimo articulo rufis. *pectinatus.*

*Cantharis pectinata.* atra, clypei lateribus, basi exteriori elytrorum latius rufo, antennis pectinatis. Linn. syst. Nat. 2. 649. 17.

Degeer Inf. 4. tab. 17. fig. 13.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Minor. Antennae compressae, nigrae, articulo vltimo rufo. Elytra striata, sulcis lineolis transuersis, subreticulatis, nigra, margine baseos rufescente.

4. L. ater, antennarum apicibus elytrisque sanguineis. Ent. syst. 2. 108. 13. * *minutus.*

Oliv. Inf. 29. tab. 1. fig. 13.

Panz. Fn. Germ. 41. tab. 11.

Panz. Naturf. 24. 31. 45. tab. 1. fig. 44.

Payk. Fn. Sv. 2. 176. 4.

Degeer Inf. 5. 47. 3.

Habitat in Europa boreali. Mus. D. Lund.

5. L. ater, thoracis margine elytrorumque lineola baseos ferrugineis, elytris laeuibus. *laeuigatus.*

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Corpus paruum. Antennae compressae, totae nigrae. Caput et thorax nigra, thoracis margine ferrugineo. Elytra laeuia, atra, basi lineola parua, ferruginea. Corpus atrum.

6. L. ater, thorace rufo, antennis simplicibus. Ent. syst. 2. 109. 15. * *ruficollis.*

Habitat ad Cap. Bon. Spei. Mus. D. de Paykull.

7. L. cyaneus, thorace rufo, antennis flabellatis. *thoracicus*

Habitat in Australasia. D. Billardiere.

Affinis videtur praecedenti, at maior. Antennae com-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

compreſſae, flabellatae. Thorax rufus. Elytra cyanea, ſulcata: ſulcis rugoſis.

*marginellus.* 38. L. ater, thoracis margine rufo, elytris ſulcatis: ſulcis rugoſis.

Habitat in Carolina. Muſ. D. Boſc.

Paruus. Antennae ſimplices, atrae. Thorax marginatus, niger, margine late rufo. Elytra ſulcata, ſulcis rugoſis. Corpus atrum.

## 122. RIPIPHORVS.

*Palpi* filiformes.
*Maxilla* breuiſſima, ouata.
*Labium* acutum.
*Antennae* flabelliformes.

*ſubdipterus.* 1. R. elytris breuiſſimis, ouatis, fornicatis, palleſcentibus. Ent. ſyſt. 2. 109. 1. *

Habitat in Galloprouincia. Muſ. D. Boſc.

*ferrugineus.* 2. R. rufus, pectore, ano pedibusque nigris. Ent. ſyſt. 2. 110. 2. *

Habitat in India orientali. Dr. Koenig.

*naſutus.* 3. R. corpore atro, immaculato. Ent. ſyſt. 2. 110. 3. *

*Mordella naſuta.* Thunb. nou. ſp. Inſ. 3. 66. fig. 77.

Habitat in Iaponia.

4. R.

---

Ripiphori corpus paruum, oblongum, compreſſum, gibbum, immarginatum, agile; *capite* ouato, adſcendente, exſerto; *oculis* ouatis, marginalibus; *antennis* breuibus, diſtantibus, cantho oculorum iuſertis; *thorace* gibbo, antice attenuato, poſtice trilobo; *ſcutello* nullo; *elytris* rigidis, attenuatis, abdomine breui, obtuſo, ſublongioribus; *pedibus* tenuibus, compreſſis; *tarſis* anticis quinque-, poſticis quadriarticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. R. ater, elytris baſi flauis, thorace poſtice dente compreſſo, recumbente. Ent. ſyſt. ſuppl. 126. 3. * *ſpinoſus.*

Habitat in Americae meridionalis Inſulis. Dom. v. Rohr.

5. R. rufus, elytris nigris. Ent. ſyſt. 2. 111. 4. * *pectinatus.*

Habitat in America. D. Drury.

6. R. niger, thoracis lateribus elytrisque teſtaceis. *paradoxus*

Ent. ſyſt. 2. 111. 5,

*Mordella paradoxa.* Linn. ſyſt. Nat. 2. 682. 1. Fn. Sv. 831.

Roſſ. Fn. Etr. 1. 244. 603.

Pauz. Fn. Germ. 26. tab. 14.

Payk. Fn. Sv. 2. 177. 1.

Habitat in Europae floribus.

7. R. teſtaceus, ore, pectore abdominisque dorſo atris. Ent. ſyſt. 2. 111. 6. * *flabellatus.*

Cyrill. Ent. Neap. 1. tab. 4. fig. 1.

Habitat in Italia. Dr. Allioni.

8. R. niger, thorace rufo, elytris teſtaceis, apice obſcurioribus. *humeratus*

Habitat in Carolina. Muſ. D. Boſc.

Magnus. Caput nigrum, antennis flabellatis. Thorax ſubpunctatus, rufus. Elytra abdomine longiora, acuta, teſtacea, apice nigra. Corpus atrum.

9. R. ater, elytris maculis duabus ſanguineis. *triſtis.*

Habitat in Carolina. Muſ. D. Boſc.

Paruus in hac familia, totus ater nitidus, elytris ſolis maculis duabus ſanguineis.

10. R. rufus, nitidus, abdominis baſi elytrisque atris. *nigricornis.*

Habitat in Carolina. Muſ. D. Boſc.

Paruus. Caput rufum, nitidum, antennis ſerratis, atris.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

atris. Thorax laeuis, rufus, nitidus, poſtice loco ſcutelli porrectus. Elytra laeuia, atra, immaculata. Corpus rufum, abdominis baſi pedibusque nigris.

*punctatus.* 11. R. thorace teſtaceo, nigro-punctato, elytris nigris: margine teſtaceo. Ent. ſyſt. 2. 111. 7.
Habitat in Cayenna. D. v. Rohr.

*6maculatus.* 12. R. ater, thorace ferrugineo, elytris teſtaceis: macula nigra. Ent. ſyſt. 2. 111. 8. *
Habitat in America. Muſ. D. Banks.

*ventralis.* 13. R. ater, abdomine ſanguineo.
Habitat in Carolina. Muſ. D. Boſc.
Statura omnino, nigricornis, at totus ater R. nitidus, abdomine ſolo ſanguineo.

*lituratus.* 14. R. capite thoraceque ferrugineis, elytris teſtaceis: litura baſeos punctoque apicis atris. Ent. ſyſt. ſuppl. 127. 8. *
Habitat in Germania australi.

*2maculatus.* 15. R. ferrugineus, pectore atro, elytris teſtaceis: macula nigra. Ent. ſyſt. 2. 112. 9. *
Habitat in Hungaria. D. Saldoner.

*dimidiatus* 16. R. ater, elytris baſi flauis. Ent. ſyſt. 2. 112. 10.*
Habitat in America boreali. Muſ. D. Yeats.

*2punctatus* 17. R. niger, elytris teſtaceis; macula apiceque nigris.
Habitat Tranquebariae. Muſ. D. Lund.
Statura praecedentium. Caput atrum, antennis flabellatis, flauis. Thorax holoſericeo-nitens, niger. Elytra laeuia, teſtacea, macula media apiceque nigris. Corpus nigrum.

*tarſatus.* 18. R. ater, pedum tarſis niueis: articulis apice nigris.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura praecedentium. Corpus totum laeue, atrum, nitidum, ſolis pedum quatuor poſteriorum tarſis niueis: articulis apice nigris. Antennae flabellatae.

19. R. ferrugineus, vertice thoracis diſco elytrisque atris. Ent. ſyſt. 2. 112. 11. ♂ *limbatus.*
Habitat - - Muſ. D. Yeats.

20. R. niger, elytris teſtaceis: faſciis tribus nigris; prima annulari. Ent. ſyſt. 2. 112. 12. ♀ *populneus.*
Habitat in Germaniae Populo. D. Smidt.

## 123. MORDELLA.

*Palpi* inaequales:
*anteriores* clauati,
*poſteriores* filiformes.
*Maxilla* bifida.
*Ligula* bifida.
*Antennae* moniliformes.

1. M. ano aculeato, corpore atro immaculato. Ent. ſyſt. 2. 113. 1. *aculeata.*
*Mordella aculeata.* Linn. ſyſt. Nat. 2. 682. 2.
Fn. Sv. 532.
Geoff. Inſ. 1. 353. 1.
Schaeff. Elem. tab. 84.
Sulz. Inſ. tab. 7. fig. 46.
~~Schaeff. Icon. tab. 127. fig. 7.~~
Payk. Fn. Sv. 2. 185. 3.

Degeer.

---

Mordellae corpus paruum, oblongum, compreſſum, gibbum, immarginatum, agile; *capite* rotundato, inflexo, exſerto; *oculis* ouatis, marginalibus; *antennis* longitudine thoracis, diſtantibus, ſub oculis inſertis; *thorace* gibbo, transuerſo; *ſcutello* minuto; *elytris* rigidis, attenuatis, longitudine abdominis, feminae ſaepius aculeati; *pedibus* validis, curſoriis; *tarſis* anticis quinque-, poſticis quadriarticulatis; *colore* minus nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer. Inf. 5. 28. 1.

Habitat in Europae floribus semifloſculoſis, vmbellatis.

E Carolina duplo maior, ſimillima, vix diſtincta in Muſ. D. Boſc.

*mixta.* 2. M. ano aculeato, cinereo atroque varia, elytris faſcia rufa.

Habitat in noua Cambria. D. Billardiere.

Magna in hoc genere. Caput cinereo-pubeſcens, macula magna, frontali, atra. Thorax cinereo-pubeſcens, linea dorſali lata maculaque vtrinque atris. Coleoptra baſi cinerea maculis quinque atris 2. 1. 2, media communi, in medio rufa, apice atra, lunulis duabus cinereis. Corpus cinereo atroque varium.

*faſciata.* 3. M. ano aculeato, nigra, elytris faſciis duabus cinereis. Ent. ſyſt. 2. 113. 2. *

Geoff. Inf 1. 354. 2.

Habitat in Europae auſtralioris Vrtica.

*hamata.* 4. M. ano aculeato, nigra, thorace lineolis duabus hamatis lunulisque duabus, baſeos cinereis, elytris lineatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo M. variegatae. Caput ferrugineo-villoſum. Thorax niger lineis duabus dorſalibus, antice hamatis, vtrinque lunula baſeos cinereis. Elytra nigra lineis abbreuiatis, baſeos cinereis. Corpus atrum, cinereo-villoſum.

*variegata.* 5. M. ano aculeato, atra, thorace rufo: macula nigra, elytris baſi macula rufa. Ent. ſyſt. ſuppl. 127. 3. *

Habitat Pariſiis. Muſ. D. Boſc.

6. M.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

6. M. ano aculeato, atra, capite thoraceque cinereo-variegatis, elytris punctis quinque albis. *10guttata.*
Habitat in Insula Iaua.
Maior M. aculeata. Caput et thorax valde variegata. Elytra atra: punctis quinque 3. oblique positis. 1. 1. Corpus atrum, abdomine striis tribus punctorum alborum. Aculeus ani valde prominet.

7. M. ano aculeato, griseo-pubescens, thorace macula dorsali, elytris puncto baseos fasciisque duabus atris. Ent. syst. suppl. 127. 3. * *pubescens.*
Habitat in America. Dom. Hybner.

8. M. ano aculeato, elytris nigris: punctis quatuor flauis; primo lunato. Ent. syst. 2. 114. 3. * *8punctata.*
Habitat in America septentrionali. Mus. D. Banks.

9. M. atra, elytris testaceis: marginibus nigris. Ent. syst. 2. 114. 4. * *testacea.*
Habitat in Africae Dauco. D. Prof. Vahl.

10. M. ano aculeato, ferruginea, elytris nigris. Ent. syst. suppl. 127. 3. * *nigripennis.*
Habitat in America. Dom. Hybner.

11. M. testacea, thorace maculis duabus nigris. Ent. syst. 4. Append. 443. 4. * *2maculata*
Habitat in Germania. D. Schneider.

12. M. ano aculeato, atra, elytris atomis numerosis albidis. *atomaria.*
Habitat Viennae. D. Scheidler. *Guttata, Payk.*
Statura omnino praecedentium, tota atra, elytris solis punctis plurimis, albidis, micantibus.

13. M. ano aculeato, atra, scutello aculeisque basi holosericeo-argenteis. *scutellaris.*
Habitat in America meridionali. Mus. D. de Seheestedt.

Paullo

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paullo maior M. aculeata. Caput et thorax ferrugineo paullo nitidula. Scutellum argenteum. Elytra atra. Aculeus elongatus, ater, basi holosericeo-argenteus. Corpus atrum, macula vtrinque ad basin abdominis holosericeo-argentea, nitida.

*vittata.* 14. M. ano aculeato, atra, thorace elytrorumque vitta abbreuiata, griseo-pubescentibus.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura omnino praecedentium. Caput et thorax nigra, pube tenuissima grisea tecta. Elytra nigra vitta ad futuram griseo-pubescente, quae tamen apicem haud attingit. Corpus nigrum, griseo-nitidulum, femoribus anticis interdum rufescentibus.

*haemorrhoidalis.* 15. M. ano aculeato, fusca, elytris apice ferrugineis.

Habitat in America meridionali. D. Smidt. Mus. D. de Schestedt.

Statura praecedentium. Corpus totum nigrum, ferrugineo parum nitidulum. Elytra apice late ferruginea. Pedes nigri, femoribus apice ferrugineis.

*2fasciata.* 16. M. ano aculeato, atra, elytris fasciis duabus rufis.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Praecedentibus minor, tota atra, elytrorum fasciis duabus late rufis.

*ferruginea* 17. M. ano aculeato, corpore testaceo immaculato.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura parua praecedentis. Corpus totum testaceum, holosericeo paullo micans, immaculatum.

13. M.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

18. M. ano aculeato, teſtacea, antennis, elytrorum apicibus aculeoque nigris. *brunnea.*
Habitat in Auſtria. D. de Meyerle.
Statura omnino praecedentium. Caput et thorax teſtacea, immaculata. Antennae apice nigrae. Elytra laeuia, teſtacea, apice obſcura. Abdomen obſcurum, aculeo nigro. Pedes teſtacei.

19. M. ano aculeato, atra, abdomine fuluo. Ent. ſyſt. 2. 114. 5. * *ventralis.*
Habitat in Germania. D. Smidt.

20. M. ano aculeato, nigra, thorace abdomineque fuluis. Ent. ſyſt. 2. 114. 6. *abdominalis.*
*Mordella abdominalis.* Payk. Fn. Sv. 2. 186. 5.
*Mordella bicolor.* Sulz. Hiſt. Inſ. tab. 7. fig. 15.
Habitat in Germania. D. de Hattorff.

21. M. atra, elytris baſi flaueſcentibus. Ent. ſyſt. 2. 114. 7. *humeralis.*
*Mordella humeralis.* Linn. ſyſt. Nat. 2. 682. 3. Fn. Sv. 833.
~~*Anaſpis.* Geoff. Inſ. 1~~. 356. 2.
Payk. Fn. Sv. 2. 187. 6.
Habitat in Europae floribus.

22. M. ano aculeato, nigra, thorace cinereo-nitidulo, elytris vnco baſeos cinereo apice ſubteſtaceis. *marmorata.*
Habitat in America meridionali. Muſ. D. de Seheſtedt.
Corpus paruum. Caput et thorax nigra, cinereo-nitidula. Elytra laeuia, nigra, baſi vnco cinereo, apice obſcura late teſtacea.

23. M. atra, ore, thoracis lateribus pedibusque teſtaceis. Ent. ſyſt. 2. 114. 8. * *lateralis.*
Habitat in Holſatiae floribus.

24. M. atra, fronte pedibusque flaueſcentibus. Ent. ſyſt. 2. 114. 9. *frontalis.*
Linn.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. syst. Nat. 2. 682. 4. Fn. Sv. 834.
Panz. Fn. Germ. 13. tab. 13.
Payk. Fn. Sv. 2. 183. 1.
Habitat in Europae borealis floribus.

*atra.* 25. M. atra, immaculata, ano inermi. Ent. syst. 2. 115. 10.
Habitat in Dania. Mus. D. de Sehestedt.

*thoracica.* 26. M. atra, capite thoraceque flauis. Ent. syst. 2. 115. 11.
Linn. syst. Nat. 2. 682. 8. Fn. Sv. 835.
Payk. Fn. Sv. 2. 184. 2.
Habitat in Europae borealis floribus.

*ruficollis.* 27. M. atra, ore, thorace pedibusque flauis. Ent. syst. 2. 115. 12. *
*Anaspis.* Geoff. Inf. 1. 356. 3.
Habitat in Gallia. Mus. D. Bosc.

*flaua.* 28. M. flaua, elytrorum apicibus nigris. Ent. syst. 2. 115. 13.
Linn. syst. Nat. 2. 682. 6. Fn. Sv. 836.
Panz. Fn. Germ. 13. tab. 14.
Payk. Fn. Sv. 2. 185. 2. β.
Habitat in Europae floribus frequens.

## 124. DONACIA.

*Palpi* filiformes.
*Maxilla* vnidentata.
*Ligula* integra.
*Antennae* setaceae.

*crassipes.* 1. D. plana, obscura, elytris striatis viridi-nitidulis, abdomine argenteo-villoso, femoribus posticis dentatis. Ent. syst. 2. 115. 1. *

Lepta-

---

Donaciae corpus depressiusculum, planum, glabrum, immarginatum, tardum; *capite* paruo, rotundato, inserto; *oculis* globo-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Leptura aquatica.* Linn. ſyſt. Nat. 2. 687. 1.
Fn. Sv. 671 var β
Degeer. Inſ. 5. 140. 18. tab. 4. fig. 14. 15.
Geoff. Inſ. 1. 229. 12.
Friſh. Inſ. 12. tab. 6. fig. 2.
Payk. Fn. Sv. 2. 191. 4.
Habitat in plantis aquaticis, Nymphaeis.

2. D. nigra, thorace elytrisque pallidis, elytris vni- *equiſeti.*
ſpinoſis. Ent. ſyſt. ſuppl. 128. 2. *
*Donacia appendiculata.* Panz. Fn. Germ. 24. II. 36
tab. 17. D. mucronata Hoppe
Habitat in Franconiae Equiſeto limoſo. D. Hoppe.

3. D. thorace rufo, nigro, bimaculato, elytris fuſcis *zoſterae.*
ruſo-ſtriatis, apice bidentatis.
*Rhagium muticum.* Ent. ſyſt. 2. 306. 11. * varièt. praecedentis
Habitat in Sueciae plantis aquaticis. Muſ. D. Lund.

4. D. coeruleo-nigra, femoribus poſticis incraſſatis *feſtucae.*
dentatis. Ent. ſyſt. 2. 116. 2. *
Panz. Fn. Germ. 29. tab. 2. III. 42.
Habitat in Germaniae feſtuca aquatica. D. Smidt. D. Illig. III. 179

5. D. violacea, nitida, femoribus poſticis, incraſſatis, violacea
dentatis.
Habitat in Germaniae aquoſis.
Affinis certe D. Feſtucae, at violacea, nitida.

6. D. aenea, elytris cupreis: margine omni viridi, fe- *dentipes.*
moribus poſticis dentatis. Ent. ſyſt. 2. 116. 3. * ——
Panz. Fn. Germ. 29. tab. 5.
Payk. Fn. Sv. 2. 195. 8. D. fasciata Hoppe III. 42
Habitat in Germaniae aquis. D. Smidt. 7. D.

---

globoſis, prominulis, marginalibus; *antennis* approximatis, thorace longioribus, inter oculos inſertis; *thorace* anguſtato, cylindrico; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis, ſaepius ſericei; *pedibus* longiuſculis, rotundatis, ſaepius femoribus clanatis; *tarſis* quadriarticulatis; *colore* ſaepius nitidulo.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Nymphaeae* 7. D. thorace elytrisque cupreis, corpore cinereo-villoso, femoribus posticis dentatis. Ent. syst. 2. 116. 4. *

Payk. Fn. Sv. 2. 195. 9.

III. 43. Panz. Fn. Germ. 29. tab. 6.

Habitat in Nymphaeae foliis.

*sagittariae* 8. D. supra aenea, elytris submicantibus, subtus villoso-aurea, femoribus posticis dentatis. Ent.

III. 43 syst. 2. 117. 5. * D. aurea Hoppe

Panz. Fn. Germ. 29. tab. 7.

Habitat in Germaniae plantis aquaticis. D. Smidt.

*nigra.* 9. D. nigra, elytris substriatis, abdomine pedibusque rufis. Ent. syst. 2. 117. 6. *

~~Payk. Fn. Sv. 2. 196. 10.~~

III. 46. ~~Panz. Fn. Germ. 29. tab.~~ 2. 3.

Habitat ad Germaniae aquas. D. Smidt.

*Variat* magis aenea, *Donacia discolor.* Hoppe Monogr.

*Lemnae.* 10. D. obscure aenea, elytris vitta submarginali, cuprea.

*Donacia limbata.* Panz. Fn. Germ. 29. tab. 12.

III. 47. Habitat in Germaniae aquosis. D. Smidt.

Media. Caput canaliculatum, cinerascens. Thorax canaliculatus, obscure aeneus. Elytra punctato-striata, obscure aenea: vitta lata, submarginali, cuprea. Pedes nigri, femoribus posticis elongatis, subclauatis, muticis.

*clavipes.* 11. D. aenea, abdomine argenteo-villoso, femoribus posticis clauatis, muticis. Ent. syst. 2. 117. 7. *

~~Panz. Fn. Germ. 29. tab. 13.~~

Payk. Fn. Sv. 2. 190. 3.

Habitat in Germaniae plantis aquaticis. D. Smidt.

12. D.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

12. D. nigra, pedibus poſticis elongatis: tibiis apice faſciculatis. Ent. ſyſt. 2. 117. 8. * *faſciculata*
Habitat in Cayenna. Muſ. D. Boſc.

13. D. aenea, nitidula, elytris ad ſuturam cupreis. Ent. ſyſt. 2. 118. 9. * *ſimplex.*
~~Panz. Fn. Germ. 29. tab. 15.~~ III. 47.
Habitat in Europae plantis aquaticis.
Confer. *Donacia ſemicuprea*: Panz. Fn. Germ. 29. tab. 14. quae vix differt.

14. D. aenea, nitida, capite abdomineque cinereo-villoſis. *Menyanthidis*
*Donacia ſimplex.* Payk. Fn. Sv. 2. 189. 2.
Habitat in Germaniae plantis aquaticis. D. Smidt. III. 43
Magnitudo D. Hydrochaeridis. Antennae rufeſcentes. Caput cinereo-tomentoſum. Thorax viridi-aeneus, nitidus. Elytra punctato-ſtriata, aenea, nitida. Corpus cinereo-tomentoſum, pedibus rufeſcentibus, muticis.

15. D. cineraſcens, ſubtus argenteo-villoſa, pedibus muticis. Ent. ſyſt. 2. 118. 10. * *Hydrochaeridis.*
Panz. Fn. Germ. 29. tab. 17.
Payk. Fn. Sv. 2. 189. 1. III. 44.
Habitat in plantis aquaticis.
Thorax vtrinque tuberculo obtuſo. Elytra punctato-ſtriata. Virginea Herbst



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

125. TRICHIVS. *Palpi* filiformes.
*Maxilla* bifida.
*Antennae* clauato lamellatae.

*Eremita.* 1. T. aeneo-ater, thorace inaequali, ſcutello ſulcato longitudinali. Ent. ſyſt. 2. 118. 1.
*Scarabaeus Eremita.* Linn. ſyſt. Nat. 2. 556. 74
Oliv. Inſ. 1. 6. 53. 71. tab. 3. fig. 17.
Roeſ. Inſ. 2. ſcarab. 1. tab. 3 fig. 6.
Schaeff. Icon. tab. 26. fig. 1.
Degeer Inſ. 4. tab. 10. fig. 22.
Panz. Fn. Germ. 41. tab. 12.
Voet. Scarab. tab. 3. fig. 21.
Herbſt. Col. 3. tab. 27. fig. 9.
Payk. Fn. Sv. 2. 198. 1.
Habitat in truncis putreſcentibus Pyri, Salicis.

*nobilis.* 2. T. auratus, abdomine poſtice albo punctato, elytris rugoſis. Ent. ſyſt. 2. 119. 2.
*Scarabaeus nobilis.* Linn. ſyſt. Nat. 2. 558. 81.
Fn. Sv. 401.
Geoff. Inſ. 1. 73. 6.
Oliv. Inſ. 6. 59. 72. tab. 3. fig. 10.
Roeſ. Inſ. 2. ſcarab. 1. tab. 3. fig. 1-5.
Schaeff. Icon. tab. 66. fig. 5.
Panz. Fn. Germ. 41. tab. 13.
Voet. ſcarab. tab. 4. fig. 28.

Dege

---

Trichii corpus ouatum, depreſſiuſculum, hirtum, immargin
tum, tardum; *capite* ouato, inſerto, labio prominulo; oc
*lis* ouatis, marginalibus, vix prominulis; *antennis* thora
breuioribus, diſtantibus, ſupter oculos inſertis; *thorace* antr
anguſtato, hirto; *ſcutello* paruo, triangulo; *elytris* rigid
includentibus, abdomine breuioribus; *pedibus* breuib
validis; *tibiis* anticis dentatis; *tarſis* quinquearticulati
*colore* variegato.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 4. 297. 26.
Herbſt. Col. 3. tab. 27. fig. 6.
Payk. Fn. Sv. 2. 199. 3.
Habitat larua in ligno putreſcente, imago in floribus. praeſertim vmbellatis.
*Mandibula* minuta, edentula, *Ligula* late emarginata.

. T. niger, thorace vtrinque puncto baſeos elytrisque duobus flauis. Ent. ſyſt. 2. 119. 3. * *spunctatus.*
*Scarabaeus variabilis.* Linn. ſyſt. Nat. 2. 358. 79.
Oliv. Inſ. 1. 6. 60. 73. tab. 4. fig. 27.
Voet. Col. tab. 5. fig. 41. 42.
Panz. Fn. Germ. 41. tab. 14.
Herbſt. Col. 3. tab. 27. fig. 7.
Payk. Fn. Sv. 2. 198. 2.
Habitat in Germaniae truncis putreſcentibus.

. T. niger, tomentoſo-flauus, elytris faſciis tribus abbreuiatis nigris. Ent. ſyſt. 2. 119. 4. *faſciatus.*
*Scarabaeus faſciatus.* Linn. ſyſt. Nat. 2. 556. 70. Fn. Sv. 395.
Geoff. Inſ. 1. 80. 16.
Oliv. Inſ. 1. 6. 1. 74. tab. 9. fig. 84.
Drury Inſ. 1. tab. 36. fig. 2.
Schaeff. Icon. tab. 1. fig. 4.
Degeer Inſ. 4. tab. 10. fig. 19.
Act. Nidroſ. 4. tab. 16. fig. 2.
Voet. Scarab. tab. 5. fig. 43.
Herbſt Col. 3. tab. 27. fig. 10.
Payk. Fn. Sv. 2. 200. 4.
Habitat imago in Europae borealis floribus vmbellatis.
*Larua* alba, brunneo-piloſa, capite brunneo, in truncis putreſcentibus Betulae.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fasciatus.* 5. T. niger, tomentoso-cinereus, elytris nigris: fasciis duabus flauis. Ent. syst. 2. 120. 5. *
Pall. Icon. tab. A. fig. 19.
Habitat in Germania. D. Hybner.

*Indus.* 6. T. thorace hirsuto, elytris liuidis, fusco-maculatis. Ent. syst. 2. 120. 6. *
*Scarabaeus Indus.* Linn. syst. Nat. 2. 556. 61.
Mus. Lud. Vlr. 27.
Oliv. Inf. 6. tab. 6. fig. 40.
Habitat in Indiis.

*2punctatus.* 7. T. capite thoraceque glabris, viridi-aeneis, elytris testaceis, abdominis vltimo segmento prominente, bipunctato. Ent. syst. 2. 120. 7. *
*Melolontha 2punctata.* Oliv. Inf. 1. 5. tab. 6. fig. 69.
Habitat in America. Mus. D. Banks.

*bibens.* 8. T. capite thoraceque viridi-aeneis, elytris testaceis, viridi-nitentibus. Ent. syst. 2. 121. 8. *
Oliv. Inf. 6. tab. 10. fig. 87.
Habitat in America. Mus. D. Tunstall.

*hemipterus.* 9. T. thorace tomentoso: rugis duabus longitudinalibus, elytris abbreuiatis. Ent. syst. 2. 121. 9. *
*Scarabaeus hemipterus.* Linn. syst. Nat. 2. 555. 63.
Geoff. Inf. 1. 78. 12.
Oliv. Inf. 1. 6. tab. 9. fig. 83.
tab. 11. fig. 103.
Schaeff. Icon. tab. 46. fig. 10. 11.
Voet. Col. tab. 10. fig. 88-90.
Habitat in Europae ligno putrescente.
*Femina* aculeo ani elongato, nigro.

*canaliculatus.* 10. T. fusco-testaceus, cinereo-maculatus, thorace canaliculato. Ent. syst. 2. 121. 10. *

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 1. 6. tab. 10. fig. 89.
Habitat in America boreali. D. Lee.

1. T. glaber, cyaneus, elytris lunulis duabus albis. Ent. ſyſt. 2. 121. 11. * *lunulatus.*
Oliv. Inſ. 1. 6. tab. 10. fig. 88.
Habitat in Carolina.

2. T. viridis, pubeſcens, ano albo, bimaculato. Ent. ſyſt. 2. 122. 12. * *viridulus.*
Oliv. Inſ. 6. tab. 9. fig. 86.
Habitat in India. Muſ. Dr. Hunter.

3. T. capite thoraceque aeneis, villoſis, elytris teſtaceis, albo-maculatis. Ent. ſyſt. 2. 122. 13. * *piger.*
Oliv. Inſ. 6. tab. 7. fig. 54.
Habitat in Marylandia.

4. T. thorace nigro: triangulo albo, elytris teſtaceis: puncto fuſco. Ent. ſyſt. 2. 122. 14. * *Delta.*
Oliv. Inſ. 6. tab. 11. fig. 107.
Forſt. Cent. Inſ. 1. 7. 7.
Drury Inſ. 2. tab. 30. fig. 1. 2.
Habitat in America boreali.

5. T. pubeſcens, thorace fuluo nigro-lineato, elytris teſtaceis: ſutura fulua. Ent. ſyſt. 2. 122. 15. * *lineatus.*
*Scarabaeus quadratus.* Degeer Inſ. 7. 645. 48. tab. 48. fig. 10. 11.
Oliv. Inſ. 6. tab. 7. fig. 63.
~~Voet. Col.~~ tab. 10. fig. 8.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

6. T. ater, nitidus, elytris ſtrigis duabus approximatis flauis, tibiis poſticis compreſſis. *retuſus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
T. Delta minor. Caput laeue, atrum, nitidum. Tho-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Thorax ater, nitidus, antice vtrinque retuſus. Elytra ſtriata, atra, ante medium ſtrigis duabus macularibus, approximatis, flauis. Pedes incraſſati, atri: vnguibus magnis, arcuatis. Tibiae poſticae magnae, compreſſae, ſubdentatae.

*nigripes.* 17. T. hirtus, fuſcus, elytris teſtaceis: margine apicis cineraſcente. Ent. ſyſt. 2. 123. 16. *

Oliv. Inſ. 6. tab. 9. fig. 85.

~~Voet. Col.~~ tab. 10. fig. 83.

Degeer Inſ. 1. tab. 48. fig. 9.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*maculatus.* 18. T. ſupra glaber, cinereo-maculatus, capite thoraceque nigris, elytris piceis. Ent. ſyſt. 2. 123. 17. *

*Cetonia craſſipes.* Oliv. Inſ. 6. 68. 83. tab. 7. fig. 62.

~~Degeer Inſ. 7.~~ tab. 48. fig. 12.

Habitat ad Cap. Bon. Spei.

*hirtus.* 19. T. hirtus, ſupra niger, ſubtus albidus. Ent. ſyſt. ſuppl. 128. 17. *

Habitat ad Cap. Bon. Spei. D. Daldorff.

*piloſus.* 20. T. hirtus, ſubteſtaceus, capite thoraceque nigris elytris piceis. Ent. ſyſt. ſuppl. 128. 17. *

Habitat ad Cap. Bon. Spei. D. Daldorff.

*minutus.* 21. T. thorace cinereo, elytris teſtaceis, cinereo-maculatis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Schéſtedt.

Paruus. Corpus cinereo-villoſum. Elytra teſtacea, cinereo-maculata.

126. CE-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

126. CETONIA. *Maxilla* apice ſetoſa.
*Ligula* emarginata, palpos tegens.
*Antennae* lamellatae.

1. C. thorace piceo, albo, lineato, elytris glaucis, clypeo porrecto, bifido. Ent. ſyſt. 2. 124. 1. * *goliata.*
*Scarabaeus goliatus.* Linn. Mant. 5. 30.
*Cetonia goliata.* Oliv. Inſ. 6. 7. 1. tab. 5. f. 33. tab. 9. f. 33.
Drury Inſ. 1. tab. 31. fig. 3. tab. 40. fig. 1.
Iabl. Col. 1. tab. 10. fig. 4.
Sulz. Hiſt. Inſ. tab. 1. fig. 2.
Habitat in Africa. Muſ. Dr. Hunter.

2. C. thorace flaueſcente, nigro-lineato, elytris albidis, nigro-maculatis, clypeo porrecto, bifido. Ent. ſyſt. 2. 124. 2. * *Cacicus.*
Oliv. Inſ. 6. 8. 2. tab. 3. fig. 22.
Iabl. Col. 1. tab. 11. fig. 2.
Voet. Col. tab. 21. fig. 51.
Habitat in America meridionali.

3. C. obſcure aenea, elytris fuſcis, albo irroratis, clypeo porrecto bifido, vertice haſtato. Ent. ſyſt. 2. 124. 3. * *bifrons.*
*Cetonia bifrons.* Oliv. Inſ. 6. tab. 6. fig. 117.

Habi-

---

Cetoniae corpus oblongum, laeue, immarginatum, tardum; *capite* ouato, inſerto; *clypeo* prominente: margine incraſſato; *oculis* globoſis, marginalibus; *antennis* diſtantibus, thorace breuioribus, ſub oculis inſertis; *thorace* antice anguſtato, elenato: *ſcutello* longiori, triangulo; *elytris* rigidis, ſornicatis, vix abdomine breuioribus; *pedibus* breuibus, validis, tibiis dentatis; *tarſis* quinquearticulatis; *colore* ſaepius nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. Muſ. D. Gigot d'Orcy.

*Polyphemus.* 4. C. viridis, albo-lineata, capite tricorni: cornu medio porrecto bifido. Ent. ſyſt. 2. 125. 4. *
Oliv. Inſ. 1. 6. 9. 3. tab. 8. fig. 61.
Habitat in Africa. Muſ. D. Banks.

*Vnca.* 5. C. thorace atro, albo-lineato, elytris brunneis, albo-irroratis, clypeo bipartito.
*Cetonia Vnca.* Web. Inſ. 66. 1.
Habitat in Peru.
C. goliata duplo minor.

*micans.* 6. C. viridis, nitens, clypeo porrecto, recurvo, bifido, tibiis anticis ſerratis. Ent. ſyſt. 2. 125. 5. *
Oliv. Inſ. 1. 6. 10. 4. tab. 1. fig. 2.
Drury Inſ. 1. tab. 32. fig. 3.
Habitat in Africa.
Hae priores ſex ſpecies ſingulares, clypeo valde elongato, recurvo, bifido, forte proprii generis, mihi haud rite examinatae.

*Chinenſis.* 7. C. aenea, clypeo emarginato, ſubſpinoſo, thorace poſtice lobato, elytris acuminatis. Ent. ſyſt. 2. 126. 6. *
Oliv. Inſ. 1. 6. 11. 5. tab. 2. fig. 5.
Forſt. Cent. Inſ. 1. 2. 2.
Voet. Col. tab. 5. fig. 40.
Habitat in China. Muſ. D. Banks.

*nigrita.* 8. C. glabra, atra, clypeo emarginato, ſubſpinoſo, thorace poſtice lobato. Ent. ſyſt. 2. 126. 7. *
Oliv. Inſ. 1. 6. 11. tab. 10. fig. 92.
Petiv. Gazoph. tab. 28. fig. 3.
Hermann. Muſ. 17. 257.
Habitat Ceyloniae. Muſ. Britann.

9. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

9. C. aurata, ſegmento abdominis primo lateribus vnidentato, elytris lineolis transuerſis albis. Ent. ſyſt. 2. 127. 8. *aurata.*
*Scarabaeus auratus.* Linn. ſyſt. Nat. 2. 557. 78. Fn. Sv. 400.
Geoff. Inſ. 1. 73.
Friſh. Inſ. 12. 25. tab. 3. fig. 1.
Roeſ. Inſ. 2. ſcarab. 1. tab. 2.
Oliv. Inſ. 1. 6. 12. 7. tab. 1. fig. 1.
Voet. ſcarab. tab. 1. fig. 1.
Schaeff. Icon. tab. 26. fig. 2. 3. 7.
Drury Inſ. 1. tab. 33. fig. 1.
Panz. Fn. Germ. 41. tab. 15.
Degeer Inſ. 4. tab. 11. fig. 1.
Herbſt. Col. 3. tab. 29. fig. 2.
Payk. Fn. Sv. 2. 202. 2.
Habitat larua in Europae ligno putreſcente, in formicarum aceruis. Imago in floribus.

10. C. aenea, nitidiſſima, immaculata. Ent. ſyſt. 2. 127. 9. * *faſtuoſa.*
*Cetonia ſpecioſiſſima.* Roſſ. Fn. Etr. 1. 25. 57.
Scopol. Del. Inſ. 1. tab. 21. fig. A.
Roeſ. Inſ. 2. ſcarab. 1. tab. 2. fig. 6.
Panz. Fn. Germ. 41. tab. 16.
Voet. Col. tab. 1. fig. 2.
Schaeff. Icon. tab. 66. fig. 4.
Habitat in Europa auſtraliori.

11. C. aenea, thorace elytrisque atomis numeroſis, albis, ſparſis. Ent. ſyſt. 2. 127. 10. * *marmorata.*
Voet. Col. tab. 1. fig. 3.
Panz. Fn. Germ. 41. tab. 17.
Herbſt. Col. 3. tab. 29. fig. 3.
Payk. Fn. Sv. 2. 201. 1.
Habitat in Germaniae quercu.

12. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*viridis.* 12. C. viridis, opaca, ſubtus nitidior, elytris albo-maculatis. Ent. ſyſt. 2. 128. 11. *

*Cetonia Hungarica.* Herbſt. Arch. tab. 29. f. 7.

Panz. Fn. Germ. 41. tab. 18.

Habitat in Italia. Dr. Allioni.

*recurua.* 13. C. aenea, nitida, ſterno cornuto, recuruo.

Habitat in Guinea. D. Iſert. Muſ. D. Lund.

Statura et magnitudo C. metallicae, tota aenea, nitida, immaculata, ſterno magno, porrecto, cornuto, recuruo.

*metallica.* 14. C. aenea, opaca, immaculata, ſubtus cuprea. Ent. ſyſt. 2. 128. 12. *

Panz. Fn. Germ. 41. tab. 19.

Habitat in Italia. Dr. Allioni.

*pubeſcens.* 15. C. obſcure aenea, thoracis ſegmentorumque margine albo, bipunctato. Ent. ſyſt. 2. 128. 13. *

*Cetonia pubeſcens.* Oliv. Inſ. 1. 6. 21. 18. tab. 11. fig. 100.

Habitat ad Cap. Bon. Spei. D. Olivier.

*opaca.* 16. C. viridis, opaca, capitis clypeo reflexo. Ent. ſyſt. 2. 128. 14. *

Habitat in Africae floribus compoſitis. D. Vahl.

*Morio.* 17. C. nigra, obſcura, corpore ſubtus nitidiore. Ent. ſyſt. 2. 129. 15. *

*Cetonia Morio.* Oliv. Inſ. 1. 6. 27. 27. tab. 2. fig. 3.

*Scarabaeus lugubris.* Voet. Col. tab. 2. fig. 14.

Habitat in Europa auſtrali. Dr. Allioni.

*cuſpidata.* 18. C. nigra, capitis clypeo porrecto, baſi vtrinque ſinuato, thoracis limbo elytrorumque margine cinereis. Ent. ſyſt. 2. 129. 16. *

Herbſt.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Col. tab. 27. fig. 8.
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

19. C. atra, nitida, capitis clypeo cordato. Ent. ſyſt. 2. 129. 17.* *cordata.*
Habitat ad Cap. Bon. Spei. Muſ D. Lund.

20. C. nigra, thorace punctis quatuor albis. Ent. ſyſt. 2. 129. 18. * *4punctata.*
Habitat in Italia. Dr. Allioni.

21. C. nigra, ano bipunctato, pedibus rufis: tibiis poſticis ſpinis annulatis. Ent. ſyſt. 2. 129. 19.* *rufipes.*
*Cetonia 4punctata.* Oliv. Inſ. 1. 6. 80. 101. tab. 10. fig. 93.
Habitat in India. Muſ. Dr. Hunter.

22. C. thorace poſtice lobato, atra, holoſericea, elytris maculis tribus marginalibus, flauis. Ent. ſyſt. 4. App. 449. 19.* *holoſericea.*
*Cetonia holoſericea.* Oliv. Inſ. 6. tab. 12. fig. 125?
Voet. Col. tab. 2. fig. 10.
Herbſt. Col. tab. 28. fig. 7.
Habitat Surinami.

23. C. thorace poſtice lobato, nigra, thorace lineolis, elytris ſtrigis abbreuiatis, inordinatis, ferrugineis. Ent. ſyſt. 4. App. 449. 19. * *ſtrigoſa.*
*Cetonia ſtrigoſa.* Oliv. Inſ. 2. 6. tab. 12. f. 124.
Voet. Col. tab. 3. fig. 19.
Herbſt. Col. tab. 31. fig. 9.
Habitat ad Cap. Bon. Spei.

24. C. thorace poſtice lobato, capite ſpina incumbente, ſterno cornuto. Ent. ſyſt. 2. 130. 20. *nitida.*
*Scarabaeus nitidus.* Linn. ſyſt. Nat. 2. 552. 52.
Muſ. Lud. Vlr. 26.

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 1. 6. 18. 14. tab. 3. fig. 16.
tab. 7. fig. 56.
~~Roef. Inf. 2. fcarab.~~ 1. tab. B. fig. 4.
Drury Inf. 1. tab. 33. fig. 5. 6.
Voet. Col. tab. 3. fig. 2. 3.
Degeer. Inf. 4. tab. 19. fig. 8. 9.
Habitat in America; praeſertim Carolina.

*lobata.* 25. C. nigra, nitida, thorace poſtice lobato, clypeo emarginato, ſterno cornuto. Ent. ſyſt. 2. 130. 21. *
*Cetonia lobata.* Oliv. Inf. 1. 6. 23. 21. tab. 4. fig. 26.
Voet. Col. tab. 3. fig. 20.
Habitat in America meridionali. Muſ. D. Olivier.

*Carmelita.* 26. C. nigro-viridis, thorace elytrisque teſtaceis, ano albo bipunctato. Ent. ſyſt. 2. 131. 22. *
Habitat in Africa. D. Saldoner.

*ſulcata.* 27. C. viridi-aenea, nitida, elytris ſulcatis. Ent. ſyſt. 2. 131. 23. *
*Cetonia ſulcata.* Oliv. Inf. 1. 6. 36. 39. tab. 5. fig. 32.
Habitat in Braſilia. Muſ. D. Banks.

*Chryſis.* 28. C. ſcutello elytris dimidio breuiore, ſterno porrecto, corpore viridi, ſubtus cupreo. Ent. ſyſt. 2. 131. 24. *
*Scarabaeus Chryſis.* Linn. ſyſt. Nat. 2. 591. 49. Muſ. Lud. Vlr. 21.
Gronov. Zooph. 423. tab. 15. fig. 1.
Oliv. Inf. 1. 6. 76. 96. tab. 4. fig. 19.
Sulz. Hiſt. Inf. tab. 1. fig. 10.
~~Voet. Col. tab. 8. fig.~~ 63.
Degeer Inf. 4. tab. 19. fig. 4.
Habitat in America meridionali.

29. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

29. C. ſcutello elytris dimidio breuiore, viridis, thorace vtrinque macula poſtica, liuida. *virens.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura omnino C. Chryſidis, at duplo minor, viridis, parum nitida. Macula thoracis vtrinque magna, liuida, micans, poſtica ad marginem exteriorem.

30. C. cyanea, thoracis margine elytrisque teſtaceis. Ent. ſyſt. 2. 141, 57. * *ſplendida.*
*Cetonia ſplendida.* Oliv. Inſ. 1. 6. 75. 95. tab. 4. fig. 21.
Voet. Col. tab. 8. fig. 62.
Habitat in America meridionali.
*Variat* maior, viridis, nitens, thoracis margine elytrisque teſtaceis.

31. C. ſcutello elytris dimidio breuiore, ſterno porrecto, corpore aeneo, nitido. Ent. ſyſt. 2. 131. 25. * *lucida.*
*Cetonia liuida.* Oliv. Inſ. 1. 6. 75. 94. tab. 7. fig. 64.
Habitat in Inſula Guadeloupe. D. Olivier.

32. C. obſcure viridis, thoracis margine flaueſcente. Ent. ſyſt. ſuppl. 129. 25. * *franciſca.*
Habitat in India. D. Daldorff.

33. C. picea, thoracis margine flaueſcente, ſterno porrecto. Ent. ſyſt. 2. 132. 26. * *capucina.*
Habitat in India. D. Daldorff.

34. C. exſcutellata, liuida, thorace punctis duobus, elytris maculis plurimis nigris, ſterno antice cornuto. Ent. ſyſt. 2. 132. 27. *Lanius.*
*Cetonia Lanius.* Oliv. Inſ. 1. 6. 19. 15. tab. 2. fig. 4.
Voet. Col. tab. 4. fig. 44.

Drury

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Drury Inſ. 1. tab. 33. fig. 8.
Sloan. Iamaic. tab. 237. fig. 7. 8.
Habitat in America meridionali.

*bajula.* 35. C. exſcutellata, liuida, thorace elytrisque fuſc variis, ſterno antice cornuto.
*Cetonia bajula.* Oliv. Inſ. 1. 6. tab. 12. fig. 126
Habitat in America meridionali. Muſ. D. de Sehe ſtedt.
Statura et ſumma affinitas C. Lanii, at thorax e elytra fuſco-variegata.

*flaveola.* 36. C. exſcutellata, liuida, capite thoracisque antic flauis, elytris nigro-variegatis, apice acuminatis
Habitat in America meridionali. D. Smidt. Muſ D. Lund.
Statura et magnitudo omnino praecedentium Caput flauum, apice liuidum. Thorax antice flauus, productus, poſtice liuidus. Elytra ni gro liuidoque variegata, apice ad ſuturam acu minata. Pectus liuidum, nigro-punctatum ſterno gibbo.

*graculus.* 37. C. exſcutellata, obſcure liuida, thorace elytrisque nigro-maculatis. Ent. ſyſt. 2. 132. 28.
*Cetonia maculoſa.* Herbſt. Col. 3. tab. 32. f. 1.
Oliv. Inſ. 6. tab. 12. fig. 120.
Voet. Col. tab. 3. fig. 22.
Habitat in America. Muſ. D. Banks.

*liturata.* 38. C. exſcutellata, obſcura, capite thoracisque margine liuidis, nigro-irroratis, elytris acuminatis: lituris flaueſcentibus.
*Cetonia liturata.* Oliv. Inſ. 1. 6. tab. 12. f. 121.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Paullo minor C. graculo. Capitis clypeus liuidus, nigro-irroratus, margine incraſſato. Thorax poſtice

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

poſtice productus, obſcurus, margine liuido, nigro-irrorato. Elytra acuminata, obſcura, lituris variis flaueſcentibus. Sternum vix porrectum.

39. C. ſcutellata rubra, elytris nigro-maculatis. Ent. ſyſt. 2. 132. 29. *carnifex.*
*Scarabaeus Lanius.* Linn. ſyſt. Nat. 2 557. 77.
Oliv. Inſ. 1. 6. 29. 16. tab. 6. fig. 43.
Roeſ. Inſ. 2. ſcarab. 1. tab. B. fig. 3.
Voet Col. tab. 2. fig. 16.
Herbſt. Col. 3. tab. 31. fig. 7.
Habitat in America meridionali.

40. C teſtacea, nitida, ſterno antice porrecto, obtuſo. Ent. ſyſt. 2. 133. 30. * *glabrata.*
*Cetonia glabrata.* Oliv. Inſ. 1. 6. 70. 86. tab. 9. fig. 80.
Habitat in Indiis. Muſ. Britann.

41. C. nigra, obſcura, elytris obſolete rufo-maculatis. Ent. ſyſt. 2. 133. 31. * *rauca.*
Habitat ad Cap. Bon. Spei. D. Vahl.

42. C. nigra, obſcura, immaculata, thorace margine antico ſubcornuto. Ent. ſyſt. 2. 133. 32. * *cornuta.*
Habitat ad Cap. Bon. Spei. D. Vahl.

43. C. nigra, albo-maculata, abdominis ſegmentis margine albis, clypeo emarginato, ſterno cornuto. Ent. ſyſt. 2. 133. 33. * *triſtis.*
*Cetonia triſtis.* Oliv. Inſ. 1. 6. 22. 20. tab. 18. fig. 91.
Habitat in America. Muſ. Dr. Hunter.

44. C. ferrugineo-flaueſcens, elytris vireſcentibus, ſterno cornuto. Ent. ſyſt. 2. 134. 34. * *ſmaragdula.*
*Cetonia ſmaragdula.* Oliv. Inſ. 1. 6. 73. 91. tab. 10. fig. 90.

Drury

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Drury Inf. 2. tab. 30. fig. 3.
Habitat in America. Muſ. Dr. Hunter.

*faſcicularis.* 45. C. thorace lineis quatuor albis, elytris viridibus, abdominis inciſuris barbatis. Ent. ſyſt. 2. 134. 35.
*Cetonia faſcicularis.* Oliv. Inſ. 1. 6. 16. 12. tab. 11. fig. 108.
*Scarabaeus faſcicularis.* Linn. ſyſt. Nat. 2. 575. 75. Muſ. Lud. Vlr. 26.
Drury Inf. 1. tab. 33. fig 2.
Voet. Col tab. 3. fig. 17.
Wulf. Inſ. Cap. tab. 1. fig. 1.
Habitat in Africa.

*aulica.* 46. C. viridis, nitida, thoracis margine elytrorumque maculis albis. Ent. ſyſt. 2. 134. 36. *
*Cetonia aulica.* Oliv. Inſ. 1. 6. 16. 11. tab. 2. fig. 6. 7.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*purpuraſcens.* 47. C. glabra, nigra, thorace elytrisque purpuraſcentibus. Ent. ſyſt. 2. 135. 37. *
*Cetonia thoracica*, nigra, thorace ferrugineo, elytris nigris purpuraſcentibus. Ent. ſyſt. 2. 138. 47. vix differt.
*Cetonia aurantia.* Herbſt. Col. 3. tab. 30. fig. 7.
Voet. Col. tab. 1. fig. 6.
Oliv. Inſ. 1. 6. tab. 3. fig. 11.
Habitat in India orientali. D. Prof. Abildgaard.

*Capenſis.* 48. C. hirta, rufa, albo-punctata. Ent. ſyſt. 2. 135. 38.
*Cetonia Capenſis.* Oliv. Inſ. 1. 6. 27. 28. tab. 6.
*Scarabaeus Capenſis.* Linn. ſyſt. Nat. 2. 556. 73. Muſ. Lud. Vlr. 30.
Drury Inſ. 1. tab. 33. fig. 3.
Voet. Col. tab. 2. fig. 11.

Roeſ.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Roeſ. Inſ. 2. ſcarab. 1. tab. B. fig. 6.
Degeer Inſ. 7. 640. 40. tab. 48. fig. 2.
Wulf. Inſ. Capenſ. tab. 1. fig. 2.
Habitat in Africa.

9. C. thorace nigro: margine albo, elytris teſta- *ſignata.*
ceis: ſutura margineque nigris. Ent. ſyſt. 2.
135. 39. *
*Cetonia ſignata.* Oliv. Inſ. 1. 6. 28. 29. tab. 5.
fig. 35.
Voet. Col. tab. 2. fig. 13.
Herbſt. Col. 3. tab. 23. fig. 8.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

0. C. glabra, atra, thoracis elytrorumque margini- *marginata*
bus rufis. Ent. ſyſt. 2. 136. 40. *
*Cetonia marginata.* Oliv. Inſ. 1. 6. 26. 26. tab.
5. fig. 34.
*Scarabaeus marginatus.* Degeer Inſ. 4. tab. 19.
fig. 10.
Drury Inſ. 2. tab. 32. fig. 1.
Voet. Col. tab. 1. fig. 4.
Habitat in Guineae Iatropha Gorcas.

1. C. glabra, obſcure purpuraſcens, thorace elytro- *ornata.*
rumque margine ſinuato, teſtaceo. Ent. ſyſt. 2.
136. 41. *
Habitat in Guinea. D. Iſert.

2. C. viridis, thoracis elytrorumque margine ferru- *marginel-*
gineo. Ent. ſyſt. 2. 136. 41. * *la.*
Habitat in Sierra Leon. D. Tunſtall.

3. C. nigra, lineola flaua a capite ad ſcutellum du- *lineola*
cta. Ent. ſyſt. 2. 137. 43. *
*Cetonia lineola.* Oliv. Inſ. 1. 6. 77. 98. tab. 5.
fig. 28.
*Scarabaeus lineola.* Linn. ſyſt. Nat. 2. 552. 53.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Roeſ. Inſ. 2. ſcarab. 1. tab. B. fig. 7.
Degeer Inſ. 4. tab. 19. fig. 5.
*Cetonia ephippium*, nigra, thoracis margine linea que dorſali, elytris macula difformi ferruginea Mant. Inſ. 1. 29. 31.
Oliv. Inſ. 1. 6. 78. 98. tab. 11. fig. 106.
Voet. Col. tab. 9. fig. 81.
Drury Inſ. 3. tab. 44. fig. 3.
Habitat in America meridionali.
Conf. *Scarabaeus Surinamenſis*. Linn. ſyſt. Nat. 2 552. 54.
Oliv. Inſ. 1. 6. 79. 99. tab. 11. fig. 104. vi differt.

*ſcutellata.* 54. C. atra, thoracis margine linea dorſali impreſſa ſcutelloque albis.
Habitat in Guinea. D. Meyer. Muſ. D. de Sehe ſtedt.
Magnitudo et ſtatura omnino C. lineolae. Capu atrum, linea baſi dentata, alba, clypeoque apice reflexo. Thorax ater, nitidus, margine late albo. Linea dorſalis impreſſa, alba, quae tamen nec anticum, nec poſticum marginem attingit. Scutellum paruum, album. Elytra laeuia, atra, immaculata. Abdominis vltimum ſegmentum prominet macula magna alba. Subtus atra, margine albo maculato.

*ſtriata.* 55. C. glabra, atra, nitida, thoracis margine teſtaceo: puncto nigro, clypeo emarginato. Ent. ſyſt. 2. 137. 44.
*Cetonia ſtriata.* Oliv. Inſ. 1. 6. 79. 100. tab. 11 fig. 102.
Habitat in Inſula Guadeloupe. D. Iſert.

*flauomaculata.* 56. C. obſcure brunnea, thorace punctis quatuor atris, elytris macula flexuoſa flaua. Ent. ſyſt. ſuppl. 129. 44.

Degeer

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer. Inſ. 7. tab. 47. fig. 19.
Oliv. Inſ. 1. 6. tab. 7. fig. 52.
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

57. C. fuſca, thorace elytrisque margine et punctis duobus flauis. Ent. ſyſt. 2. 138. 45. * *ſinuata.*
*Cetonia ſinuata.* Oliv. Inſ. 1. 6. 25. 24. tab. 8. fig. 78.
*Scarabaeus punctato-marginatus.* Degeer. Inſ. 7. 639. 38. tab. 47. fig. 20.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

58. C. nigra, thoracis margine elytrisque teſtaceis, elytris ſutura faſciaque dentata nigris. Ent. ſyſt. 2. 138. 46. * *faſciata.*
Habitat Alexandriae. D. Prof. Forſkahl.

59. C. teſtacea, thorace lineis punctisque duobus, elytris maculis quatuor transuerſis nigris. Ent. ſyſt. 2. 138. 48. * *oliuacea.*
*Cetonia oliuacea.* Oliv. Inſ. 1. 6. 37. 41. tab. 8. fig. 69.
*Scarabaeus cordatus.* Drury Inſ. 2. tab. 32.
Voet. Col. 1. tab. 1. fig. 7.
Habitat in Sierra Leon Africae. Muſ. D. Tunſtall.

60. C. thoracis margine lineaque dorſali, elytrorum faſciis interruptis teſtaceis. Ent. ſyſt. 2. 139. 49. * *interrupta*
*Cetonia interrupta.* Oliv. Inſ. 1. 6. 35. 42. tab. 8. fig. 70.
Habitat in Senegallia. Muſ. D. Olivier.

61. C. nigra, thorace quinquelineato, elytris acuminatis, albo-maculatis. Ent. ſyſt. 2. 139. 50. * *picta.*
Habitat in Oriente. D. Prof. Forſkahl.

62. C. nigra, thorace lineis tribus albis, elytris fa- *trilineata.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſcia flexuoſa ſcutelloque albis. Ent. ſyſt. 2. 139. 51. *
Habitat ad Cap. Bon. Spei. D. Lee.

*elata.* 63. C. atra, thorace lineis tribus albis, elytris oluaceis: margine atro albo-punctato. Ent. ſyſt. 2. 139. 52. *
Habitat in Guinea. D. v. Rohr.

*ſemipunctata.* 64. C. viridis, nitida, thorace quadrilineato, elytris baſi lineatis, apice punctatis. Ent. ſyſt. 2. 139. 53. *
Herbſt. Col. 3. tab. 30. fig. 11.
Voet. Col. tab. 22. fig. 152.
Habitat ad Cap. Bon. Spei. D. Boeber.

*ſ lineata.* 65. C. nigra, thorace quinquelineato, elytris albo maculatis, clypeo reflexo ſpinaque incumbente. Ent. ſyſt. 2. 104. 54. *
*Cetonia ſ lineata.* Oliv. Inſ. 1. 6. 56. 36. tab. 1 fig. 76.
Habitat in India. Muſ. D. Banks.

*atromaculata.* 66. C. nigra, nitida, elytris atro-maculatis, macula que poſtica alba. Ent. ſyſt. 2. 141. 55. *
Habitat Tranquebariae. Muſ. D. de Seheſtedt.

*-limbata.* 67. C. nigra, thoracis margine ferrugineo. Ent. ſyſt. 2. 141. 56. *
Habitat in Aegypto. D. Prof. Forſkahl.

*trigona.* 68. C. viridis, nitida, thoracis margine ſcutelliqui triangulo flauis.
*Cetonia trigona.* Herbſt. Col. tab. 26. fig. 12.
Voet. Col. tab. 10. fig. 84.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura et magnitudo C. Chryſidis. Caput viride nitidum, margine poſtico parum flaueſcente

Th

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Thorax aeneus, nitidus, margine laterali et poſtico flauo. Scutellum elongatum, triangulum, viride; triangulo ante marginem flauo. Elytra laeuia, nigra. Corpus ferrugineum.

69. C. viridis, nitidiſſima, elytrorum ſutura punctoque apicis atris. Ent. ſyſt. 2. 142. 58. * *elegans.*

*Cetonia elegans.* Oliv. Inſ. 1. 6. 29. 30. tab. 4. fig. 25.

Habitat in Coromandel. Muſ. D. Banks.

70. C. viridis, fuluo micans, elytris maculis duabus nigris, ſterno obtuſo. Ent. ſyſt. 2. 142. 59. * *4maculata.*

*Cetonia 4maculata.* Oliv. Inſ. 1. 6. 30. 31. tab. 8. fig. 73.

Habitat in Africa aequinoctiali. Muſ. D. Banks.

71. C. viridis, nitida, thorace elytrisque punctis duobus nigris, elytris punctis nigris ſtriatis. *6maculata.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Maior praecedente. Antennae nigrae. Caput viride, clypeo apice reflexo, emarginato. Thorax viridis, nitidus, punctis nigris, numeroſis, minutis impreſſis; maculisque duabus atris. Elytra viridia, nitida, ſtriis e punctis nigris impreſſis maculisque duabus atris, altera ad ſcutellum, altera verſus apicem. Corpus viride, nitidum, punctis nigris impreſſis abdominisque ſegmentis, apice atris, ſterno porrecto.

72. C. cuprea, nitida, thorace, ſcutello elytrisque viridibus. Ent. ſyſt. 2. 142. 60. * *cuprea.*

Habitat Surinami. Muſ. D. Tunſtall.

73. C. aenea, nitens, capitis ſpina incumbente, ſterno porrecto, elytris punctis nigris ſtriatis. Ent. ſyſt. 2. 142. 61. * *Africana.*

*Ceto-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cetonia Africana.* Oliv. Inſ. 1. 6. 31. 32. tab. 8. fig. 70.
*Scarabaeus Africanus.* Drury Inſ. 2. tab. 30. fig. 3.
Voet. Col. tab. 4. fig. 25.
Habitat in Africa, Sierra Leon.

*laeta.* 74. C. viridis, nitidiſſima, thorace punctato, elytris punctis nigris ſtriatis.
Habitat in India orientali. Muſ. D. Lund.
Nimis affinis videtur C. Africanae, at clypeus reflexus et omnino absque ſpina, et thorax punctis nigris impreſſus. Abdominis vltimum ſegmentum prominet lineolis numeroſis, transuerſis, vndatis, nigris. Sternum valde porrectum.

*ſuturalis.* 75. C. thorace nigro: margine rufo, elytris aeneo-nitidis: ſutura nigra. Ent. ſyſt. 2. 143. 62. *
*Cetonia ſuturalis.* Oliv. Inſ. 1. 6. 32. 34. tab. 8. fig. 74.
Habitat in Africa. D. Drury.

*vittata.* 76. C. hirta, viridi-aenea, elytris teſtaceis: vitta media nigra ſuturaque aenea. Ent. ſyſt. 2. 143. 63. *
*Melolontha Zebra.* Oliv. Inſ. 1. 5. 54. 73. tab. 7. fig. 75.
Voet. Col. tab. 7. fig. 60.
Habitat in Africa. Muſ. D. Banks.

*fulgida.* 77. C. aenea, nitidiſſima, abdominis vltimo ſegmento prominente, quadripunctato. Ent. ſyſt. 2. 144. 64. *
*Cetonia fulgida.* Oliv. Inſ. 1, 6. 33. 35. tab. 8. fig. 75.
Voet. Col. tab. 3. fig. 24.
Habitat in Carolina. Muſ. D. Boſc.

78. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

78. C. viridis, nitidiſſima, fuſco micans. Ent. ſyſt. 2. 144. 65. * *Iris.*

*Cetonia Iris.* Oliv. Inſ. 1. 6. 31. 33. tab. 8. fig. 77.
Voet. Col. tab. 4. fig. 25.
Habitat Surinami. Muſ. D. Banks.

79. C. atra, nitida, clypeo truncato, reflexo, ſterno obtuſo. Ent. ſyſt. 2. 144. 66. * *Gagates.*

*Cetonia Gagates.* Oliv. Inſ. 1. 6. 25. 25. tab. 4. fig. 20. tab. 11. fig. 20.
*Scarabaeus carbonarius.* Degeer Inſ. 4. 824. 18.
Forſt. Cent. Inſ. 6. 6.
Habitat in Sierra Leon. Muſ. D. Banks.

80. C. atra, ſcutello elytris dimidio breuiore, pedibus triunguiculatis: pollice fixo. Ent. ſyſt. 2. 144. 67. * *tetradactyla.*

*Cetonia tetradactyla.* Oliv. Inſ. 1. 6. 74. 93. tab. 2. fig. 8. tab. 7. fig. 53.
*Scarabaeus tetradactylus.* Linn. Mant. 530.
Drury Inſ. 1. tab. 33. fig. 7.
Sloan. Carn. 2. 237. fig. 2.
Habitat in America meridionali.

81. C. viridi-cuprea, nitida, elytris teſtaceis, ſterno antice cornuto. Ent. ſyſt. 2. 145. 68. * *clauata.*

Oliv. Inſ. 1. 6. 72. 89. tab. 8. fig. 58.
Habitat in America meridionali. Muſ. D. Banks.

82. C. atra, nitida, thoracis margine omni elytrorumque vittis duabus flauis. Ent. ſyſt. 2. 145. 69. * *fucata.*

*Cetonia 4vittata.* Oliv. Inſ. 1. 6. 92. 7. tab. 8. fig. 65.
Drury Inſ. 3. tab. 44. fig. 4.
Habitat in America meridionali. Muſ. D. Banks.

83. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*lurida.* 83. C. nigro-aenea, elytris lineis duabus elevatis alboque maculatis. Ent. syst. 2. 145. 70. *

*Cetonia lurida.* Oliv. Inf. 1. 6. 43. 50. t. 9. f. 81

Habitat in Brasilia. Mus. D. Banks

*mixta.* 84. C. obscura, cinereo variegata, scutello margine albo.

*Cetonia mixta.* Web. Inf. 69. 5.

Habitat in Sumatra. D. Daldorff.

Statura et magnitudo omnino C. acuminatae, at alia et distincta. Corpus supra fuscum, obscurum, vndique atomis cinereis variegatum. Scutellum margine albo. Elytra obtusa.

*lunulata* 85. C. thorace obscure viridi: margine albo, elytris acuminatis testaceis, macula nigra lunulisque albis.

*Cetonia ciliata.* Oliv. Inf. 1. 6. tab. 12. fig. 12. Web. Inf. 70. 7.

Habitat in Sumatra. D. Daldorff.

Statura praecedentium. Caput obscure viride. Thorax obscure viridis, margine late albo: puncto minuto viridi. Scutellum obscure viride. Elytra obscure testacea, macula vna alteraue nigra lunulisque albis. Corpus lanugine albida tectum.

*aurichalcea.* 86. C. cuprea, nitida, elytris acuminatis, albo-maculatis. Ent. syst. 2. 146. 71. *

*Cetonia aurichalcea.* Oliv. Inf. 1. 6. 42. 49. tab. 9. fig. 78.

Habitat ad Suratta Indiae. Mus. Britann.

*nitidula.* 87. C. viridis, testaceo-nitida, clypeo emarginato, elytris acuminatis. Ent. syst. 2. 146. 72. *

Habitat in Africa. D. Lee.

*Philippensis.* 88. C. aenea, nitida, thorace margine punctisque duobus albis, elytris acuminatis, maculatis. Ent. syst. 2. 146. 73. *

*Ceto-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cetonia Philippenſis.* Oliv. Inſ. 1. 6. 34. 37. tab. 10. fig. 97.
Petiv. Gazoph. tab. 26. fig. 9.
Habitat in Inſulis Philippinis. Muſ. Britann.

89. C. viridi-fuſca, thorace margine punctisque, elytris acuminatis: faſcia media maculisque niueis. Ent. ſyſt. 2. 146. 74. * *ſtolata.*
*Cetonia ſtolata.* Oliv. Inſ. 1. 6. 43. 51. tab. 7. fig. 59.
Habitat in noua Hollandia. Muſ. D. Banks.

90. C. obſcure viridis, albo-irrorata, elytris acuminatis, teſtaceo-lineatis. *atomaria.*
Habitat in China. Muſ. D. Lund.
Statura et magnitudo C. feſtiuae. Caput, thorax, elytra obſcure viridia, atomis numeroſiſſimis, albis irrorata. Elytra apice ad ſuturam in ſpinam acutam, compreſſum excurrunt. Pectus albo-villoſum. Abdomen aeneum, albo-maculatum.

91. atra, albo-maculata, elytris acuminatis: vitta lata, teſtacea. Ent. ſyſt. 2. 147. 75. * *feſtiua.*
Habitat Tranquebariae. Muſ. D. Lund.

92. C. atra, nitida, coleoptris faſcia lata, interrupta, flaua. Ent. ſyſt. 2. 147. 76. * *abbreuiata*
Habitat in Senegallia. Muſ. D. Geoffroy.

93. C. aenea, nitida, thorace vtrinque macula elytris plurimis albis, ſterno obtuſo. Ent. ſyſt. 2. 147. 77. * *maculata.*
*Cetonia maculata.* Oliv. Inſ. 1. 6. 36. 40. tab. 7. fig. 66.
Voet. Col. tab. 1. fig. 8.
Habitat in Coromandel. Muſ. D. Banks.

94. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*difformis.* 94. C. cyanea, thorace vtrinque macula vna difformi, elytris plurimis albis.
Habitat in India orientali. Muſ. D. de Seheſtedt.
Vix a C. maculata ſatis diſtincta. Differt tamen colore cyaneo elytrisque nullo modo acuminatis.

*acuminata* 95. C. obſcure aenea, pallido maculata, elytris acuminatis. Ent. ſyſt. 2. 148. 78. *
*Cetonia acuminata.* Oliv. Inſ. 1. 6. 42. 48. tab. 8. fig. 72.
Voet. Col. tab. 5. fig. 37.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*mandarina* 96. C. atra, thorace elytrisque acuminatis, albo-maculatis.
*Cetonia mandarina.* Web. Inſ. 68. 4.
Habitat in Sumatra. D. Daldorff.
Nimis affinis C. acuminatae, at diſtincta, colore toto atro.

*haemorrhoidalis.* 97. C. nigra, elytris viridibus, nitidis, thoracis margine anoque rufis. Ent. ſyſt. 2. 148. 79. *
*Cetonia haemorrhoidalis.* Oliv. Inſ. 1. 6. 55. 66. tab. 4. fig. 24. tab. 11. fig. 24.
Degeer. Inſ. 7. 642. 43.
Herbſt. Arch. tab. 43. fig. 8.
Habitat ad Cap. Bon. Spei. Muſ. Dr. Hunter.

*adſperſa.* 98. C. ferruginea, elytris viridibus, albo-punctatis.
Habitat ad Cap. Bon. Spei. Muſ. D. de Seheſtedt.
Statura parua C. haemorrhoidalis. Caput et thorax obſcure rufa, albo-punctata. Elytra ſtriata, viridia, albo-punctata, margine vittaque ferrugineis. Corpus ferrugineum.

*areata.* 99. C. nigra, pubeſcens, elytris diſco rufo. Ent. ſyſt. 2. 148. 80. *

*Cete-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cetonia areata.* Oliv. Inſ. 1. 6. 48. 58. tab. 9. fig. 82.
Habitat in Virginia. Muſ. Britann.

100. C. nigricans, hirta, elytris pallido-maculatis, thorace carinato. Ent. ſyſt. 2. 148. 81. *birta.*
*Cetonia birta.* Oliv. Inſ. 1. 6. 52. 63. tab. 8. fig. 36. 44.
*Scarabaeus hirtellus.* Linn. ſyſt. Nat. 2. 556. 69.
Panz. Fn. Germ. 1. tab. 3.
Schaeff. Icon. tab. 198. fig. 8.
Voet. Col. tab. 4. fig. 34.
β. *Scarabaeus ſqualidus.* Linn. ſyſt. Nat. 2. 556. 68.
Voet. Col. tab. 4. fig. 33.
Habitat in Europa auſtraliori.

101. C. glabra, nigra, thorace elytrisque albo-punctatis. Ent. ſyſt. 2. 149. 82. * *funeſta.*
Habitat in Italia. Dr. Allioni.

102. C. clypeo emarginato, nigra, albo-maculata, abdomine ſubtus punctis quatuor albis. Ent. ſyſt. 2. 149. 83. *ſtictica.*
*Cetonia ſtictica.* Oliv. Inſ. 1. 6. 53. 64. tab. 7. fig. 57.
*Scarabaeus ſticticus.* Linn. ſyſt. Nat. 2. 552. 54.
Geoff. Inſ. 1. 79. 14.
Herbſt. Arch. tab. 19. fig. 27.
Voet. Col. tab. 22. fig. 82.
Panz. Fn. Germ. 1. tab. 4.
Payk. Fn. Sv. 2. 205. 4.
Degeer. Inſ. 4. 301. 29. tab. 10. fig. 22.
~~Herbſt. Col. tab. 80. fig. 5.~~
Habitat in Europae Carduis frequens.

103. C. thorace ferrugineo; punctis quatuor nigris, *albo-punctata.*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

gris, elytris ferrugineis: limbo atro, albo-punctatis. Ent. syst. suppl. 129. 89. *
Habitat ad Cap. Bon. Spei. Mus. D. Lund.

*cinerascens* 104. C. thoracis dorso atro: linea cinerascente, elytris cinereis nigro-maculatis. Ent. syst. suppl. 129. 89. *
Habitat ad Cap. Bon. Spei. Mus. D. Lund.

*irrorata.* 105. C. nigra, elytris striatis, ferrugineis. nigro-maculatis. Ent. syst. suppl. 130. 89. *
Habitat ad Cap. Bon. Spei. Mus. D. de Seheſtedt.

*furuata.* 106. C. thorace elytrisque testaceis, nigro-punctatis. Ent. syst. suppl. 130. 89. *
Habitat ad Cap. Bon. Spei. Mus. D. de Seheſtedt.

*sepulcralis.* 107. C. clypeo reflexo, integro, tomentosa, fusca, elytris albo-maculatis.
Habitat in Carolina. Mus. D. Bosc.
Statura et magnitudo omnino C. sticticae, at omnino distincta. Clypeus reflexus, integer. Caput et thorax tomentosa, fusca, immaculata. Elytra substriata, fusca, albo obscuro maculata. Abdomen obscure aeneo-nitidulum, maculis aliquot albis, lateralibus.

*punctulata* 108. C. cinerea, elytrorum striis albo-punctatis. Ent. syst. 2. 149. 84. *
*Cetonia punctata.* Oliv. Inf. 1. 6. 54. 64. tab. 6. fig. 47.
Habitat in Senegallia. Mus. D. Geoffroy.

*floralis.* 109. C. glabra, nigra, thoracis, elytrorum abdominisque margine albo. Ent. syst. 2. 149. 85. *
Habitat in Africae floribus compositis. D. Prof. Vahl.

*14maculata* 110. C. capite thoraceque aureis, elytris punctis quatuordecim albis. Ent. syst. 2. 150. 86. *
Habitat - - Mus. D. Lee.

111.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

111. C. nigra, thorace rufo, nigro-maculato, elytris nigris, albo-punctatis: dorso rufo. Ent. syst. 2. 150. 87. * *verſicolor.*

*Cetonia verſicolor.* Oliv. Inſ. 1. 6. 46. 54. tab. 4. fig. 23.

*Scarabaeus cruentus.* Pall Icon. tab. B. fig. 24.

Herbſt. Arch. tab. 19. fig. 28.

Acta Berol. tab. 1. fig. 8.

Habitat in Aegypto. D. Prof. Forſkahl.

112. C. atra, thorace, margine elytrisque maculis albis ſparſis. Ent. ſyſt. 2. 151. 88. * *variegata.*

*Cetonia variegata.* Oliv. Inſ. 1. 6. 47. 56. tab. 5. fig. 30. 31. b.

*Scarabaeus albellus.* Pall. Icon. 1. 17. tab. A. fig. 18.

Herbſt. Arch. tab. 19. fig. 29.

Habitat Tranquebariae. Dr. Koenig.

113. C. atra, thoracis margine poſtico ſanguineo, elytris flauo-maculatis. *torquata.*

Habitat - - - Dr. Pflug. Muſ. D. Lund.

Statura depreſſa C. aequinoctialis. Caput atrum, clypeo emarginato. Thorax ater, margine poſtico late ſanguineo. Scutellum laeue, anguſtum. Elytra ſubſtriata, atra, punctis duobus parcus, approximatis, ad baſin maculisque duabus pone medium flauis. Corpus nigrum, ferrugineo flauoque maculatum.

114. C. atra, thoracis margine albo, coleoptris teſtaceis: macula ſcutellari apiceque atris. Ent. ſyſt. 2. 151. 89. * *aequinoctialis.*

*Cetonia aequinoctialis.* Oliv. Inſ. 1. 6. 50. 60. tab. 6. fig. 42.

Habitat in Senegallia. Muſ. D. Olivier.

115.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſanguinolenta.* 115. C. atra, elytris acuminatis: macula magna marginali fulua. Ent. ſyſt. 2. 151. 90. *

*Cetonia ſanguinolenta.* Oliv. Inſ. 1. 6. tab. 6 fig. 41.

Habitat in Senegallia. Muſ. D. Geoffroy.

*diſcoidea.* 116. C. atra, elytris fuluis: macula ſcutellari apiceque atris, faſcia mediaque purpurea. Ent. ſyſt. 2. 151. 91. *

Habitat in Bengalia. D. de Paykull.

*lugubris.* 117. C. glabra, atra, elytris macula laterali anoque albis. Ent. ſyſt. 2. 152. 92. *

*Cetonia lugubris.* Oliv. Inſ. 1. 6. 44. 52. tab. 7. fig. 60.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*biſtrio.* 118. C. teſtacea, capite thoracis lineis duabus, elytrorum ſutura maculisque tribus nigris. Ent. ſyſt. 2. 152. 93. *

*Cetonia biſtrio.* Oliv. Inſ. 1. 6. 45. 53. tab. 10. fig. 94.

Habitat in Aegypto. D. Prof. Forſkahl.

*modeſta.* 119. C. ſupra viridis, opaca, albo-punctata, capite emarginato atro. Ent. ſyſt. 2. 152. 94. *

Habitat Tranquebariae. Muſ. D. de Seheſtedt.

*glorioſa.* 120. C. teſtacea, nitida, lineis abbreuiatis nigris. Ent. ſyſt. 2. 153. 95. *

*Melolontha Dorcyi.* Oliv. Inſ. 1. 5. 34. 38. tab. 4 fig. 41.

Habitat in Inſula St. Domingo.

*maura.* 121. C. depreſſa, atra, thorace poſtice punctis duobus impreſſis.

Habitat in Guinea. Muſ. D. Lund.

Statura elongata, depreſſa C. hottentottae. Caput nigrum, clypeo obtuſo. Thorax planus, ſubpuncta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

punctatus, ater, poſtice punctis duobus impreſſis. Elytra ſubſtriata, nigra, nitida. Corpus nigrum, ſterno haud porrecto.

Statura omnino C. hottentottae; forte proprii generis, mihi haud rite examinata.

122. C. atra, glabra, elytris poſtice punctis duobus albis. Ent. ſyſt. 2. 153. 96. * *hottentotta.*

*Cetonia hottentotta.* Oliv. Inſ. 1. 6. 56. 68. tab. 7. fig. 55.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

123. C. atra, thorace elytrisque ſulcis villoſis albis. *regia.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Media, reliquis magis oblonga, antice poſticeque attenuata. Caput nigrum, clypeo apice emarginato antennisque teſtaceis. Thorax planus, punctatus, ater, ſulcis tribus dorſalibus, villoſo-albis: lateralibus parum abbreviatis. Scutellum impreſſum, villoſo-album. Elytra plana, ſulcis duobus baſeos, longiore ad ſuturam, verſus apicem incuruo, maculaque pone medium impreſſis, albo-villoſis. Puncta praeterea quatuor in margine. Corpus atrum, lateribus flauo-villoſis. Pedes rufi.

124. C. glabra, atra, thorace poſtice maculis duabus elytrisque ſanguineis. Ent. ſyſt. 2. 153 97. * *cruenta.*

*Cetonia cruenta.* Oliv. Inſ. 1. 6. 57. 69. tab. 6. fig. 37. tab. 7. fig. 58.

Habitat - - Muſ. D. Banks.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

127. MELOLONTHA. *Maxilla* breuis, cornea, apice multidentata: dentibus ſimplicibus, acutis.
*Antennae* lamellatae.

*ſtigma.* 1. M. fuſco-ferruginea, elytris ſtigmate oblongo apicis albo. ~~Ent. ſyſt. ſuppl. 131. 1.~~
~~Drury Inſ. 3. tab. 44. fig. 1.~~
~~Herbſt. Col. 3. tab. 28. fig. 1.~~
Habitat in Iaua.

*alba.* 2. M. farinoſo-albida.
~~*Melolontha alba.* Oliv. Inſ. 1. 5. 10. 2. tab. 8. fig. 1.~~
~~*Scarabaeus albus* Pall. Icon. 1. tab. B. fig. 21.~~
*Melolontha alba.* Web. Inſ. 71. 1.
Habitat in Sumatra. D. Daldorff.
Statura omnino praecedentis, tota albido-farinoſa, immaculata.

*Fullo.* 3. M. teſtacea, albo-maculata, ſcutello macula duplici, antennis heptaphyllis. Ent. ſyſt. 2. 154. 1.
*Scarabaeus Fullo.* Linn. ſyſt. Nat. 2. 553. 57.
Geoff. Inſ. 1. 69. 2.
Oliv. Inſ. 1. 5. 9. 1. tab. 3. fig. 28.
Schaeff. Icon. tab. 23. fig. 2.
Petiv. Gazoph. tab. 138. fig. 1.
Voet. Scarab. tab. 6. fig. 48. 49.
Petagna Inſ. Calabr. 3. 11. tab. 1. fig. 1.

Friſh.

---

Melolonthae corpus oblongum, gibbum, tomentoſum, laeue, immarginatum, tardum, *capite* ouato, inſerto, labio rotundato; *oculis* globoſis, marginalibus; *antennis* diſtantibus, thorace breuioribus, ſubter oculos inſertis; *thorace* transuerſo; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis; *tibiis* dentatis; *tarſis* quinquearticulatis; *colore* variegato.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Friſh. Inſ. 11. 22. tab. 1. fig. 1.
Roeſ. Inſ. 4. tab. 30.
Aub. Miſcell. tab. 16. fig. med.
Payk. Fn. Sv. 2. 206. 1.
Degeer Inſ. 4. 272. 19. tab. 10. fig. 14.
Herbſt. Col. 3. tab. 22. fig. 1. 2.
Habitat in Europae ſyluis arenoſis.

4. M. fuſca, atomis numeroſiſſimis griſeis. *rorida.*
*Melolontha rorida.* Web. Inſ. 71. 2.
Habitat in Sumatra. D. Daldorff.
Paullo minor, quam M. Fullo. Corpus totum fuſcum, obſcurum, punctatum, et in ſingulo puncto pilus breuiſſimus, griſeus, vnde corpus punctis numeroſiſſimis griſeis irroratum videtur. Pili ſaepius detriti, et tunc totum corpus fuſcum apparet.

5. M. obſcure teſtacea, clypeo emarginato, thoracis margine ſerrato. Ent. ſyſt. 2. 155. 2. * *ſerrata.*
*Melolontha ſerrata.* Oliv. Inſ. 1. 5. 11. 4. tab. 1. fig. 5.
Habitat in Coromandel. Muſ. D. Banks.

6. M. teſtacea, thorace villoſo, inciſuris abdominis albis. Ent. ſyſt. 2. 155. 3. *vulgaris.*
*Scarabaeus Melolontha.* Linn. ſyſt. Nat. 2. 554. 60. Fn. Sv. 392.
Geoff. Inſ. 1. 70. 1.
Oliv. Inſ. 1. 5. 12. 5. tab. 1. fig. 1.
Petiv. Gazoph. 29. tab. 19. fig. 2.
Roeſ. Inſ. 2. ſcarab. 1. tab. 1.
Degeer Inſ. 4. tab. 10. fig. 14.
Voet. Scarab. tab. 6. fig. 45. 46.
Schaeff. Icon. tab. 93. fig. 1. 2.
tab. 102. fig. 3. 4.
Habitat in Europae borealis graminoſis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Variat* thorace rufo et nigro, et vtraque varietas alternat quotannis.

Animalculum inter omnia noſtratia maxime fatale.

*Larua* mollis, griſea, capite pedibusque teſtaceis, per plures annos ſub terra degit, et deuorat radices vegetabilium, agros prataque deſolans. *Imago* veſperi volitans cum ſuſurro deuaſtat folia arborum, tilia intacta.

Deuoratur a ſuibus, coruis, cornicibus, picis, gallinis, meleagridibus, carabis maioribus. Per diem arboribus concuſſis facillime colligitur a pueris.

*Hippocaſtani.* 7. M. nigra, thorace villoſo, elytris teſtaceis, abdomine maculis marginalibus albis.

~~*Melolontha Hippocaſtani.* Oliv. Inſ. 1. 5. 13. tab. 1. fig. 3. a. b. c.~~

Habitat in Aeſculo hippocaſtano Europae auſtralioris.

Nimis affinis M. vulgari. Caput nunc nigrum, nunc rufum, villoſum, reflexum. Thorax villoſus, nunc niger, nunc rufus. Elytra teſtacea. Corpus nigrum, abdomine maculis magnis, marginalibus, triangularibus, albis. Pedes nunc ~~albi~~, nunc nigri.

*villoſa.* 8. M. teſtacea, clypeo marginato, reflexo, corpore ſubtus lanato, ſcutello albo. Ent. ſyſt. 2. 156. 4.

*Scarabaeus ſepicola.* Linn. ſyſt. Nat. 2. 156. 17.

*Scarabaeus cerealis.* Scopoli Flor. Inſ. 1. 49. tab. 21. fig. B.

Oliv. Inſ. 1. 5. 13. 6. tab. 1. fig. 4.

Panz. Fn. Germ. 31. tab. 19.

~~Herbſt. Col. 3. tab. 22. fig. 8.~~ (*pilosa*)

Habitat in Europa auſtraliori. Dr. Allioni.

*pilosa.* 9. M. villoſa, nigricans, clypeo reflexo, ſubemarginato, antennis ferrugineis. Ent. ſyſt. 2. 156. 5.

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 31. tab. 20.
Habitat in Hungaria. Muſ. D. de Seheſtedt.

10. M. thorace pubeſcente, teſtacea, elytris lineis quatuor niueis. Ent. ſyſt. 2. 156. 6. * *occidentalis.*
~~*Scarabaeus occidentalis.* Linn. ſyſt. Nat. 2. 555. 64.~~
Oliv. Inſ. 1. 5. 14. 7. tab. 1. fig. 7.
~~Drury Inſ. 2. tab. 32. fig. 2.~~
Habitat in Europa auſtraliori.

11. M. glabra, teſtacea, clypeo emarginato. Ent. ſyſt. 2. 156. 7. * *feruida.*
~~Oliv. Inſ. 1. 5. 24. 21. tab. 9. fig. 109.~~
Habitat in America ſeptentrionali.

12. M. villoſa, teſtacea, clypeo reflexo: linea media eleuata transuerſa. *transuerſa.*
Habitat in Auſtria. Dom. de Meyerle.
Statura et magnitudo M. feruidae. Corpus ſupra teſtaceum, ſubtus villoſum, pedibus ferrugineis. Clypeus reflexus, ſubemarginatus, in medio linea transuerſa, eleuata, vti in ſcarabaeis feminiinis.

13. M. glabra, teſtacea, clypeo reflexo. Ent. ſyſt. 2. 156. 8. * *reflexa.*
~~Oliv. Inſ. 1. 5. 24. 22. tab. 4. fig. 32.~~
Habitat in Africa. Muſ. D. Banks.

14. M. fuluo-hirta, clypeo reflexo emarginato, elytris glabris nigris. Ent. ſyſt. 2. 157. 9. * *Alopex.*
*Melolontha Alopex.* Oliv. Inſ. 1. 5. 18. 9. tab. 4. fig. 35.
~~*Scarabaeus vertumnus.* Pall. Itin. 1. 10. 28.~~
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

15. M. cinereo-hirta, elytris nigris, clypeo reflexo, integro. Ent. ſyſt. 2. 157. 10. * *tomentoſa.*
Habitat in India orientali. D. Lee.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſolſtitialis.* 16. M. teſtacea, thorace villoſo, elytris luteo-pallidis: lineis tribus pallidioribus. Ent. ſyſt. 2. 157. 11.

*Scarabaeus ſolſtitialis.* Linn. ſyſt. Nat. 2. 554. 61. Fn. Sv. 393.

Oliv. Inſ. 1. 5. 16. 10. tab. 2. fig. 8. 11.

Friſh. Inſ. 9. 30. tab. 15. fig. 3.

Schaeff. Icon. tab. 93. fig. 3.

Voet. Scarab. tab. 6. fig. 51. 52.

Degeer. Inſ. 4. tab. 10. fig. 15.

Payk. Fn. Sv. 2. 208. 3.

Herbſt. Col. 3. tab. 22. fig. 9.

Habitat in Europae arboretis.

*bidens.* 17. M. atra, antennis pedibusque teſtaceis, clypeo bidentato, elytris ſtriatis.

Habitat in Carolina.

Paullo maior M. atra. Caput nigrum, clypeo bidentato. Antennae teſtaceae. Thorax punctatus, ater. Elytra crenato-ſtriata, nigra. Pectus ferrugineo-hirtum. Pedes ferrugínei.

*candida.* 18. M. clypeo rotundato, corpore villoſo, albo. Ent. ſyſt. 2. 157. 12.

*Melolontha candida.* Oliv. Inſ. 1. 5. 15. 8. tab. 8. fig. 98.

Habitat in India orientali. D. Lee.

*atra.* 19. M. clypeo reflexo, villoſa, nigra, antennis ferrugineis. Ent. ſyſt. 2. 158. 13. Cat. 3844 37

*Melolontha atra.* Herbſt. Arch. tab. 24. fig. 1.

N. 596 *Melolontha fuſca.* Oliv. Inſ. 1. 5. 99. 13. tab. 2. fig. 10.

Habitat in Europa auſtraliori. D. Prof. Vahl.

*aequinoctialis.* 20. M. villoſa, teſtacea, clypeo ſubreflexo. Ent. ſyſt. 2. 158. 14.

*Melolontha aequinoctialis.* Herbſt. Arch. tab. 22. fig. 11. Col. 3. 62. 10.

Melo

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Melolontha aestiva.* Oliv. Inf. 1. 5. tab. 2. fig. 9. a. b.

Habitat in Hungaria. D. Prof. Helwig.

21. M. nigra, clypeo, thoracis lateribus, elytris pedibusque pallidis. Ent. syst. 2. 158. 15. * *Pini.*

*Melolontha Pini* Oliv. Inf. 1. 5. tab. 2 fig. *q. a b.*

Habitat in Barbaria. Mus. D. Desfontaines.

22. M. oblonga, glabra, testacea, thorace vtrinque macula abdomineque flauis. Ent. syst. 2. 158. 16. * *2maculata*

Oliv. Inf. 1. 5. 33. tab. 5. fig. 45.

Habitat in China. Mus. D. Banks.

23. M. oblonga, villosa, pallida, elytris sutura apiceque nigris, clypeo reflexo. Ent. syst. 2. 159. 17. * *atriplicis.*

Oliv. Inf. 1. 5. 28. 28. tab. 8. fig. 99.

Habitat in Barbariae Atriplice humilifolia. Prof. Vahl.

24. M. oblonga, glabra, nigra, elytris obsolete striatis. Ent. syst. 2. 159. 18. * *oblonga.*

*Scarabaeus oblongus.* Scopol. Carn. 10.

Oliv. Inf. 1. 5. 19. 14. tab. 4. fig. 34.

Habitat in Germania. *est variet. nigra* Fn. Suec.

25. M. villosa, fusca, elytris testaceis, abdomine albicante. Ent. syst. 2. 159. 19. * *ruficornis.*

Voet. Scarab. tab. 6. fig. 50. *marginata, Herbst. tab. 24. f. 2.*

Habitat in Germania. D. de Hattorff.

26. M. subtus lanata, capite thoraceque aureis, elytris luteis. Ent. syst. 2. 159. 20. * *lanigera.*

*Scarabaeus laniger.* Linn. syst. Nat. 2. 555. 67. Mus. Lud. Vlr. 20.

Oliv. Inf. 1. 5. 21. 17. tab. 4. fig. 39.

Drury Inf. 1. tab. 34. fig. 6.

Voet. Col. tab. 7. fig. 59.

Habitat in India.

27. M.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*longicornis.* 27. M. ſupra glabra, nigra, elytris ferrugineis, antennarum claua elongata. Ent. ſyſt. 2. 160. 21.
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

*punctata.* 28. M. teſtacea, elytris punctis tribus fuſcis, diſtantibus. Ent. ſyſt. 2. 160. 22.
*Scarabaeus punctatus.* Linn. ſyſt. Nat. 2. 557. 76. Muſ. Lud. Vlr. 23.
Oliv. Inſ. 1. 5. 22. 18. tab. 1. fig. 6.
Drury Inſ. 1. tab. 34. fig. 6.
Gronov. Gazoph. 438.
Voet. Scarab. tab. 8. fig. 65.
Habitat in America.

*viridis.* 29. M. glabra, ſupra viridis, ſubtus aurea. Ent. ſyſt. 2. 160. 23. *
Oliv. Inſ. 1. 5. 29. 31. tab. 3. fig. 21. b. fig. 18.
Habitat in India orientali. Muſ. D. Banks.

*aenea.* 30. M. glabra, aenea, abdominis lateribus albis, elytris acuminatis. Ent. ſyſt. 2. 160. 24. *
Habitat in noua Hollandia. Muſ. D. Banks.

*futuralis.* 31. M. viridis, elytrorum futura flaueſcente. Ent. ſyſt. 2. 161. 25. *
Oliv. Inſ. 1. 5. 31. 34. tab. 7. fig. 85.
Habitat in noua Zelandia. Muſ. D. Banks.

*bicolor.* 32. M. glabra, viridis, ſubtus teſtacea, pedibus apice aureis. Ent. ſyſt. 2. 161. 26. *
Oliv. Inſ. 1. 5. 30. 33. tab. 7. fig. 86.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*geminata.* 33. M. glabra, atra, thorace punctato, elytris punctatis: ſtriis geminatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura omnino M. barbatae. Caput laeue, atrum, niti-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nitidum. Thorax punctatus, ater. Elytra punctata, ſtriisque per paria approximatis notata. Corpus atrum.

34. M. glabra, nigra, nitida, abdominis vltimo ſegmento prominulo, barbato. Ent. ſyſt. 2. 161. 27. * *barbata.*

Habitat in India orientali.

35. M. glabra, atra, thorace punctato, elytris ſubſtriatis. Ent. ſyſt. ſuppl. 131. 27. * *Horia hottentotta*

Habitat in Nympheae floribus Indiae orientalis. Muſ. D. Lund.

36. M. glabra, nitida, caſtanea; elytris laeuibus, teſtaceis. *caſtanea.*

*Melolontha caſtanea.* Oliv. Inſ. 1. 5. tab. 10. fig. 124.

Habitat in America meridionali. Muſ. D. de Seheſtedt.

Statura omnino M. barbatae. Corpus totum caſtaneum, nitidiſſimum, elytris laeuibus, teſtaceis. Pedes antici parum elongati, vngue validiſſimo.

*Variat* corpore atro, nitidiſſimo.

37. M. glabra, atra, thorace punctato, elytris punctato-ſtriatis, ferrugineis. *ruſipennis.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Praecedentibus affinis. Thorax punctatus, ater, nitidus. Elytra punctato-ſtriata, ferruginea.

38. M. glabra, teſtacea, capite elytrorum ſutura tibiisque poſticis nigris, clypeo reflexo. Ent. ſyſt. 2. 161. 28. * *elata.*

Habitat Tranquebariae. Muſ. D. de Seheſtedt.

39. M.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*dorſalis.* 39. M. glabra, teſtacea, capite, thoracis coleoptrorumque dorſo nigris. Ent. ſyſt. 2. 162. 29. *
Herbſt. Arch. tab. 22. fig. 12.
Habitat Tranquebariae. Dr. Koenig.

*glacialis.* 40. M. glabra, picea, antennis pedibusque teſtaceis. Ent. ſyſt. 2. 162. 30. *
Oliv. Inſ. 1. 5. 78. 46. tab. 6. fig. 61.
Habitat in Terra del Fuego. Muſ. D. Banks.

*lurida.* 41. M. thorace nigro: margine pallido, elytris griſeis, nigro-punctatis. Ent. ſyſt. 2. 162. 31. *
Habitat - - Muſ. D. Tunſtall.

*ſtriata.* 42. M. glabra, aenea, elytris ſtriatis: ſutura lineisque quatuor cupreis. Ent. ſyſt. 2. 162. 32. *
Oliv. Inſ. 1. 5. 39. 49. tab. 6. fig. 65.
Habitat in Terra del Fuego. Muſ. D. Banks.

*triſtis.* 43. M. thorace villoſo, obſcuro, elytris laeuibus, teſtaceis, ſcutello albo. Ent. ſyſt. 2. 162. 33. *
Habitat in America boreali. Muſ. D. Blackburn.

*teſtacea.* 44. M. glabra, teſtacea, capite obſcuriore, elytris ſtriatis. Ent. ſyſt. 2. 163. 34. *
Oliv. Inſ. 1. 5. 39. 47. tab. 5. fig. 49.
Habitat in Terra del Fuego. Muſ. D. Banks.

*birticollis.* 45. M. hirta, nigra, elytris punctatis, glabris, rufis. Ent. ſyſt. 2. 163. 35. *
Habitat in Africa. D. Prof. Vahl.

*pallida.* 46. M. glabra, teſtacea, capite elytrorumque ſutura nigris. Ent. ſyſt. 2. 163. 36. *
*Melolontha ſpinipes.* Herbſt. Arch. 13. 4. tab. 19. fig. 21.
~~Oliv. Inſ. 1. 5 25. 24. tab. 9. fig. 103.~~
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.
*Variat* minor elytris ſtriatis.

47. M.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

47. M. glabra, punctata, ferrugiuea, elytris teſtaceis. Ent. ſyſt. 2. 163. 37. * *ruficollis.*
Oliv. Inſ. 1. 5. 27. 26. tab. 9. fig. 111.
Habitat in Coromandel. Dr. Koenig.

48. M. glabra, obſcura, elytris teſtaceis, clypeo quadridentato. Ent. ſyſt. ſuppl. 131. 37. * *quadridens.*
Habitat in India orientali. D. Hybner.

49. M. villoſa, nigra, thorace elytrisque varioloſis. Ent. ſyſt. 2. 164. 38. * *varioloſa.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

50. M. teſtacea, ſupra albo-villoſa, elytris lineis tribus eleuatis. *lanata.*
Habitat in Isle de France. D. Daldorff. Muſ. D. de Seheſtedt.
Statura et magnitudo omnino M. ſignatae. Capitis clypeus rotundatus, integer, albo-villoſus. Thorax teſtaceus, albo-villoſus: linea dorſali eleuata. Elytra albo-villoſa, lineis tribus eleuatis. Corpus teſtaceum.

51. M. glabra, pallida, thorace lineis duabus, elytris maculis tribus nigris. Ent. ſyſt. 2. 164. 39. * *ſignata.*
*Scarabaeus nigrocephalus.* Degeer Inſ. 4. tab. 19. fig. 6.
~~Oliv. Inſ. 1. 5. tab. 4. fig. 33.~~
Voet. Col. tab. 9. fig. 77.
Sloane Iamaic. 206. tab. 237. fig. 38.
Habitat in Iamaica. Muſ. D. Banks.
*Variat* thorace immaculato, aut elytris punctis duobus, aut nullis.

52. M. glabra, aenea, elytris teſtaceis: ſutura margineque aeneis. Ent. ſyſt. 2. 164. 40. * *marginata.*
Habitat in America meridionali. Muſ. D. Banks.

53. M. glabra, coeruleo-nigra, elytris teſtaceis: ſutura ferruginea. M. coeruleocephala. Ent. ſyſt. 2. 164. 41. * *cyanocephala.*
Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 1. 5. 36. 43. tab. 6. fig. 59.
Habitat - - Muſ. D. Banks.

*brunnea.* 54. M. glabra, teſtacea, elytris ſtriatis, thorace vtrinque puncto notato. Ent. ſyſt. 2. 165. 42. *
*Scarabaeus brunneus.* Linn. ſyſt. Nat. 2. 556. 72.
Oliv. Inſ. 1. 5. 43. 55. tab. 4. fig. 33.
Geoff. Inſ. 1. 83. 22.
Degeer Inſ. 4. tab. 10. fig. 17.
Roſſ. Fn. Etr. 1. 22. 50.
Voet. Scarab. tab. 7. fig. 53. 54.
~~Sulz. Hiſt. Inſ. tab. 1. fig. 9.~~
~~Fn. Sv.~~ 1v. 6 Payk. Fn. Sv. 2. 209. 4.
Herbſt Col. 3. tab. 24. fig. 3.
Habitat in Europae Arboretis.
*Larua* mollis, albida, capite pedibusque teſtaceis, deuorat radices, *imago* folia.

*melanocephala.* 55. M. glabra, rufeſcens, capite nigro, elytris pallidis. Ent. ſyſt. 2. 165. 43. *
~~Oliv. Inſ. 1. 5. 43. 55. tab. 7. fig. 83.~~
Voet. Scarab. tab. 9. fig. 80.
Habitat in Braſilia. Muſ. D. Banks.

*ferruginea nigriceps* 56. M. glabra, punctata, rufa, capite nigro. Ent. ſyſt. ſuppl. 131. 43. *
Habitat in Cayenna. D. Richard.

*pallens.* 57. M. glabra, pallida, capitis baſi thoracisque punctis duobus nigris. Ent. ſyſt. ſuppl. 131. 44. *
Habitat in Cayenna. D. Richard.

*erythrocephala.* 58. M. glabra, pallida, capite rufo, baſi nigro. Ent. ſyſt. 2. 165. 44. *
Oliv. Inſ. 1. 5. 41. 51. tab. 7. fig. 80.
Habitat in Coromandel. Muſ. D. Banks.

*tridentata.* 59. M. glabra, teſtacea, thorace rufo, clypeo tridentato.

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Inſula Guadeloupe. Muſ. D. Lund.
Statura et ſumma affinitas M. melanocephalae.
Caput rufum, baſi magis obſcurum, clypeo paruo, tridentato. Thorax punctatus, rufus, lateribus pallidioribus. Elytra et corpus teſtacea.

60. M. nigricans, thorace pallido: punctis duobus nigris. Ent. ſyſt. 2. 165. 45. * *mixta.*
Habitat in Africa aequinoctiali. D. Lee.

61. M. obſcure picea, immaculata, pilis breuiſſimis, erectis, albis. Ent. ſyſt. 2. 166. 46. * *obſcura.*
Oliv. Inſ. 1. 5. 42. 53. tab. 6. fig. 64.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

62. M. glabra, rufeſcens, elytris teſtaceis, clypeo quinquedentato. Ent. ſyſt. 2. 166. 47. * *rufa.*
~~Oliv. Inſ. 1. 5. 4. 3. tab. 7. fig. 29.~~
~~Voet. Col. tab. 9. fig. 70.~~
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

63. M. ſupra glabra, viridis, thorace linea dorſali, elytris futura nigris. Ent. ſyſt. 2. 166. 48. * *feſtiua.*
Oliv. Inſ. 1. 5. 47. 62. tab. 5. fig. 48.
Habitat in noua Zelandia. Muſ. D. Banks.

64. M. ſupra glabra, aurea, thoracis linea dorſali ſcutelloque ſanguineis. Ent. ſyſt. 2. 166. 49. * *lacta.*
Oliv. Inſ. 1. 5. 48. 63. tab. 6. fig. 56.
Habitat in noua Zelandia. Muſ. D. Banks.

65. M. ſupra viridis, holoſericea, ſubtus cuprea, elytris lineis eleuatis quatuor. Ent. ſyſt. 2. 167. 50. * *holoſericea.*
Habitat in Ruſſia auſtraliori. Muſ. D. Boeber.

66. M. viridi-aenea, elytris ſubſtriatis, obſcuris. Ent. ſyſt. 2. 167. 51. * Voet. Col. tab. 7. fig. 58. *Iulii.*
*Melolontha dubia.* Herbſt. Col. 3. tab. 25. f. 9. ~~XIV. 7~~

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 2. 210. 5.
Habitat in Germania. D. Smidt.

*nigrita.* 67. M. glabra, nigra, clypeo reflexo, elytris obſolete ſtriatis. Ent. ſyſt. 2. 167. 52. *
Habitat in America.

*Friſchii.* 68. M. nigro-aenea, nitida, elytris teſtaceis. Ent ſyſt. 2. 167. 53. *
*Scarabaeus vitis.* Friſh. Inſ. 4. 29. tab. 14.
Oliv. Inſ. 1. 5. 35. tab. a. fig. 29.
Degeer Inſ. 4 tab. 10. fig. 16.
Schaeff. Icon. tab. 23. fig. 3. 7.
Habitat in vite, roſa, Germaniae.

*vitis.* 69. M. viridis, thoracis lateribus flauis. Ent. ſyſt. 2. 167. 54.
Oliv. Inſ. 1. 5. 34. 39. tab. 2. fig. 12.
Schaeff. Icon. tab. 259. fig. 6.
Petagn. Inſ. Calabr. 5. 19.
~~Voet. Scarab. tab. 6. fig. 56. 57.~~
~~Sulz. Hiſt. Inſ. tab. 1. fig. 11.~~
Habitat in Europae et Americae vite, roſa.

*aurata.* 70. M. viridis, nitida, elytrorum margine tenuiſſime purpuraſcente, clypeo ſubemarginato.
Habitat in Carneolia. D. de Meyerle.
Maior M. Friſhii, at affinis. Caput et thorax laeuia, viridia, nitida.

*cardui.* 71. M. glabra, viridi-aenea, abdomine rufo, cinereo-villoſo. Ent. ſyſt. 2. 167. 55. *
*Melolontha maura.* Oliv. Inſ. 1. 5. 38. 45. tab. 8. fig. 90.
*Scarabaeus maurus.* Linn. ſyſt. Nat. 2. 548. 304.
Habitat in Carduo pycnocephalo Africae. Dom. Vahl.

72. M.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

72. M. villoſa, viridis, elytris ſerratis acuminatisque, femoribus poſticis incraſſatis. Ent. ſyſt. 2. 168. 56. * *ſerratulae.*

Habitat in Barbaria. Muſ. D. Desfontaines.

73. M. viridi-aenea, nitida, elytris vittis duabus aureis. Ent. ſyſt. 2. 168. 57. * *faſciatoſa.*

*Melolontha Leei.* Oliv. Inſ. 1. 5. 30. tab. 8. fig. 87.

*Scarabaeus Leei.* Sweder. Act. Holm. 1787. 188. 4.

Habitat in India orientali. D. Lee.

74. M. teſtacea, thorace maculis duabus nigris, pedibus ruſis. Ent. ſyſt. 2. 169. 58. * *errans.*

Oliv. Inſ. 1. 5. 45. 58. tab. 8. fig. 92.

Habitat in Anglia. Muſ. D. Tunſtall.

75. M. thorace aeneo-nitido: margine pallido, elytris teſtaceis, nigro-punctatis. *varians.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Parua. Caput viridi-aeneum, clypeo ſubreflexo rufeſcente. Thorax viridi-aeneus, nitidus, margine pallido, interdum puncto viridi. Elytra ſtriata, obſcure teſtacea: punctis plurimis nigris, ſparſis. Corpus teſtaceum.

*Variat* elytris nigro-variis.

76. M. ſubteſtacea, griſeo-villoſa, pedibus teſtaceis. Ent. ſyſt. 2. 169. 59. * *umbroſa.*

Habitat in Guinea. D. Iſert.

77. M. villoſa, vireſcens, capite nigro. Ent. ſyſt. 2. 169. 60. *rupicola.*

Oliv. Inſ. 1. 5. 68. 92. tab. 6. fig. 70.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

78. M. glabra, nigra, thoracis margine, abdomine femoribusque teſtaceis. Ent. ſyſt. 2. 169. 61. * *innuba.*

*Melo-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Melolontha innuba.* Oliv. Inſ. 1. 5. 46. 59. tab. 8. fig. 93.
Habitat in India. Muſ. Dr. Hunter.

*elongata.* 79. M. capite thoraceque nigris, elytris nitidulis, corpore teſtaceo. Ent. ſyſt. 2. 170. 62. *
Habitat - - D. Lee.

*rauca.* 80. M. clypeo reflexo, thorace elytrisque punctatis nigro-aeneis. Ent. ſyſt. 2. 170. 63.
Oliv. Inſ. 1. 5. 41. 56. tab. 6. fig. 62.
Habitat in Coromandel. Muſ. D. Banks.

*ferruginea* 81. M. rufa, thorace elytrisque vage punctatis. Ent. ſyſt. 2. 170. 64. *
Oliv. Inſ. 1. 5. 44. 56. tab. 7. fig. 82.
Habitat in Coromandel. Muſ. D. Banks.

*moerens.* 82. M. rufa, elytris teſtaceis, pectore nigro. Ent. ſyſt. ſuppl. 132. 64. *
Habitat in America boreali. D. v. Rohr.

*ſplendida.* 83. M. atra, elytris vitta abbreuiata, aurea. Ent. ſyſt. 2. 170. 65. *
Oliv. Inſ. 1. 5. 49. 65. tab. 5. fig. 50.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*aulicola.* 84. M. cuprea, nitida, clypeo emarginato. Ent. ſyſt. 2. 170. 66. *
Oliv. Inſ. 1. 5. 49. tab. 7. fig. 81.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

*lucicola.* 85. M. glabra, nigra, thoracis margine elytrisque teſtaceis. Ent. ſyſt. ſuppl. 132. 66. *
Habitat in America boreali. D. v. Rohr.

*atrata.* 86. M. glabra, atra, elytris ſtriatis. Ent. ſyſt. ſuppl. 132. 66. *
Habitat in America boreali. D. v. Rohr.

*arboricola* 87. M. nigra, hirta, elytris ſtriatis, teſtaceis. Ent. ſyſt. 2. 171. 67. *

*Melo-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Melolontha nitidula.* Oliv. Inſ. 1. 5. 46. 60. tab. 9. fig. 102.
Habitat in America.

8. M. capite thoraceque coeruleo-piloſis, elytris teſtaceis, pedibus nigris. Ent. ſyſt. 2. 171. 68. *horticola*
*Scarabaeus horticola.* Linn. ſyſt. Nat. 2. 554.
Fn. Sv. 391.
Geoff. Inſ. 1. 75. 8.
Oliv. Inſ. 1. 5. 26. tab. 2. fig. 17.
Schaeff. Icon. tab. 32. fig. 4.
Degeer Inſ. 4. tab. 10. fig. 18.
Payk. Fn. Sv. 2. 211. 6.
Herbſt. Col. tab. 25. fig. 1.
Habitat in Europae hortis.
*Larua* deuaſtat radices Braſſicae Botrydis et capitatae, intactis viridis et ſavoyae, *imago* Pomonam intacto Pyro communi.

9. M. thorace punctato-viridi: margine cupreo, elytris ſtriatis, teſtaceis. *nitidula.*
Habitat in America boreali. Muſ. D. Boſc.
Statura et magnitudo omnino M. horticolae. Caput nigrum, clypeo parum reflexo. Thorax punctatus, viridis, nitidulus, margine exteriori et poſtico cupreis. Scutellum nigrum, viridi nitidulum. Elytra crenato-ſtriata, teſtacea. Corpus nigrum.

10. M. capite thoraceque coeruleo-piloſis, elytris teſtaceis, abdomine villoſo, albo. Ent. ſyſt. 2. 171. 69. * *abdomina-lis.*
Habitat in Italia. Dr. Allioni.

11. M. cyanea, nigra, glabra, elytris rufis, abdomine punctis faſciculatis, albis, clypeo reflexo. Ent. ſyſt. 2. 171. 70. * *floricola.*

Melo-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

~~*Melolontha auſtriaca.* Herbſt Arch. tab. 20. f. 8.~~
Habitat in Africa. D. Prof. Vahl.

*aruicola.* 92. M. hirta, capitis clypeo reflexo. thorace coeruleſcente, corpore nigro. Ent. ſyſt. 2. 172. 71. *
Oliv. Inſ. 1. 5. 6. 4. tab. 7. fig. 84.
Habitat in Ruſſiae regionibus auſtralioribus.

*praticola.* 93. M. thorace ſubpiloſo, nigro flauoque vario, elytris teſtaceis. Ent. ſyſt. 2. 172. 72. *
Oliv. Inſ. 1. 5. 60. 83. tab. 7. fig. 74.
Habitat in Sibiria. Muſ. D. Banks.

*fruticola.* 94. M. capite thoraceque coeruleo-piloſis, elytris liuidis, clypeo apice reflexo. Ent. ſyſt. 2. 172. 73.
~~Oliv. Inſ. 1. 5. 36. 56. tab. 2. fig. 13.~~
Petagn. Inſ. Calabr. tab. 1. fig. 2.
*IV. 48* Schaeff. Icon. tab. 63. fig. 1.
Herbſt. Arch. 15. 11. tab. 19. fig. 24.
~~16. 12. tab. 19. fig. 26.~~
Voet. Col. tab. 8. fig. 67.
Habitat in Europae ſpicis ſecalinis.
*Femina* macula quadrata coleoptrorum, nigra, ſcutellum cingente.

*agricola.* 95. M. thorace villoſo, elytris liuidis: limbo faſciaque nigris, clypeo apice reflexo. Ent. ſyſt. 2. 173. 74. *
*IV. 48* *Scarabaeus agricola.* Linn. ſyſt. Nat. 2. 553. 58.
Oliv. Inſ. 1. 5. 61. 84. tab. 2. fig. 19.
Habitat in Germaniae Graminibus.

*lineata.* 96. M. villoſa, capite thoraceque obſcure aeneis, elytris teſtaceis: marginibus lineolisque duabus nigris. Ent. ſyſt. ſuppl. 133. 75. *
Habitat in Africae borealis arboribus. D. Schousboe.

*nuricola.* 97. M. atra, ſericea, elytris rufis: margine omni ~~rufo.~~ *nigro* Ent. ſyſt. 2. 173. 75.

Geoff.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 80. 15.
Oliv. Inf. 1. 5. 52. 71. tab. 3. fig. 25.
Panz. Naturf 24. tab. 1. fig. 10.
Herbſt. Arch. tab. 43. fig. 7.
Habitat in Europae graminoſis. D. Lee.

98. M. albo-farinoſa, thorace canaliculato atro, elytris faſcis, abdomine albo: punctis lateralibus atris. Ent. ſyſt. 2. 173. 76. * *atomaria.*
Oliv. Inf. 1. 5. 71. 98. tab. 5. fig. 52.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

99. M. ſupra coeruleo-, ſubtus argenteo-ſquamoſa, nitens, clypeo integro. Ent. ſyſt. 2. 173. 77. * *farinoſa.*
~~*Scarabaeus farinoſus.* Linn. ſyſt. Nat. 2. 555. 64. Fn. Sv. 399.~~
Geoff. Inf. 1. 79. 13.
Drury Inf. 2. tab. 32. fig. 4. ~~III. 48 IV. 9~~
Oliv. Inf. 1. 5. 66. 90. tab. 2. fig. 14. a. c.
Panz. Fn. Germ. 28. tab. 16.
Voet. Col. tab. 9. fig. 73.
Degeer Inf. 4. tab. 10. fig. 23.
Habitat in Europa auſtraliori.

100. M. ſupra viridi-ſquamoſa, ſubtus pallido-nitens, capite fuſco. Ent. ſyſt. 2. 174. 78. *ſquamoſa.*
~~*Melolontha minuta.* Panz. Naturf. 24. 9. 11. tab. 1. fig. 11.~~
*Melolontha argentea.* Laichart. 1. 42. 7.
Oliv. Inf. 1. 5. 65. 89. tab. 2. fig. 14. b. ~~III. 48 IV. 10~~
Herbſt. Arch. tab. 20. fig. 6.
Degeer Inf. 4. tab. 10. fig. 23.
Panz. Fn. Germ. 28. tab. 17.
Voet. Col. tab. 9. fig. 71. 72.
Payk. Fn. Sv. 2. 212. 7.
Habitat in Germania. Dom. Hybner.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bilineata.* 101. M. ſupra viridi-ſquamoſa, capite thoracis lineis duabus pedibusque fuſcis.

Habitat in Tanger. Muſ. D. de Seheſtedt.

Statura et ſumma affinitas M. ſquamoſae. Caput fuſcum clypeo reflexo. Thorax vireſcens lineis duabus latis fuſcis. Elytra viridia. Corpus ſubtus viridi-ſquamoſum, pedibus apice nigricantibus.

In hac et praecedentibus ſpeciebus articulus vltimus palporum anteriorum magis incraſſatus, at vix tamen proprii generis.

*regia.* 102. M. ſquamoſa, ſupra lutea, capite nigro, ſubtus cinerea. Ent. ſyſt. 2. 174. 79. *

*Scarabaeus aulicus.* Linn. ſyſt. Nat. 2. 555. 65. Oliv. Inſ. 1. 5. 44. 88. tab. 9. fig. 106.

Habitat in Africa. D. Prof. Vahl.

*glabrata.* 103. M. atra, nitida, elytris ſtriatis, flauo-maculatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paullo maior, quam M. argentea. Caput et thorax laeuia, atra, immaculata. Elytra abbreuiata, atra, flauo-maculata. Pedes ſimplices, picei.

*videns.* 104. M. atra, nitida, elytris ſtriatis teſtaceis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Affinis praecedenti, at paullo minor, et elytra minus ſtriata, teſtacea, ſutura nigra.

*argentea.* 105. M. clypeo marginato, nigra, ſubtus argenteo-nitens, elytris teſtaceis. Ent. ſyſt. 2. 174. 80. *

*Scarabaeus argenteus.* Scopol. Carn. 9.

Oliv. Inſ. 5. 67. 91. tab. 8. fig. 82.

Panz. Fn. Germ. 28. tab. 19.

Voet.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Voet. ſcarab. tab. 8. fig. 68.
Habitat in Angliae floribus frequens.

106. M. argenteo-ſquamoſa, clypeo reflexo. Ent. ſyſt. 2. 175. 81. * *graminicola.*
*Melolontha farinoſa.* Herbſt. Arch. tab. 25. f. 3.
Panz. Fn. Germ. 28. tab. 19.
Habitat in Germania. D. Prof. Helwig.

107. M. glabra, atra, elytris baſi maculaque media pallidis. Ent. ſyſt. ſuppl. 133. 81. * *pygmaea.*
Habitat in Carolina. Muſ. D. Boſc.

108. M. ſubrotunda, pubeſcens, obſcure ferruginea, antennis pedibusque teſtaceis. Ent. ſyſt. 2. 175. 82. *chryſomelina.*
*Scarabaeus chryſomeloides.* Schranck. Ent. 16. 25.
Panz. Fn. Germ. 34. tab. 11.
Habitat in Auſtria. D. Prof. Helwig.

109. M. ſupra nigro-, ſubtus albo-hirta, clypeo bifido. *capicola.*
Habitat ad Cap. Bon. Spei. D. Daldorff.
Statura et magnitudo omnino M. argenteae. Capitis clypeus porrectus, bifidus. Corpus ſupra nigrum, pilis nigris, ſubtus pilis albis denſiſſimis hirtum.

110. M. nigro-aenea, nitida, elytris brunneis: margine nigro. Ent. ſyſt. 2. 175. 83. * *cincta.*
*Melolontha marginata.* Oliv. Inſ. 1. 5. 70. 96. tab. 3. fig. 23.
Habitat in Inſula Guadeloupe. Muſ. D. Olivier.

111. M. clypeo porrecto ſubreflexo, hirta, nigra, elytris teſtaceis: margine nigro. Ent. ſyſt. 2. 175. 84. * *proboſcidea.*
Oliv. Inſ. 1. 3. 59. 81. tab. 8. fig. 96.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in India, in Africae floribus. D. Prof. Vahl.

*ſpinipes.* 112. M. nigra, immaculata, pedibus poſticis elongatis, femoribus baſi vniſpinoſis. Ent. ſyſt. 2. 176. 85. *

Oliv. Inſ. 1. 5. 72. 100. tab. 6. fig. 63.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*dentipes.* 113. M. nigra, elytris teſtaceis, clypeo quadridentato, femoribus tibiisque poſticis ſpinoſis. Ent. ſyſt. 2. 176. 86. *

Oliv. Inſ. 1. 5. 73. 102. tab. 6. fig. 66.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*podagrica.* 114. M. nigra, clypeo tridentato, femoribus tibiisque poſticis dentatis. Ent. ſyſt. 2. 176. 87. *

Oliv. Inſ. 1. 5. 75. 104. tab. 5. fig. 51.

Habitat in Coromandel. Muſ. D. Banks.

*arthritica.* 115. M. nigra, elytris griſeis, clypeo tridentato, femoribus tibiisque poſticis incraſſatis, ſubinermibus. Ent. ſyſt. 2. 176. 88. *

Oliv. Inſ. 1. 5. 75. 104. tab. 5. fig. 44.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*gonagra.* 116. M. griſea, pedibus rufis: femoribus poſticis incraſſatis muticis. Ent. ſyſt. 2. 177. 89. *

Oliv. Inſ. 1. 5. 64. 103. tab. 6. fig. 68.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*craſſipes.* 117. M. atra, albo-maculata, pedibus elongatis craſſiſſimis. Ent. ſyſt. 2. 177. 90. *

Oliv. Inſ. 1. 5. 72. 99. tab. 2. fig. 15.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*calcarata.* 118. M. nigra, elytris flauis: faſcia atra, femoribus tibiisque poſticis ſpinoſis rufis. Ent. ſyſt. ſuppl. 133. 89. *

Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

119. M.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

119. M. atra, albo-irrorata, femoribus poſticis elongatis craſſiſſimis, apice ſpinoſis, tibiis vnidentatis. Ent. ſyſt. 2. 177. 91. * *cancroides.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

120. M. thorace nigro: margine albo, elytris teſtaceis. Ent. ſyſt. 2. 178. 92. * *marginella.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

121. M. corpore polline vireſcenti-argenteo, elytris pedibusque teſtaceis. Ent. ſyſt. 2. 178. 93. * *puluerulenta.*
Oliv. Inſ. 1. 5. 69. 49. tab. 7. fig. 78.
Panz. Fn. Germ. 28. tab. 20.
Herbſt. Arch. tab. 25. fig. 7.
Sulz. Hiſt. Inſ. tab. 1. fig. 7.
Habitat in Germania.

122. M. hirta, atra, clypeo reflexo. Ent. ſyſt. 2. 178. 94. * *Morio.*
Habitat in Barbaria. Muſ. D. Desfontaines.

123. M. clypeo reflexo integro, corpore glabro nigro. Ent. ſyſt. 2. 178. 95. * *ſyluicola.*
Habitat in noua Hollandia. Muſ. D. Banks.

124. M. flaueſcens, pedibus rufis, thorace ſubſpinoſo. Ent. ſyſt. 2. 178. 96. * *ſubſpinoſa.*
*Melolontha elongata.* Herbſt. Arch. tab. 26. f. 3.
Oliv. Inſ. 1. 5. 70. 97. tab. 7. fig. 73.
Habitat in Americae meridionalis Inſulis.

125. M. capite thoraceque atris, elytris villoſis pedibusque teſtaceis, abdomine breui retuſo. Ent. ſyſt. 2. 179. 97. * *longipes.*
Oliv. Inſ. 1. 5. 76. 106. tab. 7. fig. 72.
Habitat - - Muſ. D. Banks.

126. M. villoſa, nigra, clypeo tridentato, elytris abbreuiatis teſtaceis. Ent. ſyſt. 2. 179. 98. * *abbreuiata*

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 1. 5. 75. 105. tab. 5. fig. 54.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*minuta.* 127. M. nigra, elytris griſeis, pedibus teſtaceis, femoribus poſticis acute dentatis. Ent. ſyſt. 2. 179. 99. *

Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

*mutabilis.* 128. M. atra, tomento cinereo villoſa. Ent. ſyſt. 2. 179. 100. *

Oliv. Inſ. 1. 5. 61. 69. tab. 3. fig. 24.
Voet. Col. tab. 9. fig. 75.
Habitat Tranquebariae. Dr. Koenig.

*variabilis.* 129. M. nigricans, elytris cinereo-nitentibus, antennis pedibusque teſtaceis. Ent. ſyſt. 2. 180. 101. *

Geoff. Inſ. 1. 74. 25.
Laichart. 1. 31. 4.
Fyesl. Heluet. 35.
Harrer tab. 21. fig. 24.
Oliv. Inſ. 1. 5. 52. 70. tab. 21. fig. 37.
Habitat in Germaniae et Americae ligno putreſcente.
*Varietas* Americana tomento aureo tuta.

*verſicolor.* 130. M. obſcure aenea, elytris tomento cinereo nitidulis. Ent. ſyſt. 2. 180. 102. *

Oliv. Inſ. 1. 5. 51. 68. tab. 9. fig. 105.
Habitat in Sierra Leon Africae. Muſ. D. Banks.

*ſplendidula* 131. M. viridi-aenea, rufo-micans, elytris ſtriatis obſcurioribus: baſi macula ferruginea.

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Statura et magnitudo M. variabilis. Antennae nigrae. Caput et thorax laeuia, viridia, rufo parum micantia. Elytra ſtriata, obſcuriora, magis

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

gis rufo-micantia maculaque diſtincta. baſeos rufa. Corpus obſcurum, abdomine ſubtus rufo.

132. M. rufa, elytris obſcuris auro-cyaneoque micantibus. *micans.*

Habitat in Amboina. D. Billardiere.

Statura omnino praecedentium. Caput et thorax rufa, immaculata. Elytra ſtriata, cyaneo auroque micantia. Corpus ferrugineum.

133. M. glabra, ferruginea, elytris ſtriatis. Ent. ſyſt. 2. 180. 103. *picea.*

Oliv. Inſ. 1. 5. 47. 61. tab. 5. fig. 43.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

134. M. gibba, nigra, thoracis macula laterali elytrorumque diſco ferrugineo. *diſcoidea.*

Habitat in Guinea. D. Meier. Muſ. D. de Seheſtedt.

Statura et magnitudo M. piceae. Caput nigrum, minime nitidum, antennis ferrugineis. Thorax niger, macula magna, laterali, ferruginea: puncto paruo, nigro. Elytra vix ſtriata, nigra, diſco vel potius vitta lata ferruginea, quae apicem haud attingit. Corpus gibbum, nigrum.

135. M. clypeo ſubreflexo, flaueſcens, elytris ferrugineis, fuſco-ſubfaſciatis. Ent. ſyſt. 2. 181. 104. * *araneoides*

Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

136. M. obſcure vireſcens, elytris lineis duabus rufis. Ent. ſyſt. 2. 181. 105. * *4lineata.*

Oliv. Inſ. 1. 5. 49. 66. tab. 6. fig. 68.

Habitat in Sierra Leon Africae. Muſ. D. Banks.

137. M. gibba, teſtacea, tomento cineraſcente, nitidula. Ent. ſyſt. 2. 181. 106. * *gibba.*

Oliv. Inſ. 1. 5. 50. 67. tab. 6. fig. 67.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

137. M.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*montieola.* 138. M. clypeo integro, reflexo, glabra, teſtacea capite atro. Ent. ſyſt. 2. 181. 107. *
Habitat in noua Hollandia. Muſ. D. Banks.

*humeralis.* 139. M. atra, elytris baſi punctoque medio pallidis Ent. ſyſt. 2. 181. 108.
Oliv. Inſ. 1. 5. 53. 72. tab. 3. fig. 26.
Panz. Fn. Germ. 34. tab. 10.
Habitat in Europa. Muſ. D. Tunſtall.

*vrſus.* 140. M. hirſutiſſima, atra, pedibus quatuor anticis teſtaceis. Ent. ſyſt. 2. 182. 9. *
Oliv. Inſ. 1. 5. 58. 79. tab. 8. fig. 88.
Wulf. Inſect. Capenſ. tab. 2. fig. 19.
Voet. Col. tab. 9. fig. 74.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.
*Variat* pedibus omnibus nigris.

*bombyliformis.* 141. M. hirſutiſſima, atra, elytris teſtaceis.
*Melolontha bombyliformis.* Pall. Icon. tab. A. fig. 17.
Habitat in Sibiria. Muſ. D. Lund.
Statura omnino et ſumma affinitas M. vrſus. Differt tamen elytris glabris, teſtaceis. Pedes omnes nigri.

*Lynx.* 142. M. hirta, nigra, elytrorum margine aureo. Ent. ſyſt. 2. 182. 110. *
Oliv. Inſ. 1. 5. 58. 80. tab. 9. fig. 101.
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

*crinita.* 143. M. hirta, ſupra viridis, ſubtus nigra. Ent. ſyſt. 2. 182. 111.
*Scarabaeus longipes* Linn. ſyſt. Nat. 2. 555. 66. Muſ. Lud. Vlr. 20.
Oliv. Inſ. 1. 5. 57. tab. 2. fig. 16.
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

*cyanipennis.* 144. M. cinereo-hirta, elytris cyaneis.
Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

Statu-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura omninó praecedentium, at diſtincta videtur. Caput et thorax cinereo-hirta. Elytra glabra, cyanea.

145. M. hirta, capite thoraceque viridibus, elytris fuſcis. Ent. ſyſt. 2. 183. 112. * *hirta.*
*Scarabaeus Alopecius.* Pall. Icon. tab. A. f. 15.
Oliv. Inſ. 1. 5. 56. 76. tab. 7. fig. 77.
Habitat in Sibiriae Leontodon Taraxaco.

146. M. aurea, fuluo-hirta, abdomine ferrugineo. Ent. ſyſt. 2. 183. 113. * *vulpes.*
Oliv. Inſ. 1. 5. 55. tab. 7. fig. 76.
Habitat in Sibiriae Leotodon Taraxaeo.
*Varietas* praecedentis obſeruante D. Boeber.

147. M. fulua, hirta, elytris abbreuiatis atris, abdomine ferrugineo. Ent. ſyſt. 2. 183. 114. * *Melis.*
Habitat in Barbaria. Muſ. D. Desfontaines.

148. M. piloſa, nigra, cinereo-hirta, elytris teſtaceis: lineis tribus apicis albidis. Ent. ſyſt. 2. 184. 115. * *Bombylius.*
Habitat in Africae plantis. Muſ. D. Vahl.

149. M. piloſa, cyanea, elytris teſtaceis: lineis tribus albicantibus. Ent. ſyſt. 2. 184. 116. * *vittata.*
*Scarabaeus acuminatus.* Lepech. Itin. 1. 508. tab. fig. 9.
*Scarabaeus oxypterus.* Pall. Itin. 1. 462. 26.
Oliv. Inſ. 1. 5. 55. 74. tab. 8. fig. 94.
Pall. Icon. 1. 14. tab. A. fig. 14.
Degeer. Inſ. 7. 644. 40. tab. 48. fig. 8.
Petiv. Gazoph. tab. 2. fig. 12.
Voet. Scarab. tab. 5. fig. 35.
Habitat in India.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

128. BVPRESTIS. *Palpi* filiformes: articulo vltimo obtuſo, truncato.
*Maxilla* obtuſa, vnidentata.
*Ligula* cylindrica, acuminata.
*Antennae* ſerratae.

*vnidentata* 1. B. elytris vnidentatis aeneis: margine aureo. Ent. ſyſt. 2. 185. 1. *
*Bupreſtis vnidentata.* Oliv. Inſ. 2. 32. tab. 8. fig. 86.
Habitat - - Muſ. D. Banks.

*bicolor. barbara* 2. B. elytris acuminatis viridi-aeneis: macula flaua, pectore abdomineque flauis. Ent. ſyſt. 2. 185. 2. *
*Bupreſtis bicolor.* Oliv. Inſ. 2. 32.
Habitat in America. D. Drury.

*4dentata.* 3. B. elytris acuminatis obſcuris, corpore aeneo-nitido, ano quadridentato. Ent. ſyſt. 2. 186. 3.*
Habitat - - Muſ. D. Banks.

*4vittata.* 4. B. elytris acuminatis punctato-ſtriatis obſcuris: vittis duabus flauis.
Habitat ad Cap. Bon. Spei. Muſ. D. de Seheſtedt.
Parua. Caput nigrum, punctis quatuor paruis obſcure

---

Bupreſtis corpus oblongum, depreſſiuſculum, rarius cylindricum, immarginatum, tardum; *capite* rotundato, inſerto, obtuſo; *oculis* magnis, oblongis, vix prominulis, marginalibus; *antennis* ſerratis, approximatis, thorace breuioribus, ſubter oculos inſertis; *thorace* transuerſo, ſaepius ſinuato; *elytris* rigidis, fornicatis: margine ſaepius dentato, longitudine abdominis; *ſcutello* paruo, rotundato; *pedibus* breuibus, validis; *tarſis* quinquearticulatis; *colore* varjo.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

obſcure flaueſcentibus. Thorax punctatus: punctis aliquot lineolaque marginali baſeos obſcure flauis. Elytra ſtriata: ſtriis punctatis obſcura, vittis duabus tenuibus, apicem haud attingentibus, flauis, apice valde acuminata. Corpus cyaneum, abdominis margine parum flauicante.

. B. elytris acuminatis, nigra, glabra, ano bidentato. Ent. ſyſt. ſuppl. 134. 3. * *regalis.* *nigrita*
Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

. B. elytris bidentatis rugoſis, thorace laeui, corpore inaurato. Ent. ſyſt. 2. 186. 4. * *gigantea.*
Linn. ſyſt. Nat. 2. 659. 1. Muſ. Lud. Vlr. 85.
Oliv. Inſ. 2. 32. tab. 1. fig. 1.
Degeer. Inſ. 4. 134. 1.
Grew. Muſ. 165. tab. 13.
Sloan. Iam. 2. tab. 236. fig. 1. 2.
Petiv. Gazoph. 147. fig. 16. 17.
Sulz. Inſ. tab. 6. fig. 38.
Menon Surin. tab. 30. fig. inf.
Seb. Muſ. 3. tab. 48. fig. 12.
Herbſt. Arch. tab. 28. fig. 1.
Habitat in America et Aſia.
*Varietas* in Muſ. D. Lund. thorace nigro-bimaculato et elytris ſerratis apice bidentatis, vix diſtincta.

7. B. elytris bidentatis punctatis: lineis quatuor eleuatis viridi-aeneis vittaque lata aurea. Ent. ſyſt. 2. 186. 5. * *vittata.*
*Bupreſtis vittata.* Oliv. Inſ. 2. 32. tab. 3. f. 17.
Sulz. Hiſt. Inſ. tab. 6. fig. 14.
Habitat in India. Muſ. D. Banks.

8. B. elytris bidentatis, punctato-ſtriatis viridibus: dorſo aureo. Ent. ſyſt. 2. 187. 6. *faſtuoſa.*

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 2. 32. tab. 8. fig. 81.
Habitat in America ſeptentrionali. Muſ. D. Banks

*regalis.* 9. B. elytris bidentatis fuſcis: faſciis duabus fl
xuoſis lineolisque duabus remotis viridibu
Ent. ſyſt. 2. 187. 7. *
Habitat in Cayenna. Muſ. D. Geoffroy.

*collaris.* 10. B. elytris bidentatis viridi-aureis, thorace fuſc
nitido: ſtrigis duabus aureis, ſterno cupre
Ent. ſyſt. 2. 187. 8. *
Oliv. Inſ. 2. 32. tab. 2. fig. 9.
Habitat in Cayenna. Muſ. D. Olivier.

*attenuata.* 11. B. elytris bidentatis attenuatis ſtriatis, viridi
aenea: ſutura cuprea. Ent. ſyſt. 2. 187. 9.
Habitat in Rio Faneiro. Muſ. D. Banks.

*punctatiſſima.* 12. B. elytris bidentatis cupreis: punctis numeroſi
violaceis. Ent. ſyſt. 2. 188. 10. *
*Bupreſtis punctatiſſima.* Oliv. Inſ. 2. 32. tab. 7
fig. 76.
Habitat in Sierra Leon Africae. Muſ. D. Banks.

*ruſipes.* 13. B. elytris ſtriatis cyaneis: maculis quatuor fla-
uis, corpore viridi-aeneo-nitido. Ent. ſyſt. 2.
188. 11. *
*Bupreſtis ruſipes.* Oliv. Inſ. 2. 32. tab. 7. f. 73.
Petiv. Gazoph. tab. 2. fig. 2.
Habitat in Marylandia. Muſ. Britann.

*Berolinenſis.* 14. B. elytris bidentatis, viridi nigroque variis, an
tridentato. Ent. ſyſt. 2. 188. 12. *
*Bupreſtis Berolinenſis.* Herbſt. Arch. tab. 28.
fig. 5.
1782 Schaeff. Icon. tab. 55. fig. 7.
Habitat in Germania.

*calcarata.* 15. B. elytris bidentatis aeneis: punctis elevatis
atris, tibiis intermediis dentatis.

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Auſtria. D. Scheidler.

Summa affinitas B. Berolinenſis, cuius forte varietas ſexus. Corpus ſupra punctatum aeneum, elytris punctis eleuatis, atris, ſcabris, ſubtus cupreum, tibiis intermediis, dente valido armatis.

16. B. elytris bidentatis ſtriatis, fuſca, ſubtus cuprea, albo-irrorata. *B. aenea Linn.* *carniolica.*

*Bupreſtis ſubrugoſa.* Payk. Fn. Sv. 2. 218. 5.

Habitat in Auſtria. D. de Meyerle.

Statura et magnitudo B. Berolinenſis. Caput et thorax punctata, obſcura, cupro parum nitidula. Elytra ſtriata, obſcura, apice bidentata. Corpus et pedes cuprea, nitida, punctis albis numeroſiſſimis irrorata.

17. B. elytris bidentatis, aenea, nitidiſſima, elytris apice abdomineque aureis. Ent. ſyſt. 2. 183. 13. * *corruſca.*

*Bupreſtis corruſca.* Oliv. Inſ. 2. 32. tab. 9. f. 99.

*Bupreſtis tereticollis.* Pall. Icon. tab. D. fig. 18.

Habitat in Iamaica. Muſ. D. Banks.

18. B. elytris bidentatis ſtriatis aeneis coeruleo-micantibus: margine aureo. Ent. ſyſt. 2. 189. 14. * *micans.*

Habitat in Italia, D. de Slanbuſch; Hafniae. Muſ. D. Lund.

19. B. elytris truncatis, bidentatis, viridibus: vitta coeruleſcente marginibusque aureis. Ent. ſyſt. 2. 189. 15. *decora.*

*Bupreſtis decora.* Oliv. Inſ. 2. 32. tab. 8. f. 82.

Habitat in America. Muſ. D. Banks.

20. B. elytris bidentatis, ſulcatis, viridibus, auro variegatis. *regia.*

*Bupre-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis equeſtris.* Oliv. Inſ. 2. 32. tab. 9. f. 10
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Statura omnino B. funeſtae. Caput punctatum v
ridi-aeneum. Thorax viridis, nitidus: punct
impreſſis aureis. Maculae praeterea tres baſe
impreſſae: intermedia maiore. Elytra bide
tata, ſulcata viridi auroque variegata. Corpu
aureum, tarſis cyaneis.

*funeſta.* 21. B. elytris bidentatis, obſcure aenea, thora
punctato, cupreo, nitido, elytris ſtriatis. En
ſyſt. 2. 189. 16. *
Habitat in Guinea. D. Iſert.

*varioloſa.* 22. B. elytris bidentatis, punctata aenea, nitida, tar
ſis cyaneis.
Habitat in Inſula Trinidad. Dr. Pflug. Muſ. D
de Seheſtedt.
Magnitudo et ſtatura B. funeſtae. Caput retuſum
viridi-aeneum, nitidum. Thorax punctatus
viridi-aeneus, nitidus. Elytra apice bidentata
viridi-aenea, nitida, punctis varioloſis, impreſ
ſis. Corpus aeneum, nitidum, tarſis cyaneis.

*umbroſa.* 23. B. elytris bidentatis, ſtriatis, nigro-aeneis
vitta marginali nitidiore. Ent. ſyſt. 2. 189
17. *
Habitat in Guinea. D. Iſert.

*lurida.* 24. B. elytris bidentatis, obſcura, ano tridentato.
Ent. ſyſt. 2. 190. 18.
Oliv. Inſ. 2. 32. tab. 8. fig. 83.
Habitat in America boreali. Muſ. D. Banks.

*obſcura.* 25. B. elytris bidentatis, obſcure aenea: punctis
eleuatis obſcurioribus. Ent. ſyſt. 2. 190. 19.
Habitat in America boreali. Muſ. D. Blackburn.

26. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. B. elytris bidentatis, aeneo-nigra, thorace punctato, elytris ſtriatis. Ent. ſyſt. ſuppl. 134. 20.* *melancholica.*

Habitat in India. Muſ. D. de Seheſtedt.

. B. elytris ſubbidentatis, ſtriatis, aenea, capite abdomineque teſtaceo-maculatis. Ent. ſyſt. 2. 190. 20. * *punctata.*

*Bupreſtis punctata.* Oliv. Inſ. 2. 32. tab. 10. fig. 114.

Habitat in Barbaria. D. Prof. Vahl.

. B. elytris bidentatis, ſtriatis, viridibus: maculis ſex teſtaceis. *12maculata.*

Habitat in Iaua. D. Billardiere.

Media. Caput punctatum, canaliculatum, viridi-aeneum, nitidum, antennis nigris. Thorax puncto vtrinque impreſſo, punctatus, nitidus. Elytra viridia, obſcuriora, ſtriata, apice bidentata, maculis ſex teſtaceis, 2 interiore maiore, antice lunata, exteriore marginali, 2 exteriore marginali, 1 transuerſa, flexuoſa, 1 marginali, lineari.

9. B. elytris bidentatis, ſtriatis, nigris: punctis numeroſis impreſſis albis. Ent. ſyſt. 2. 190. 21. * *morbilloſa.*

*Bupreſtis morbilloſa.* Oliv. Inſ. 2. 32. tab. 8. fig. 84.

Habitat in America meridionali. Muſ. D. Banks.

0. B. elytris bidentatis, viridibus: maculis numeroſis, aureis, impreſſis, pedibus cyaneis. Ent. ſyſt. 2. 191. 22. * *aurifera.*

*Bupreſtis aurifer.* Oliv. Inſ. 2. 32. tab. 9. f. 95.

Habitat in Cayenna. Muſ. D. Olivier.

11. B. elytris bidentatis, viridi-aeneis, apice cyaneis: faſciis duabus ferrugineis. Ent. ſyſt. 2. 191. 23. * *faſciata.*

*Bupre-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis faſciata.* Oliv. Inſ. 2. 32. tab. 9. f. 92.
Habitat in America boreali. Muſ. Dr. Hunter.

*lucidula.* 32. B. elytris bidentatis, aenea nitidiſſima: vitta abbreuiata abdomineque fulno aeneis. Ent. ſyſt. 2. 191. 24. *
Habitat Surinami. Muſ. D. Boſc.

*rauca.* 33. B. elytris bidentatis, ſubſtriatis, corpore obſcure aeneo, immaculato. Ent. ſyſt. 2. 191. 25. *
Habitat in Barbaria. D. Prof. Vahl.

*ſtriata.* 34. B. elytris truncatis, ſubbidentatis, cupreis: [illegible] neis eleuatis quatuor. Ent. ſyſt. 2. 191. 26.
*Bupreſtis ſtriata.* Oliv. Inſ. 2. 32. tab. 7. f. 77.
Habitat in Penſyluania. Muſ. D. Banks.

*rutilans.* 35. B. elytris tridentatis ſerratisque viridibus, nigro-maculatis: margine aureo. Ent. ſyſt. 2. 191. 27. *
*Bupreſtis rutilans.* Oliv. Inſ. 2. 32. tab. 5. f. 45.
IV. 11
*Bupreſtis faſtuoſa.* Jaqu. Miſcell. 2. tab. 23. f. 2.
II. 56.
Schaeff. Icon. tab. 35. fig. 6.
Pall. Icon. tab: D. fig. 12.
Habitat in Auſtria. Dr. Schulz.

*ignita.* 36. B. elytris tridentatis, thorace impreſſo, corpore inaurato. Ent. ſyſt. 192. 28. *
*Bupreſtis ignita.* Linn. ſyſt. Nat. 2. 669. 3.
Oliv. Inſ. 2. 32. tab. 4. fig. 33.
Sulz. Hiſt. Inſ. tab. 6. fig. 14.
Petiv Gazoph. tab. 26. fig 8.
Paempf. Iaponenſ. 1. tab. 10. fig. 7.
Habitat in India orientali.

*lineata.* 37. B. elytris truncatis, ſubtridentatis, nigro-aeneis vittis duabus fuluis. Ent. ſyſt. 2. 192. 29. *
*Bupreſtis lineata.* Oliv. Inſ. 2. 32. tab. 8. f. 80.
Habitat in America. Muſ. D. Banks.

38. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

38. B. viridi-nitens, elytris tridentatis: maculis duabus aureis ocellarique flaua. Ent. ſyſt. 2. 193. 30. * *ocellata.*

*Bupreſtis ocellata.* Oliv. Inſ. 2. 32. tab. 1. fig. 3.
Sulz. Hiſt. Inſ. tab. 6. fig. 15.
Degeer Inſ. 7. 633. 30. tab. 47. fig. 12.
Habitat in India orientali.

39. B. elytris tridentatis, corpore aeneo immaculato. Ent. ſyſt. 2. 193. 31. * *aenea.*

*Bupreſtis tricuſpidata.* Oliv. Inſ. 2. 32. tab. 8. fig. 87.
Habitat in Coromandel. Muſ. D. Banks.

40. B. elytris tridentatis cyaneis flauo-maculatis. Ent. ſyſt. 2. 193. 32. * *maculata.*

*Bupreſtis maculata.* Oliv. Inſ. 2. 32. t. 6. f. 61. t. 10. f. 61.
Habitat in Sibiria. Muſ. D. Banks.

41. B. elytris truncatis, dentatis, ſtriatis nigris: maculis quatuor flauis. Ent. ſyſt. 2. 193. 33. * *flauomaculata.*

*Bupreſtis flauomaculata.* Degeer Inſ. 4. 129. 1. 2.
Herbſt. Arch. tab. 28. fig. 8.
Payk. Fn. Sv. 2. 213. 1.
Habitat in Suecia. D. de Paykull.

42. B. elytris ſubtridentatis nigris: lineis tribus baſeos vnicaque apicis interrupta impreſſis albis. Ent. ſyſt. 2. 194. 34. * *interrupta*

*Bupreſtis interrupta.* Oliv. Inſ. 2. 32. tab. 4. fig. 28.
Habitat in Senegallia. Muſ. D. Olivier.

43. B. elytris ſerrato-tridentatis: punctis cinereis impreſſis, ſterno porrecto conico. Ent. ſyſt. 2. 194. 35. * *ſternicornis.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis ſternicornis.* Linn. ſyſt. Nat. 2. 660. 5. Muſ. Lud. Vlr. 88.
Oliv. Inſ. 2. 32. tab. 6. fig. 52.
Herbſt Arch. tab. 46. fig. 2.
Grew Muſ. 167. tab. 13.
Habitat in India orientali.
*Variat* punctis elytrorum conſolidatis, excepto puncto latiori baſeos.

*Chryſis.* 44. B. elytris ſerrato-tridentatis caſtaneis, ſterno conico porrecto. Ent. ſyſt. 2. 194. 36. *
*Bupreſtis ſternicornis.* Linn. ſyſt. Nat. 2. 660. 5. β. Muſ. Lud. Vlr. 88. β.
*Bupreſtis Chryſis.* Oliv. Inſ. 2. 32. tab. 2. f. 8.
Degeer Inſ. 4. 136. 2. tab. 17. fig. 25.
Herbſt. Arch. tab. 28. fig. 2.
Habitat in India orientali.

*caſtanea.* 45. B. elytris ſerrato-tridentatis, caſtaneis: puncto baſeos ferrugineo impreſſo, thorace atro, ferrugineo-reticulato. Ent. ſyſt. 2. 195. 37. *
*Bupreſtis caſtanea.* Oliv. Inſ. 2. 32. tab. 2. fig. 3. b.
Habitat in Senegallia. Muſ. D. Banks.

*violacea.* 46. B. elytris truncato-quadridentatis, violaceis, corpore elongato cyaneo. Ent. ſyſt. 2. 195. 38. *
Habitat in Sierra Leon Africae. Dr. Pflug.

*equeſtris.* 47. B. elytris ſerratis cyaneis: faſcia poſtica teſtacea. Ent. ſyſt. 2. 195. 39. *
Oliv. Inſ. 2. 32. tab. 10. fig. 109.
Habitat in Braſilia. Muſ. D. Lee.

*depreſſa.* 48. B. elytris ſerratis ſulcatis, corpore depreſſo obſcuro. Ent. ſyſt. 2. 195. 40. *
Habitat in Oriente. D. Prof. Forſkahl.

49. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

49. B. elytris ſerratis, longitudinaliter rugoſis: maculis duabus impreſſis, thorace ſulcato. Ent. ſyſt. 2. 195. 41. * *mariana.*

*Bupreſtis mariana.* Linn. ſyſt. Nat. 2. 660. 6.
Fn. Sv. 754. Muſ. Lud. Vlr. 89.
Oliv. Inſ. 2. 32. tab. 1. fig. 4.
Herbſt. Arch. tab. 28. fig. 4.
Drury Inſ. 1. tab. 30. fig. 3.
Schaeff. Icon. tab. 49. fig. 1.
tab. 128. fig. 3.
Payk. Fn. Sv. 2. 220. 7.
Habitat in Europa boreali.

50. B. elytris ſerratis viridibus: ſulco abbreuiato ſuturali impreſſa. Mant. Inſ. 1. 178. 28. * *farinoſa.*
Ent. ſyſt. 2. 196. 42. *
Habitat in noua Hollandia. Muſ. D. Banks.

51. B. elytris ſerratis, viridibus punctatis, thorace aeneo: ſulco maculisque aureis impreſſis. Ent. ſyſt. 2. 196. 43. * *ventricoſa.*

*Bupreſtis ventricoſa.* Oliv. Inſ. 2. 32. tab. 6. fig. 63.
Habitat in Madagaſcar. Muſ. D. Banks.

2. B. elytris ſerratis, viridi-aenea, thorace vtrinque macula impreſſa aurea. Ent. ſyſt. 2. 196. 44. * *ſmaragdula.*

*Bupreſtis ſmaragdula.* Oliv. Inſ. 2. 32. tab. 1. fig. 2.
Habitat in India orientali. Muſ. D. Olivier.

3. B. elytris ſerratis, aenea nitidiſſima, elytris apice aureis. Ent. ſyſt. 2. 196. 45. * *fulminans.*
Habitat in Indiis. Muſ. D. Zſchuck.

4. B. elytris ſerratis viridi-aenea: ſutura cyanea apice aureis. *ſuturalis.*
Habitat in India.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:24 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo omnino praecedentis, omnino diſtincta. Caput retuſum, aeneum Thorax aeneus, medio canaliculatus lateribus que depreſſis. Elytra punctato-ſtriata, viridi apice quinquedentata, aurea ſuturaque cyanea Corpus aeneum, nitidum, ano concolore.

*cyanipes.* 55. B. elytris ſerratis ſtriatis, viridi-aeneis, thorac plano coeruleo. Ent. ſyſt. 2. 196. 46.

*Buprestis cyanipes.* Oliv. Inſ. 2. 32. tab. 7. fig 104.

Habitat in America meridionali. Muſ. Dr. Hunter

*modeſta.* 56. B. elytris ſerratis, corpore obſcurioribus: maculis tribus impreſſis aureis. Ent. ſyſt. 2. 197 47. *

*Buprestis modeſta.* Oliv. Inſ. 2. 32. tab. 7. f. 72

Habitat in Braſilia. Muſ. D. Banks.

*blanda.* 57. B. elytris ſerratis, attenuatis, ſulcatis, viridi-aeneis, ſulcis cupreis. Ent. ſyſt. 2. 197. 48.

*Buprestis blanda.* Oliv. Inſ. 2. 32. tab. 9. f. 94.

Habitat in America meridionali. Muſ. Dr. Hunter.

*aurata.* 58. B. elytris ſerratis, aurea, thorace aeneo. Ent. ſyſt. 2. 197. 49. *

*Buprestis aurata.* Oliv. Inſ. 2. 32. tab. 9. f. 93.

Habitat in America. Muſ. Dr. Hunter.

*teſtacea.* 59. B. elongata, elytris ſerratis, ſtriatis teſtaceis punctis duobus fuſcis.

Habitat in America meridionali. D. Smidt. Muſ D. de Seheſtedt.

Statura valde elongata, magnitudine praeceden tis. Caput fuſcum, ſubretuſum. Thorax laeuis, teſtaceus, maculis duabus approximatis dorſalibus nigris. Elytra ſerrata, ſtriata, teſtacea, punctis duobus remotis, fuſcis. Pectus teſta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717 Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

teſtaceum. Abdomen atro-cyaneum, nitidum, ſegmentis abdominis teſtaceis. Pedes atri.

60. B. elytris ſerratis viridi-aenea, thorace vittis duabus, elytris vnica rufo-cupreis. Ent. ſyſt. 2. 197. 50. ♀ *fulgida*

*Bupreſtis fulgida.* Oliv. Inſ. 2. 32. tab. 11. fig. 124.

Habitat in India orientali. Muſ. D. Geoffroy.

61. B. elytris acute ſerratis, aenea, nitida, elytris vitta aurea. *cyaniceps.*

Habitat in Sumatra. D. Daldorff.

Maior. Antennae cyaneae, apice nigrae. Caput cyaneum, valde canaliculatum, canali viridi. Thorax punctatus, canaliculatus, viridi-aeneus. Elytra valde et acute ſerrata, punctato-ſubſtriata, viridi-aenea: vitta media lata, aurea. Corpus aeneum, ſuluo-puluerulentum.

62. B. elytris ſerratis, ſtriatis cupreis: punctis tribus impreſſis aureis. Ent. ſyſt. 2. 198. 51. ♀ *3punctata.*

*Bupreſtis 3punctata.* Oliv. Inſ. 2. 32. tab. 2. fig. 10.

Habitat in America meridionali. Muſ. Dr. Hunter.

63. B. elytris ſerratis, aenea, nitida, thorace lineis duabus elytrorumque vnica aureis. Ent. ſyſt. 2. 198. 52. ♀ *elegans.*

*Bupreſtis elegans.* Oliv. Inſ. 2. 32. tab. 8. f. 89.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

64. B. elytris ſerratis, nigro-aenea, pubeſcens, elytris faſciis tribus flauis: anteriore margine dilatata. Ent. ſyſt. 2. 198. 53. ♀ *trifaſciata*

Habitat in America meridionali.

65. B. elytris ſerratis, nigra, fronte puncto vnico, thorace duobus elytris tribus flauis. Ent. ſyſt. 2. 198. 54. ♀ *9maculata*

Bupre- *Vt. 17.*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis 9maculata.* Linn. ſyſt. Nat. 2. 662. 17.
Oliv. Inſ. 2. 32. tab. 4. fig. 30.
Panz. Naturf. 24. 33. 47. tab. 1. fig. 48.
Sulz. Hiſt. Inſ. tab. 6. fig. 17.
Habitat in Europa auſtrali. Muſ. D. Zſchuck.

*porcata.* 66. B. elytris ſerratis porcatis, thorace angulato, corpore fuſco nitido. Ent. ſyſt. 2. 199. 55. *
*Bupreſtis depreſſa.* Linn. Mant. 2. 583.
Oliv. Inſ. 2. 32. tab. 2. fig. 15.
Habitat in Oriente. D. Prof. Forſkahl.

*dorſata.* 68. B. elytris ſerratis, obſcure aenea, ano bidentato, femoribus anticis incraſſatis vnidentatis. Ent. ſyſt. 2. 199. 56. *
Habitat in America meridionali.
Femora antica valde incraſſata, acute vnidentata.

*chryſoſtigma.* 69. B. elytris ſerratis, longitudinaliter ſulcatis: maculis duabus aureis impreſſis, thorace punctato. Ent. ſyſt. 2. 199. 57. *
*Bupreſtis chryſoſtigma.* Linn. ſyſt. Nat. 2. 660. 7. Fn. Sv. 755.
Geoff. Inſ. 1. 125. 1.
Degeer Inſ. 4. 129. 3.
Oliv. Inſ. 2. 32. tab. 6. fig. 54.
Herbſt. Arch. tab. 28. fig. 6.
Schaeff. Elem. tab. 31. fig. 1. 2.
Sulz. Inſ. tab. 6. fig. 39.
Payk. Fn. Sv. 2. 221. 8.
Habitat in Europae nemoribus.

*impreſſa.* 70. B. elytris ſerratis, laeuibus, aeneis: punctis tribus impreſſis. Ent. ſyſt. 2. 200. 58.
*Bupreſtis impreſſa.* Oliv. Inſ. 2. 32. tab. 5. fig. 42.
Habitat in India. Muſ. D. Bernoulli.

71. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

71. B. elytris ſubſerratis, laeuibus, nigris: punctis tribus impreſſis aureis. Ent. ſyſt. 4. App. 449. 58. * *affinis.*

Habitat in Germania. D. Schneider.

72. B. elytris ſerratis nigris: maculis ſex viridibus, fronte impreſſa. Ent. ſyſt. 2. 200. 59. * *frontalis.*

*Bupreſtis frontalis.* Oliv. Inſ. 2. 32. tab. 5. f. 44.

Habitat in Carolina auſtraliori. Muſ. D. d'Orcy.

73. B. elytris ſerratis, obſcuris, albo-maculatis. Ent. ſyſt. 2. 200. 60. * *ornata.*

*Bupreſtis ornata.* Oliv. Inſ. 2. 32. tab. 7. f. 67.

Habitat in Penſyluania. Muſ. D. Banks.

74. B. elytris ſerratis, ſupra nigricans, thorace lineis tribus, elytris punctis quatuor impreſſis viridibus. *8punctata.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Magnitudo B. ſerratae. Caput canaliculatum, obſcurum. Thorax niger, lineis tribus impreſſis, viridibus. Elytra ſerrata, laeuia, nigra, punctis quatuor impreſſis, 1. 1. 2. Corpus aeneum, tarſis cyaneis.

75. B. elytris ſerratis, aenea, fronte tuberculo transuerſo eleuato, abdomine ſerrato. Ent. ſyſt. 2. 200. 61. * *ſerrata.*

Habitat in Guinea. D. Iſert.

76. B. elytris ſerratis, cinereo-villoſa, elytris nigris: faſciis duabus ferrugineis. Ent. ſyſt. 2. 201. 62. * *taeniata.*

*Bupreſtis blanda.* Oliv. Inſ. 2. 32. tab. 5. fig. 41. 49.

Roſſ. Fn. Etr. 1. 191. 474.

Pill. It. Poſ. Scl. 84. tab. 7. fig. 12.

Habitat in Hyperici floribus Europae auſtralioris.

77. B.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*elongata.* 77. B. elytris ſerratis, vireſcentibus, corpore elongato cupreo. Ent. ſyſt. 2. 201. 63. *
*Bupreſtis elongata.* Oliv. Inſ. 2. 32. tab. 9. fig. 102.
Habitat in Barbaria. Muſ. D. Olivier.

*pulcra.* 78. B. elytris ſerratis: faſciis tribus ſanguineis abbreuiatis, corpore atro. Ent. ſyſt. 2. 201. 64. *
Habitat in Marocco. Muſ. D. de Seheſtedt.

*gibboſa.* 79. B. elytris ſerratis, ſtriatis, baſi gibbis, viridi-aenea, thorace canaliculato. Ent. ſyſt. 2. 201. 65. *
*Bupreſtis gibboſa.* Oliv. Inſ. 2. 32. tab. 6. f. 59.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*nobilis.* 80. B. elytris ſerratis, viridi-aenea, capite thoraceque fuſco-maculatis, elytris atris: faſciis tribus lineolaque apicis viridibus. Ent. ſyſt. 2. 202. 66. *
*Bupreſtis nobilis.* Oliv. Inſ. 2. 32. tab. 5. f. 43.
Habitat in Cayenna. D. v. Rohr.

*cylindrica.* 81. B. elytris ſerratis, ſtriatis, thorace villoſo, corpore cylindrico nigro. Ent. ſyſt. 2. 202. 67. *
Habitat in Oriente. D. Prof. Forſkahl.

*voluulus.* 82. B. elytris ſerratis, flauo-maculatis, thorace inaequali, villoſo, corpore cylindrico, villoſo. Ent. ſyſt. 2. 202. 68. *
Habitat in Americae meridionalis Inſulis.

*tubulus.* 83. B. elytris ſerratis, crenato-ſtriatis, flauo-maculatis, thorace punctato, corpore cylindrico nigro.
Habitat in Carolina. Muſ. D. Boſc.
Statura omnino B. voluuli, at duplo minor. Caput et thorax punctata, nigra, immaculata. Elytra glabra, crenato-ſtriata, nigra: punctis

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ctis flauis 2. 2. 1. transuerſo, maiori 1. Corpus atrum.

84. B. elytris integris, fronte ſulcata, corpore ſcabro-aureo. Ent. ſyſt. 2. 202. 69. * *ſcabra.*

*Bupreſtis ſcabra.* Oliv. Inſ. 2. 32. tab. 3. f. 25.
Habitat in India. Dr. Mallet.

85. B. elytris integris, eleuato-ſtriatis fuſcis: punctis quinque flauis. Ent. ſyſt. 2. 203. 71. * *decoſtigma*

Panz. Fn. Germ. 29. tab. 18. 19.
Habitat in Auſtria. D. Saldoner.
*Variat* punctis adhuc duobus ad baſin elytrorum.

86. B. elytris integris: maculis quatuor albis, corpore coeruleo. Ent. ſyſt. 2. 203. 72. *8guttata.*

*Bupreſtis 8guttata.* Linn. ſyſt. Nat. 2. 659. 2.
Fn. Sv. 753.
Oliv. Inſ. 2. 32. tab. 4. fig. 36.
Geoff. Inſ 1. 126. 2.
Degeer Inſ. 4. 132. 5. tab. 4. fig. 20.
Herbſt Arch. tab. 28. fig. 7.
Schaeff. Icon. tab. 204. fig. 4.
Payk. Fn. Sv. 2. 215. 2.
Habitat in Europae nemoribus.

87. B. elytris integris, obſcure cuprea, elytris ſulcis quatuor albo-villoſis. *Andreae.*

Oliv. Inſ. 2. 32. tab. 1. fig. 6.
Habitat in Perſia. Muſ. D. Boſc.
Magna. Caput et thorax ſcabra, obſcura, hinc inde cupro nitidula. Elytra varioloſa, et praeterea ſulcis quatuor albo-villoſis, quorum ſecunda nec baſin nec apicem attingit. Corpus viride, minime nitens.

88. B. elytris integris ſtriatis: punctis faſciculato-piloſis, corpore hirſuto inaurato. Ent. ſyſt. 2. 203. 73. *faſcicularis.*

*Bupre-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis faſcicularis.* Linn. ſyſt. Nat. 2. 661. 12. Muſ. Lud. Vlr. 90.
Oliv. Inſ. 2. 32. tab. 4. fig. 38.
Seb. Muſ. 2. tab. 20. fig. 5.
Sulz. Inſ. tab. 6. fig. 4.
Degeer Inſ. 7. tab. 47. fig. 6.
Herbſt. Arch. tab. 28. fig. 9.
Wulf. Inſ. Cap. tab. 1. fig. 12.
Habitat ad Cap. Bon. Spei.

*manipularis.* 89. B. elytris integris violaceis: ſtrigis duabus e punctis faſciculatis albis. Ent. ſyſt. ſuppl. 134. 73.
Habitat in Tanger. Muſ. D. Lund.

*variolaris.* 90. B. elytris integris obſcuris: punctis impreſſis numeroſis, thorace carinato. Ent. ſyſt. 2. 204. 74. *
*Bupreſtis variolaris.* Oliv. Inſ. 2. 32. tab. 8. fig. 85.
Pall. Icon. tab. D. fig. 2.
Habitat in Onopordine et Echinope Ruſſiae.

*Onopordinis.* 91. B. elytris integris, obſcure aeneis: ſulcis albo-villoſis, thorace ſcabro. Ent. ſyſt. 2. 204. 75. *
Habitat in Barbariae Onopordine. D. Prof. Vahl.
*Varietas* paullo maior vix diſtincta, elytris magis glabris, ſulco tantum vnico alteroue albo-villoſo in Muſ. D. de Seheſtedt.

*piloſa.* 92. B. elytris integris, aenea, hirta, pedibus concoloribus. Ent. ſyſt. ſuppl. 134. 75. *
— Habitat in Marocco. Muſ. D. Lund.

*birta.* 93. B. elytris integris, pedibus ferrugineis, corpore hirto. Ent. ſyſt. 2. 204. 76. *
*Bupreſtis birta.* Linn. ſyſt. Nat. 2. 661. 13. Muſ. Lud. Vlr. 91.
Oliv. Inſ. 2. 32. tab. 3. fig. 18. 19.
Habitat in Africa, Barbaria.

94. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

94. B. elytris integris, ſupra obſcura, thorace vtrinque macula baſeos aurea, ſcutello porrecto ſubulato. *ſcutellaris.*

*Bupreſtis ſcutellaris.* Web. Inſ. 72. 1.
Habitat in Sumatra. D. Daldorff.
Affinis videtur B. canaliculatae. Antennae ſerratae, nigrae. Caput obſcure viride. Thorax obſcurus, poſtice vtrinque ante angulum lunula impreſſa angulo ipſo aureo, nitidiſſimo. Scutellum ſingulare, elongatum, ſubulatum. Elytra ſubpunctata, obſcura, lineis quatuor eleuatis, ante apicem coëuntibus. Corpus viridi-aeneum.

95. B. elytris integris aeneis: macula aurea, apice fuſcis, ſcutello elongato triangulari. *ſtigma.*

*Bupreſtis ſtigma.* Web. Inſ. 73. 2.
Habitat in Sumatra. D. Daldorff.
Statura omnino praecedentis, at minor. Caput obſcurum orbita oculorum viridi. Thorax punctatus, aeneus, nitidus, lunula vtrinque magna impreſſa. Scutellum elongatum, ſubulatum, aeneum. Elytra punctata lineisque quatuor eleuatis, ante apicem coëuntibus, baſi aenea, in medio macula aurea, nitida, apice obſcura. Alae cyaneae. Corpus aeneum, nitidum.

96. B. elytris integris, ſulcatis aeneis, capite thoraceque viridibus. Ent. ſyſt. 2. 204. 77. *auſtriaca.*

*Bupreſtis auſtriaca.* Linn. ſyſt. Nat. 2. 661. 9.
Oliv. Inſ. 2. 32. tab. 10. fig. 113.
Habitat in Auſtria. D. Scopoli.

97. B. elytris integris atris: vittis duabus impreſſis, cinereo-tomentoſis. Ent. ſyſt. 2. 204. 78. * *ſibirica.*

Oliv. Inſ. 2. 32. tab. 8. fig. 79.

Herbſt.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Arch. tab. 28. fig. 11.
Habitat in Sibiria. Muſ. D. Banks.

*imperialis.* 98. B. elytris integris, atra, thorace margine elytris faſciis abbreuiatis teſtaceis.
Habitat in noua Hollandia. D. de Meyerle.
Magna. Caput nigrum. Thorax punctatus, ſubcanaliculatus, ater, margine teſtaceo. Scutellum vix vllum. Elytra punctato-ſtriata, atra, nitida, faſciis quatuor teſtaceis, baſeos interrupta, dentata, ſecunda ſinuata, futuram haud attingente, tertia tenuiore abbreuiata, quarta breuiſſima. Margo apicis teſtaceus. Corpus nigrum, albo-villoſum.

*ſplendens.* 99. B. elytris integris viridibus: margine omni aureo lineisque tribus eleuatis. Ent. ſyſt. 2. 205. 79. *
Habitat in China.

*fuſca.* 100. B. elytris integris, obſcure fuſcis, ſubtus aenea, abdomine punctis lateralibus fuluis. Ent. ſyſt. 2. 205. 80. *
*Bupreſtis fuſca.* Oliv. Inſ. 2. 32. tab. 8. fig. 88.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*ruſtica.* 101. B. elytris emarginatis ſtriatis, obſcure aeneis. Ent. ſyſt. 2. 205. 81. *
*Bupreſtis ruſtica.* Linn. ſyſt. Nat. 2. 660. 8. Fn. Sv. 756.
Geoff. Inſ. 1. 126. 3.
Degeer Inſ. 4. 130. 4. tab. 4. fig. 10.
Schaeff. Icon. tab. 2. fig. 1.
Oliv. Inſ. 2. 32. tab. 3. fig. 22.
Payk. Fn. Sv. 2. 216. 3.
Habitat in Europae nemoribus.

*canaliculata.* 102. B. elytris integris, obſcura, abdomine ſubtus cana-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

canaliculato: ano quadridentato. Ent. ſyſt. 2. 205. 82.

Habitat in Africa aequinoctiali. Muſ. D. Banks.

103. B. elytris integris elongato-attenuatis truncatis, aenea, thorace elytrisque reticulato-punctatis. Ent. ſyſt. ſuppl. 135. 82. * *reticulata.*

Habitat in Italia. Dr. Allioni.

104. B. elytris integris attenuato-acuminatis obſcuris, corpore cupreo. Ent. ſyſt. 2. 206. 83. * *acuminata*

*Bupreſtis acuminata.* Oliv. Inſ. 2. 32. tab. 5. fig. 46.

Degeer Inſ. 4. 128. tab. 4. fig. 18.

Petiv. Gazoph. tab. 2. ſig. 2.

Pall. Icon. tab. D. ſig. 10.

Habitat in Ruſſia meridionali. D. Boeber.

105. B. elytris integris, nigra, obſcura, thorace macula baſeos vtrinque impreſſa. Ent. ſyſt. 2. 206. 84. * *excauata.*

*Bupreſtis excauata.* Oliv. Inſ. 2. 32. tab. fig.

Habitat Tranquebariae. Dr. Pflug.

106. B. elytris integris truncatis, ſupra obſcura, ſubtus cuprea, thorace rugoſo. Ent. ſyſt. 2. 206. 85. * *moeſta.*

Habitat in Saxonia. D. Hybner.

107. B. elytris integris ſtriatis, obſcuris, nigro-ſcabris, corpore ſubtus cupreo. Ent. ſyſt. 2. 206. 86. * *lugubris.*

*Bupreſtis lugubris.* Oliv. Inſ. 2. 32. tab. 10. fig. 106.

*Bupreſtis oxyptera.* Pall. Icon. tab. D. fig. 11.

Panz. Fn. Germ. 1. tab. 21.

Habitat in Europa auſtraliori. D. Schulz.

108. B. elytris integris, atra, antennis albis, thorace varioloſo. Ent. ſyſt. 2. 207. 87. * *cariosa.*

*Bupre-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis carioſa.* Oliv. Inſ. 2. 32. tab. 7. f. 68.
*Bupreſtis Brutia.* Petagn. Spec. Inſ. Cal. inf. 22. 108. tab. 1. fig. 20.
Habitat in Ruſſia auſtraliori. D. Boeber.

*vndata.* 109. B. viridi-aenea, elytris integris, apice obſcuris: ſtrigis vndatis albis. Ent. ſyſt. 2. 207. 88. *
Habitat in Germania. Muſ. Dr. Schulz.

*ſanguinea.* 110. B. elytris integris ſtriatis coeruleo-maculatis, corpore rufo. Ent. ſyſt. ſuppl. 135. 88. *
Habitat in Mogodor. D. Schousboe. Muſ. D. Lund.

*6punctata.* 111. B. elytris integris, exſcutellata, aenea, elytris obſcurioribus: punctis tribus impreſſis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo B. chryſoſtigmae. Caput aeneum, impreſſum. Thorax vix punctatus, aeneus, nitidus: punctis duobus dorſalibus, impreſſis, poſtice productus absque ſcutello. Elytra integra, laeuia, obſcuriora: punctis tribus impreſſis. Pedes aenei, tarſis cyaneis.

*tenebricoſa* 112. B. elytris integris ſubſtriatis, obſcura, abdomine ſubtus cyaneo, auro-punctato. Ent. ſyſt. 2. 207. 89. *
*Bupreſtis tenebricoſa.* Oliv. Inſ. 2. 32. tab. 5. fig. 48.
*Bupreſtis Tenebrionis.* Roſſ. Fn. Etr. 1. 184. 459.
Habitat in Gallia meridionali. Muſ. D. Olivier.

*Tenebrionis.* 113. B. elytris integris truncatis, atra, thorace dilatato varioloſo. Ent. ſyſt. 2. 207. 90. *
*Bupreſtis Tenebrionis.* Linn. ſyſt. Nat. 2. 661. 11.
Sulz. Hiſt. Inſ. tab. 6. fig. 16.
Oliv. Inſ. 2. 32. tab. 6. fig. 16.

Herbſt.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Arch. tab. 28. fig. 15.
Cyrilli Ent. Neapol. 1. tab. 1. fig. 13.
Habitat in Europa, Africa.

114. B. elytris integris, viridi-aenea, thorace punctis duobus, elytris quinque atris. Ent. ſyſt. ſuppl. 136. 90. * *4punctata*

Habitat in Galliae Alpibus. D. Brogniard.

115. B. elytris integris, faſciis cinereis vndatis, corpore cylindrico nigro. Ent. ſyſt. 2. 208. 91. * *Rubi.*

*Bupreſtis Rubi.* Linn. ſyſt. Nat. 2. 602. 14.
Oliv. Inſ. 2. 32. tab. 4. fig. 29.
Panz. Fn. Germ. 29. tab. 20.
Habitat in Europae Rubo.

116. B. elytris integris: baſi faſciisque vndatis cinereis, corpore aeneo cinereo-villoſo. Ent. ſyſt. ſuppl. 136. 91. * *angulata.*

Habitat in India. Muſ. D. de Seheſtedt.

117. B. elytris integris: margine aureo, ſupra obſcure viridis, ſubtus cuprea. Ent. ſyſt. 2. 208. 92. * *aurulenta.*

*Bupreſtis aurulenta.* Oliv. Inſ. 2. 32. tab. 5. fig. 51.
Habitat Halae Saxonum. D. Hybner.

118. B. elytris integris, viridis, thorace punctato, pedibus apice cyaneis. Ent. ſyſt. 2. 208. 93. * *transſata.*

Habitat in Marocco. Muſ. D. de Seheſtedt.

119. B. elytris integris, obſcure viridibus, thorace lineis duabus obſcurioribus, antennis coeruleis. Ent. ſyſt. 2. 208. 94. * *cyanicornis.*

*Bupreſtis Trochilus*, elytris integris, aurea nitidiſſima, thoracis dorſo, elytris pedibusque viridibus. Ent. ſyſt. 2. 205. 70. * vix differt.

*Bupre-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis cyanicornis.* Oliv. Inſ. 2. 32. tab. 2. fig. 11. tab. 3. fig. 20.
*Bupreſtis Stephanelli.* Petagn. ſpec. Inſ. Cal. 23. 110.
*Bupreſtis femorata.* Vill. Ent. 1. 338. 34. tab. 1. fig. 40.
Panz. Fn. Germ. 29. tab. 21.
Habitat in Gallia meridionali. Muſ. D. Olivier.

*2maculata* 120. B. elytris integris ſtriatis: macula rubra, corpore fuſco-viridi. Ent. ſyſt. 2. 209. 95. *
*Bupreſtis 2maculata.* Linn. ſyſt. Nat. 2. 662. 16.
Muſ. Lud. Vlr. 92.
Oliv. Inſ. 2. 32. tab. 12. fig. 140.
Sulz. Hiſt. Inſ. 57. tab. 6. fig. 15.
Habitat in India.

*4maculata* 121. B. elytris integris, viridis, thorace poſtice elytrisque maculis duabus aureis. Ent. ſyſt. 2. 209. 96. *
*Bupreſtis 4maculata.* Oliv. Inſ. 2. 32. tab. 10. fig. 110.
Habitat in India. Dr. Fothergill.

*femorata.* 122. B. elytris integris, obſcura, elytris maculis duabus aureis impreſſis confluentibus, femoribus anticis vnidentatis.
Habitat in Carolina. Muſ. D. Boſc.
Statura omnino B. chryſoſtigmae. Caput obſcurum, aeneo parum nitidulum. Thorax punctatus, canaliculatus, concolor. Elytra ſubſtriata, obſcura: maculis aliquot obſcure aeneis impreſſa. Corpus obſcure metallicum, nitidulum. Femora antica incraſſata, vnidentata.

*triſtis.* 123. B. elytris integris marginatis, corpore cupreo, abdomine punctis decem nigris. Ent. ſyſt. 2. 209. 97. *

*Bupre-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis triſtis.* Linn. ſyſt. Nat. 2. 662. 18.
Muſ. Lud. Vlr. 93.
Habitat in Indiis.

124. B. elytris integris, thoracis lateribus ſcabris. *cuprea.*
ſcutello tuberculari, corpore cupreo. Ent. ſyſt. 2. 209. 97.
*Bupreſtis cuprea.* Linn. ſyſt. Nat. 2. 662. 20.
Muſ. Lud. Vlr. 94.
Habitat in America.

125. B. elytris integris, nigra, thorace punctis vtrinque tribus marginalibus, elytris ſeptem flauescentibus. Ent. ſyſt. ſuppl. 136. 98. * *14guttata.*
Habitat - - D. Weber.

126. B. elytris integris, purpuraſcentibus: punctis quinque viridibus impreſſis vivaciſſimis. *hybernata.*
Habitat in Carolina. Muſ. D. Boſc.
Statura B. candentis, at maior. Caput et thorax punctata, obſcure purpuraſcentia, minime nitida. Elytra punctata, laetius purpuraſcentia: punctis quinque 1. baſeos, 2. 2. impreſſis, viridibus, nitidis. Corpus obſcure purpuraſcens.

127. B. elytris integris, ſubſtriatis, corpore aeneo, immaculato. Ent. ſyſt. 2. 210. 100. * *metallica.*
Habitat in Gallia meridionali.

128. B. elytris integris: ſtrigis duabus inauratis, vndatis, corpore fuſco. Ent. ſyſt. 2. 210. 101. * *nobilitata.*
Habitat in Indiis.

129. B. elytris integris, ſubſtriatis, ſubtus aenea, ſupra cuprea. Ent. ſyſt. 2. 210. 103. * *bicolor.*
*Bupreſtis barbara.* Oliv. Inſ. 2. 32. tab. fig.
Habitat in Barbaria. D. Prof. Vahl.
Puncta duo minuta, approximata, thoracis baſi impreſſa.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*exaltata.* 130. B. elytris integris, ſupra viridis, thoracis ma gine corporeque ſubtus aureis. Ent. ſyſt. ſupp 137. 98.
Habitat in Cayenna.

*vmbellatarum.* 131. B. elytris integris laeuibus, corpore obſcu cupreo, immaculato. Ent. ſyſt. 2. 210. 104.
*Bupreſtis vmbellatarum.* Oliv. Inſ. 2. 32. tab. fig. 23.
Habitat in Barbariae floribus vmbellatis. D. Pro Vahl.

*adſperſa.* 132. B. elytris integris laeuibus, ferrugineo-irr ratis, corpore nigro obſcuro. Ent. ſyſt. ſupp 137. 103. *
Habitat in Africa. Muſ. D. de Seheſtedt.

*Morio.* 133. B. elytris integris laeuibus, corpore nigro, im maculato. Ent. ſyſt. 2. 210. 105. *
*Bupreſtis appendiculata*, elytris integris ſub acuminatis, corpore laeui atro, immaculato Ent. ſyſt. 2. 210. 102. *
~~Payk. Fn. Sv. 2. 230.~~ 17.
Habitat Hafniae. Muſ. D. Lund.

*4punctata.* 134. B. elytris integris punctatis, thorace puncti quatuor impreſſis, corpore obſcuro. Ent. ſyſt 2. 211. 106.
*Bupreſtis 4punctata.* Linn. ſyſt. Nat. 2. 662. 23 Fn. Sv. 759.
Degeer Inſ. 4. 134. 8.
Oliv. Inſ. 2. 32. tab. 10. fig. 117.
Herbſt. Arch. tab. 28. fig. 17.
Payk. Fn. Sv. 2. 230. 18.
Habitat in Calthae paluſtris floribus frequens

*uciata.* 135. B. coleoptris integerrimis aureis: punctis duo bus cruceque poſtico nigris. Ent. ſyſt. 2. 211 107. *

*Bupr*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Bupreſtis cruciata.* Oliv. Inſ. 2. 32. tab. 7. f. 74.
Habitat in noua Hollandia. Muſ. D. Banks.

136. B. elytris integris ſubſtriatis, obſcure aenea, thorace ſtriis impreſſis albidis. Ent. ſyſt. 2. 211. 108. * *lineata. ſtriola*

Habitat in Barbaria. Muſ. D. Desfontaines.

137. B. elytris integris obſcuris, thorace aureo: ſtriis duabus nigris. Ent. ſyſt. 2. 211. 110. * *manca.*
*Bupreſtis manca.* Linn. ſyſt. Nat. 2. 1067. 10.
Geoff. Inſ. 1. 127. 4.
Oliv. Inſ. 2. 32. tab. 2. fig. 12.
Herbſt. Arch. tab. 28. fig. 18.
Habitat in Europae nemoribus.

138. B. elytris integris fuſco-nitidis, capite thoraceque cupreis, corpore aureo. Ent. ſyſt. 2. 212. 112. * *meditabunda.*
*Bupreſtis meditabunda.* Oliv. Inſ. 2. 32. tab. 10. fig. 107.
Habitat in America boreali. Muſ. Dr. Hunter.

139. B. elytris integris, cylindrica atra, thorace vtrinque impreſſo aureo. *ignara.*
Habitat in America. Muſ. D. Boſc.
Magnitudo B. mancae, at tota cylindrica. Caput canaliculatum. Thorax vtrinque impreſſus, obſcure aureus, antice parum nigricans. Elytra laeuia, atra, obſcura. Corpus atrum.

140. B. elytris integris linearibus: linea media eleuata, corpore aeneo: atomis numeroſis cineraſcentibus. Ent. ſyſt. 2. 212 13. * *atomaria.*
Habitat in Surinamo. Muſ. D. Boſc.

141. B. elytris integris linearibus: lineis duabus eleuatis, thorace inaequali, corpore atro. Ent. ſyſt. ſuppl. 137. 113. * *plana.*
Habitat in Senegallia. Muſ. D. de Seheſtedt.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*cyanoptera.* 142. B. elytris integris linearibus, basi impressis cyaneis, capite thoraceque obscure viridibus.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura elongata B. viridis. Antennae serratae, nigrae. Caput valde canaliculatum, viride, obscurum. Thorax obscure viridis, linea dorsali maculaque magna, lateralli impressis. Elytra linearia, cyanea, basi valde impressa. Corpus cyaneum, pedibus nigris.

*viridis.* 143. B. elytris integris linearibus punctatis, corpore elongato viridi. Ent. syst. 2. 213. 114.
*Buprestis viridis.* Linn. syst. Nat. 2. 663. 25. Fn. Sv. 762.
Geoff. Inf. 1. 127. 5.
Degeer Inf. 4. 133. 6. tab. 5. fig. 1.
Oliv. Inf. 2. 32. tab. 11. fig. 124.
Herbst. Arch. tab. 28. fig. 21.
Payk. Fn. Sv. 2. 225. 12.
Habitat in Europae Betula alba.
*Variat* corpore obscuriore et cupreo.

*4guttata.* 144. B. elytris integris linearibus viridibus: puncto albo, abdomine cyaneo: punctis vtrinque tribus albis. Ent. syst. 2. 213. 115.
Oliv. Inf. 2. 32. tab. 7. fig. 75.
Roff. Fn. Etr. 191. 473.
Herbst. Arch. tab. 28. fig. 22.
Payk. Fn. Sv. 2. 224. 11.
Schaeff. Icon. tab. 279. fig. 6.
Habitat in Anglia. D. Yeats.

*deplana.* 145. B. elytris integris linearibus, corpore depresso, plano obscuro.
Habitat in Amboina. D. Billardiere.
Corpus totum depressum, planum, obscure fuscum

146. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

146. B. elytris integris linearibus viridibus, capite thoraceque obſcure aureis. Ent. ſyſt. 2. 213. 116. ✲ *linearis.*

Payk. Fn. Sv. 2. 226. 13.

Habitat in Dania. Muſ. D. de Seheſtedt.

147. B. elytris integris linearibus, corpore elongato atro. Ent. ſyſt. 2. 213. 117. ✲ *atra.*

*Bupreſtis atra.* Linn. ſyſt. Nat. 2. 663. 26.

Habitat in Germania.

148. B. elytris linearibus apice ſpinoſis, corpore obſcure aeneo. *acuta.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

A B. ſpinoſa omnino differt. Corpus magnitudine B. biguttatae, obſcure aeneum, minime nitidum. Caput valde retuſum. Thorax canaliculatus, lateribus depreſſis.

149. B. elytris integris linearibus, corpore elongato aeneo-nitido. Ent. ſyſt. 2. 213. 118. ✲ *elata.*

~~*Bupreſtis Hyperici.* Creutz. Inſ.~~ 122. 14. tab. 3. fig. 26.

Habitat Halae Saxonum. Dom. Hybner.

150. B. elytris integris linearibus, coeruleo-fuſca, ſubtus aenea. Ent. ſyſt. 2. 214. 119. ✲ *pauida.*

Habitat - - Muſ. D. de Seheſtedt.

151. B. elytris integris linearibus, atra, capite ſulcato. Ent. ſyſt. 2. 214. 120. ✲ *emarginata.*

Habitat in Gallia. Muſ. D. Boſc.

152. B. elytris integris linearibus, nigra, thorace cupreo. Ent. ſyſt. 2. 214. 121. ✲ *ruficollis.*

*Bupreſtis ruficollis.* Oliv. Inſ. 2. 32. tab. 9. fig. 101.

Habitat in America. Muſ. Dr. Hunter.

153. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*innuba.* 153. B. elytris integris, nigra, capite thoraceque obſcure aeneis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Parua, vix linearis. Caput obſcure aeneum, nitidulum, in medio canaliculatum. Thorax obſcure aeneus, nitidulus, vtrinque impreſſus. Elytra integra, vix ſtriata, nigra, immaculata. Corpus nigrum.

*capitata.* 154. B. elytris integris, atra, capite thoracisque margine aeneis nitidis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura parua B. innubae. Caput viridi-aeneum, nitidum. Thorax laeuis, ater, margine laterali, aeneo, nitido. Elytra laeuia et corpus atra.

*armata.* 155. B. elytris integris ſublinearibus, apice ſpinoſis, corpore cyaneo.

Web. Inſ. 74. 4.

Habitat in Sumatra. D. Daldorff.

Affinis certe B. ſpinoſae, at paullo maior et corpus totum cyaneum, nitidum. Forte mera ſexus varietas.

*ſpinoſa.* 156. B. elytris integris ſublinearibus viridibus, apice ſpinoſis.

Habitat in Amboina. D. Billardiere.

Parua, ſublinearis. Caput ſulco impreſſo obſcure aeneum. Thorax obſcure aeneus, maculis tribus baſeos impreſſis. Elytra punctata, viridia, apice ſpina, valida, armata. Corpus obſcure aeneum, nitidum.

*feſtiua.* 157. B. elytris integris: maculis ſex coeruleis, corpore elongato viridiſſimo. Ent. ſyſt. 2. 214. 122. *

*Bupre-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Buprestis festiua.* Linn. syst. Nat. 2. 663. 27.
Oliv. Ins. 2. 32. tab. 9. fig. 100.
Habitat in Africa.

158. B. elytris integris, thorace marginato vtrinque impresso, corpore supra viridi, subtus atro. *nitens.*
*Buprestis lucidula.* Ent. syst. 4. App. 451. 128. *
Habitat Kiliae.

159. B. elytris integris, thorace submarginato, obscura, antennis pedibusque nigris. *sepulchralis*
Habitat in Germania.
Corpus paruum obscurum, aeneo-paullo nitidulum. Thorax submarginatus, nullo modo canaliculatus. Antennae et pedes nigri.

160. B. elytris integris, thorace marginato vtrinque depresso, corpore viridi nitido. Ent. syst. 2. 214. 123. *nitidula.*
*Buprestis nitidula.* Linn. syst. Nat. 2. 662. 21.
Fn. Sv. 757.
Oliv. Ins. 2. 32. tab. 11. fig. 122.
Schaeff. Icon. tab. 50. fig. 7.
Herbst. Arch. tab. 28. fig. 20.
Payk. Fn. Sv. 2. 231. 19.
Habitat in Europae plantis.

161. B. thorace laeui, fusco-bimaculato, corpore viridi. *millefolii.*
Habitat in Austria. D. de Meyerle.
*Varietatem* β. nitentis credidi, at distincta videtur. Corpus totum subpunctatum, viride. Thorax maculis duabus dorsalibus nunc maioribus nunc minoribus.

162. B. elytris integris viridibus, capite thoraceque aureis. Ent. syst. 2. 214. 124. * *laeta.*
*Buprestis laeta.* Oliv. Ins. 2. 32. tab. 5. fig. 50.
Payk. Fn. Sv. 2. 232. 19. β.

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat Halae. D. Schaller.
Forte praecedentis mora varietas.

*ſalicis.* 163. B. elytris integris, viridis, nitens, coleoptris aureis, baſi viridibus. Ent. ſyſt. 2. 215. 125. *
Oliv. Inſ. 2. 32. tab. 2. fig. 13.
Panz. Fn. Germ. 1. tab. 12.
Herbſt. Arch. tab. 46. fig. 5.
Schaeff. Icon. tab. 31. fig. 12.
Habitat in Europae Salice. Dr. Schulz.
*Bupreſtis nitida* Roſſ. Fn. Etr. vix differt.

*Sophorae.* 164. B. elytris integerrimis, ſupra viridibus, ſubtus nigra, abdomine ſupra aureo. Ent. ſyſt. 2. 215. 126. *
Habitat in Guineae Sophora occidentali. Dr. Iſert.

*diſcoidea.* 165. B. elytris integris, aenea, villoſa, elytrorum diſco teſtaceo. Ent. ſyſt. 2. 215. 127. *
Oliv. Inſ. 2. 32. tab. 7. fig. 65. tab. 10. fig. 65.
Habitat in Barbariae plantis vmbellatis. D. Prof. Vahl.

*quercata.* 166. B. elytris integris, viridis, thorace macula elytris vitta fuſcis.
Habitat in Carolinae Quercu. Muſ. D. Boſc.
Parua, viridis. Thorax macula nunc maiore, nunc minore media, fuſca. Elytra laeuia, viridia, vitta nunc latiore, nunc anguſtiore, fuſca.

*paruula.* 167. B. elytris integris ſubſtriatis, cuprea, thorace cylindrico. Ent. ſyſt. ſuppl. 137. 127. *
Habitat in Tanger. D. Schousboe. Muſ. D. de Seheſtedt.

*cyanea.* 168. B. elytris integris rugoſis, corpore cyaneo. Ent. ſyſt. 2. 215. 128. *
*Bupreſtis tarda.* Ent. ſyſt. 2. 209. 99. *
Habitat Halae Saxonum. D. Schaller.

169. B.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

169. B. elytris integris, viridi-aenea, coleoptris aureis, maculis ſeptem viridibus. *purpuraſcens.*

Habitat in noua Cambria. D. Billardiere.

Statura et magnitudo omnino B. candentis. Caput viridi-aeneum. Thorax aeneus, lineis duabus latis, fuſcis. Elytra aurea, nunc purpuraſcentia, nunc rufo-aurea, maculis viridibus: duabus baſeos linearibus; interiore communi, media transuerſa, tertia lineari.

170. B. elytris integris obſcuris: maculis tribus viridibus impreſſis. Ent. ſyſt. ſuppl. 138. 128. * *thoracica.*

Habitat in Inſula St. Thomas Americae. Muſ. D. Lund.

171. B. elytris integris, cyanea, thorace canaliculato, elytris aureis: taenia ſuturali baſeos cyanea. Ent. ſyſt. 4. App. 451. 128. * *candens.*

*Bupreſtis candens.* Panz. Fn. Germ. 1. tab. 9.

Habitat in Germania. Dr. Panzer.

172. B. elytris integris nigris: faſciis tribus cupreis, capite thoraceque cupreis. *tricincta.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Minuta. Caput et thorax laeuia, cuprea, immaculata. Elytra laeuia, nigra, faſciis tribus cupreis, baſeos abbreuiata, ſecunda in medio, vltima apicis. Corpus atrum.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

129. TRACHYS. *Palpi* quatuor breuiſſimi, aequales.
*Maxilla* bifida.
*Antennae* moniliformes.

*teſſellata.* 1. T. nigra, elytris albo-teſſellatis.
Habitat in Carolina. Muſ. D. Boſc.
*Os* maxillis palpisque. *Palpi* quatuor breuiſſimi, aequales, filiformes, *anteriores* vix longiores, quadriarticulati; articulis ſubaequalibus, adhaerentes maxillae dorſo, *poſteriores* triarticulati: articulis aequalibus, adnati ligulae medio exteriori. *Mandibula* breuis, cornea. edentula, acuta. *Maxilla* membranacea, bifida: laciniis aequalibus, obtuſis. *Ligula* breuis, membranacea, integra. *Labium* breue, corneum, transuerſum, integrum.
Maior quam T. pygmaea. Color capitis thoracisque variat, nunc niger, nunc obſcure aurulentus. Elytra ſubſtriata ſaepius concolora, albo-teſſellata. Corpus atrum, nitidum.

*ephippium.* 2. T. cuprea, coleoptris macula dorſali ouata, atra.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo T. pygmaeae. Caput et thorax laeuia, glabra, cuprea. Scutellum atrum.
Elytra

---

Trachydis corpus paruum, oblongum, depreſſum, immarginatum: *capite* paruo, retuſo ſiue canaliculato, obtuſo, inſerto; *oculis* magnis, oblongis, prominulis, ſubmarginalibus; *antennis* thorace breuioribus, diſtantibus, reconditis, ſubter oculos inſertis; *thorace* ſubtrilobo: margine deflexo; *ſcutello* paruo, rotundato; *elytris* rigidis, ante medium coarctatis, apice anguſtatis, longitudine abdominis; *pedibus* breuibus, validis; *tarſis* quinquearticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Elytra laeuia, cuprea: macula magna, communi atra, quae baſin at haud apicem attingit.

. T. aenea, nitida, elytris cyaneis. *pygmaea.*
*Bupreſtis pygmaea.* Ent. ſyſt. 2. 211. 110.
Oliv. Inſ. 2. 32. tab. 4. fig. 34.
Habitat in Barbariae vmbellatis. D. Prof. Vahl.

.. T. nigra, elytris rugoſo-impreſſis. *rugoſa.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo T. minutae, tota nigra, obſcura, at diſtincta, elytris rugis plurimis, inaequalibus impreſſis.

;. T. nigra, elytris rugis, transuerſis, albidis, thorace ſubtrilobo, laeuia. *minuta.*
*Bupreſtis minuta.* Ent. ſyſt. 2. 212. 111.
Linn. ſyſt. Nat. 2. 663. 24. Fn. Sv. 760.
Geoff. Inſ. 1. 128. 6. Illr. 29. IV. 21
Oliv. Inſ. 2. 32. tab. 2. fig. 14.
Herbſt. Arch. tab. 28. fig. 19.
Payk. Fn. Sv. 2. 232. 20.
Habitat in Europae Corylo.

. T. nigra, elytris cyaneis, apice aureis. *terminans.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et ſumma affinitas T. exiguae. Caput et thorax laeuia, nigra, aeneo-nitida. Elytra cyanea, apice aurea, nitida.

. T. elytris nigris, immaculatis, capite thoraceque atris, nitidulis. *faſtigiata.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura omnino T. exiguae, at poſtice magis attenuata. Caput et thorax atra, nitida. Elytra nigra, immaculata.

8. T.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*pulchella.* 8. T. capite thoraceque viridibus, elytris cyaneis: maculis duabus viridibus impreſſis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura parua T. minutae. Caput et thorax laeuia, viridia, nitida Elytra laeuia, cyanea, maculis duabus magnis impreſſis, viridibus. Corpus nigrum.

*exigua.* 9. T. capite thoraceque viridi-aeneis, elytris cyaneis immaculatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Praecedenti affinis, at elytra cyanea, omnino immaculata. Corpus atrum.

*puſilla.* 10. T. aenea, elytris ſtrigis cinereis.
Habitat Tranquebariae. D. Daldorff. Muſ. D. Lund.
Statura omnino T. minutae, at duplo adhuc minor, tota aenea, ſtrigis aliquot elytrorum vndatis cinereis.

*nana.* 11. T. atra, nitida, elytris punctatis, thorace ſubtrilobo.
*Bupreſtis nana.* Payk. Fn. Sv. 2. 233. 21.
*Bupreſtis puſilla.* Oliv. Inſ. 2. 32. tab. 12. f. 133.
Habitat in Europae borealis Corylo.
T. minuta duplo minor, tota atra, nitida. Frons valde retuſa. Elytra punctata, immaculata.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

30. ELATER. *Palpi* securiformes.
*Maxilla* vnidentata, obtusa.
*Ligula* bifida.
*Antennae* filiformes.

1. E. fuscus, elytris sulcatis: sulcis rugosis, antennis flabelliformibus. *gigas.*

Habitat in Guinea. D. Meier. Mus. D. Lund.

Maximus in hoc genere et forte proprii generis. Palpi haud securiformes, nec sternum porrectum. Corpus elongatum, subcylindricum. Antennae flabellatae. Caput impressum. Thorax niger, punctis duobus dorsalibus impressis anguloque postico subspinoso. Elytra fusca, sulcata: sulcis lineolis eleuatis transuersis, rugosa. Corpus fuscum.

2. E. fuscus, antennarum fasciculo flabelliformi. Ent. syst. 2. 216. 1. * *flabellicornis*

*Elater flabellicornis*. Linn. syst. Nat. 2. 651. 1.
Oliv. Ins. 2. 31. tab. 3. fig. 28.
Drury Ins. 3. tab. 47. fig. 1.
Voet. Col. 1. tab. 45. fig. 34.

Habitat in India.

*Femina* maior, antennis tantum serratis.

3. E.

---

Elateris corpus elongatum, subcylindricum, immarginatum, tardum; *capite* paruo, rotundato, inserto; *oculis* globosis, marginalibus, vix prominulis; *antennis* longitudine thoracis, serratis, distantibus, subter oculos insertis; *thorace* plano: angulo postico vtrinque prominulo, acuto, pectorisque dente e fouea abdominis resiliente; *scutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis; *tarsis* quinquearticulatis; *colore* saepius obscuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fascicularis.* 3. E. thorace murino, elytris pallidis fusco-vndatis, antennarum fusciculo flabelliformi. Ent. syst. 2. 216. 2. *

*Elater fasciicularis.* Oliv. Inf. 2. 31. tab. 5. f. 56.
Habitat in America. Muf. Dr. Hunter.

*fpecioſus.* 4. E. albus, nigro-maculatus. Ent. fyſt. 2. 216. 3.
*Elater fpecioſus.* Linn. fyſt. Nat. 2. 652. 2.
Voet. Col. tab. 45. fig. 32.
Oliv. Inf. 2. 31. tab. 7. fig. 70.
Habitat in India.

*luridus.* 5. E. cinereo-tomentoſus, elytris ſubſtriatis, plantis fuſcis. Ent. ſyſt. 2. 217. 4. *
*Elater luridus.* Oliv. Inf. 2. 31. tab. 5. fig. 57.
Habitat in Coromandel. Muf. D. Banks.

*tomentoſus.* 6. E. niger, ferrugineo-tomentoſus, elytris ſtriatis. Ent. ſyſt. ſuppl. 138. 4. *
Habitat in India orientali.

*foueatus.* 7. E. niger, thorace punctis duobus impreſſis albis, elytris ſtriatis. Ent. ſyſt. 2. 217. 5. *
Habitat in Guinea. D. Iſert.

*myops.* 8. E. thorace ocellis duobus atris, corpore fuſco cinereoque variegato.
Habitat in Carolina. Muf. D. Boſc.
Statura magna E. oculati et affinis, at magis oblongus. Ocelli thoracis magis oblongi et corpus totum fuſco cinereoque rarium.

*oculatus.* 9. E. thorace ocellis duobus atris, corpore nigro albo-maculato. Ent. ſyſt. 2. 217. 6.
*Elater ocellatus.* Linn. ſyſt. Nat. 2. 651. 3.
Oliv. Inf. 2. 31. tab. 3. fig. 34.
Petiv. Gazoph. tab. 10. fig. 4.
Act. Angl. 246. 396. 13. et 271. 841. 32.
Voet. Col. 1. tab. 45. fig. 33.

Degeer

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 4. 159. 1. tab. 17. fig. 28.
Habitat in Americae meridionalis Quercu.

10. E. thorace ocellis duobus atris, corpore nigro immaculato. Ent. ſyſt. 2. 217. 7. * *luſcus.*
*Elater luſcus.* Oliv. Inſ. 2. 31. tab. 6. fig. 64.
Habitat in America. D. Drury.

11. E. ater, thorace maculis duabus ocellaribus, elytris ſtrigis duabus arcuatis punctoque poſtico ferrugineis. Ent. ſyſt. 2. 217. 8. * *coccus.*
Habitat in Africa. D. Lee.

12. E. ater, thorace elytrisque rufo-lineatis. Ent. ſyſt. 2. 218. 9. * *lineatus.*
*Elater lineatus.* Oliv. Inſ. 2. 31. tab. 6. fig. 63.
Habitat in Braſilia. Muſ. Gallic.

13. E. thoracis lateribus, macula flaua glabra. Ent. ſyſt. 2. 218. 10. *noctilucus*
*Elater noctilucus.* Linn. ſyſt. Nat. 2. 657. 4.
Muſ. Lud. Vlr. 82.
Oliv. Inſ. 2. 31. tab. 2. fig. 14.
Degeer. Inſ. 4. 160. 2. tab. 13. fig. 1.
Brown. Iamaic. 432. tab. 44. fig. 10.
Sloan. Iamaic. 2. 206. tab. 237. fig. 1.
Voet. Coleopt. 1. tab. 43. fig. 16.
Habitat in Americae meridionalis Inſulis.

14. E. thorace poſtice maculis duabus glabris, flauis. Ent. ſyſt. 2. 218. 11. *phoſphoreus.*
*Elater phoſphoreus.* Linn. ſyſt. Nat. 2. 652. 5.
Oliv. Inſ. 2. 31. tab. 2. fig. 20. tab. 2. fig. 14.
Degeer. Inſ. 4. 161. 3. tab. 18. fig. 2.
Habitat in Americae Inſulis.

15. E. fuſcus, thorace nigro: margine flauo. Ent. ſyſt. 2. 218. 12. * *ignitus.*
*Elater ignitus.* Oliv. Inſ. 2. 31. tab. 8. fig. 78.

*Ela-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Elater Indicus*. Herbſt. Arch. tab. 27. fig. 3.
Voet. Col. 1. tab. 43. fig. 18.
Habitat in Cayenna. D. v. Rohr.

*lambens.* 16. E. fuſcus, ferrugineo-tomentoſus, thorace punctato, elytris ſtriatis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Affinis certe E. ignito, at margo thoracis concolor. Corpus totum obſcurum tomento tenerrimo, ferrugineo tectum.

*fuſcipes.* 17. E. ater, antennis pedibusque fuſcis. Ent. ſyſt. 2. 218. 13. *
*Elater fuſcipes*. Oliv. Inſ. 2. 31. tab. 3. fig. 21.
Habitat in India orientali. Dr. Koenig.

*4maculatus.* 18. E. cinereus, thorace punctis quatuor, elytris faſciis duabus nigris. Ent. ſyſt. 2. 219. 14. *
*Elater quadrimaculatus*. Oliv. Inſ. 2. 31. tab. 8. fig. 79.
*Elater faſciatus*. Drury Inſ. 3. tab. 47. fig. 2.
Habitat in Africa aequinoctiali.

*rubiginoſus.* 19. E. nigricans, griſeo-maculatus, thoracis limbo elytrorumque macula magna pallidis. Ent. ſyſt. 2. 219. 15. *
Habitat in Sierra Leon Africae. Dr. Pflug.

*ligneus.* 20. E. ferrugineus, elytris mucronatis, antennis nigris. Ent. ſyſt. 2. 219. 16. *
*Elater ligneus*. Linn. ſyſt. Nat. 2. 652. 7. Amoen. acad. 6. 395. 24.
Oliv. Inſ. 2. 31. tab. 2. fig. 15.
Drury Inſ. 3. tab. 47. fig. 4.
Voet. Col. 1. tab. 42. fig. 9.
Habitat in America meridionali.

*lacunoſus.* 21. E. niger, thorace punctato quinquefoueato, elytris ſtriatis.

Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Carolina. Muſ. D. Boſc.
Magnitudo fere E. rufi. Caput et thorax punctata, nigra, immaculata. Thorax praeterea foueolis quinque dorſalibus, impreſſis: 2. 1. 2: intermedio maiore. Elytra ſtriata, nigra. Plantae parum ferrugineae.

22. E. ater, elytris ſtriatis, ſcutello griſeo. Ent. ſyſt. ſuppl. 138. 16. * *Morio.*
Habitat in America. D. Hybner.

23. E. ater, nitidus, antennis pedibusque fuſcis, elytris laeuiſſimis. Ent. ſyſt. ſuppl. 138. 16. * *laeuigatus.*
Habitat in America. D. Hybner.

24. E. ferrugineus, capite thoraceque obſcurioribus. Ent. ſyſt. 2. 219. 17. * *rufus.*
Panz. Fn. Germ. 10. tab. 10.
*Elater rufus.* Payk. Fn. Sv. 3. 1. 1.
Degeer Inſ. 4. tab. 5. fig. 13.
Habitat in Germania. D. Smidt.

25. E. thorace elytrisque ferrugineis, thoracis margine poſtico corporeque nigris. Ent. ſyſt. 2. 220. 18. * *ferrugineus.*
*Elater ferrugineus.* Linn. ſyſt. Nat. 2. 654. 20.
Fn. Sv. 729.
Geoff. Inſ. 1. 130. 1. tab. 2. fig. 4.
Panz. Fn. Germ. 10. tab. 11.
Oliv. Inſ. 2. 31. tab. 3. fig. 35.
Schaeff. Icon. tab. 19. fig. 1.
Habitat in Europa.

26. E. viridi-aeneus, elytris ſulcatis: ſulcis villoſo-albis. Ent. ſyſt. 2. 220. 19. *porcatus.*
*Elater porcatus.* Linn. ſyſt. Nat. 2. 652. 6.
Oliv. Inſ. 2. 31. tab. 7. fig. 74.
Drury Inſ. 3. tab. 47. fig. 6.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Sulz. Hiſt. Inſ. tab. 6. fig. 7.
Habitat in America meridionali.

*ſulcatus.* 27. E. villoſo-albidus, elytris ſtriis tribus elevatis glabris. Ent. ſyſt. 2. 220. 20. *
*Elater ſulcatus.* Oliv. Inſ. 2. 31. tab. 2. fig. 10.
Habitat in America meridionali. D. Spengler.

*ſtriatus.* 28. E. nigricans, opacus, elytris ſtriatis: ſtriis albo-pubeſcentibus.
*Elater ſtriatus.* Linn. ſyſt. Nat. 2. 652. 8.
Habitat in America meridionali. Muſ. D. Lund.
Affinis E. virenti, at diſtinctus videtur. Elytra ſtriis nouem, villis albis tectis.

*virens.* 29. E. villoſus, nigro-viridis, antennis nigris. Ent. ſyſt. 2. 220. 21. *
*Elater virens.* Oliv. Inſ. 2. 31. tab. 2. fig. 19. tab. 9. fig. 55.
Voet. Col. 1. tab. 42. fig. a.
Habitat in Americae Inſulis. Muſ. Dr. Hunter.

*fulgens.* 30. E. cyaneo-nitidus, ſubtus cupreus, elytris acuminatis. Ent. ſyſt. 2. 220. 22. *
*Elater fulgens.* Oliv. Inſ. 2. 31. tab. 4. fig. 43.
Habitat in India. Muſ. Britt.

*inaequalis.* 31. E. cyaneo-niger. thorace inaequali, elytris ſtriatis rubris. Ent. ſyſt. 2. 221. 23. *
Oliv. Inſ. 2. 31. tab. 5. fig. 47.
Habitat in Italia. Muſ. D. Boſc.

*bicornis.* 32. E. ater, nitidus, capite bidentato, elytris ſtriatis apice acuminatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Corpus elongatum, atrum, nitidum. Antennae ſerratae. Caput antice dentibus duobus porrectis, acutis. Thorax punctatus: margine poſtico

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

postico tridentato. Elytra substriata, apice acuminata.

33. E. obscurus, thorace rugoso bidentato. *bidens.*
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Corpus medium, elongatum, nigrum. Antennae serratae. Caput paruum, ferrugineo-pubescens. Thorax strigis transuersis rugosus pone marginem anticum, dentibus duobus eleuatis, acutis. Elytra rugosa, obscura, nigra, griseo parum mixta.

34. E. thorace atro nitido, elytris striatis nigris. *aterrimus.*
Ent. syst. 2. 221. 24.
*Elater aterrimus.* Linn. syst. Nat. 2. 653. 17. Fn. Sv. 726.
Geoff. Inf. 1. 136. 73.
Oliv. Inf. 2. 31. tab. 5. fig. 53.
Habitat in Europae hortis.

35. E. niger, opacus, elytris striatis, antennis pedibusque concoloribus. Linn. syst. Nat. 2. 221. 25. * *niger.*
*Elater niger.* Linn. syst. Nat. 2. 656. 36. Fn. Sv. 743.
Oliv. Inf. 2. 31. tab. 6. fig. 65.
Habitat in Europae hortis.

36. E. nigricans, cinereo varius, thorace canaliculato. *marmoratus.*
Habitat in Carolina. Mus. D. Bosc.
Statura omnino E. murini, at maior. Caput et thorax punctata, nigra, cinereo maculata, thorace in medio canaliculato. Antennae serratae, piceae. Scutellum cinerenm. Elytra substriata, nigro cinereoque varia. Corpus cum pedibus nigricans.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*murinus.* 37. E. thorace obſcure cinereo, elytris cinereo-nebuloſis, plantis rufis. Ent. ſyſt. 2. 221. 26.
*Elater murinus.* Linn. ſyſt. Nat. 2. 655. 22. Fn. Sv. 738.
Oliv. Inſ. 2. 31. tab. 2. fig. 9.
Schaeff. Icon. tab. 4. fig. 6.
Voet. Col. 1. tab. 44. fig. 26.
Payk. Fn. Sv. 3. 34. 38.
Degeer Inſ. 4. 150. 10.
Habitat in Europae hortis frequens.

*pulueru-lentus.* 38. E. niger, atomis numeroſiſſimis cinereis, pedibus teſtaceis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura E. murini paullo minor. Caput, thorax, elytra, corpus nigra, pileis rarioribus, cinereis, vti atomis irrorata. Pedes teſtacei. Caput impreſſum. Thorax poſtice retuſus.

*boloſericeus.* 39. E. obſcurus, thorace elytrisque cinereo fuſcoque nebuloſis ſubholoſericeis. Ent. ſyſt. 2. 222. 27.
Oliv. Inſ. 2. 31. tab. 3. fig. 33.
tab. 7. fig. 69.
Geoff. Inſ. 1. 135. 9.
Voet Col. tab. 44. fig. 27.
Payk. Fn. Sv. 3. 35. 39.
Habitat in Germania. D. Smidt.

*fuſcus.* 40. E. villoſus, fuſcus, thorace punctato, elytris crenato-ſtriatis.
Habitat in Amboina. D. Billardiere.
Statura et magnitudo E. murini ruſticus, totus obſcurus, villoſus, fuſcus. Pedes rufo parum nitent.

41. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

11. E. elytris aeneis: maculis pallidioribus confertis, vnguibus rubris. Ent. syst. 2. 222. 28. * *tessellatus.*

*Elater tessellatus.* Linn. syst. Nat. 2. 659. 22.
Fn. Sv. 738.
Oliv. Inf. 2. 31. tab. 3. fig. 22.
Herbst Arch. tab. 27. fig. 5.
Schaeff. Icon. tab. 4. fig. 7.
Voet. Col. 1. tab. 44. fig. 27.
Payk. Fn. Sv. 3. 7. 9.
Degeer Inf. 4. 148. 7.
Habitat in Europae hortis.

12. E. corpore atro, atomis numerosissimis albis. *atomarius.*
Ent. syst. suppl. 139. 28. *

*Elater atomarius.* Linn. syst. Nat. 2. 655. 28.
Fn. Sv. 739.
*Elater carbonarius.* Oliv. Inf. 2. 31. tab. 2. f. 11.
*Elater pulverulentus.* Panz. Fn. Germ.
Habitat in Germania. Dr. Panzer.

13. E. thorace nigro pallidoque vario, elytris nigricantibus: fascia vndata alba. Ent. syst. 2. 222. 29. *fasciatus.*

*Elater fasciatus.* Linn. syst. Nat. 2. 655. 27.
Fn. Sv. 737.
Oliv. Inf. 2. 31. tab. 5. fig. 46.
Panz. Naturf. 24. 32. 45. tab. 1. fig. 46.
Payk. Fn. Sv. 3. 15. 18.
Habitat in Europae hortis.

14. E. niger, thoracis margine elytrorumque basi fasciaque postica villoso-flauescentibus. Ent. syst. 2. 222. 30. * *varius.*

*Elater varius.* Oliv. Inf. 2. 31. tab. 3. fig. 26.
tab. 2. fig. 68.
Herbst. Arch. tab. 27. fig. 11.
Habitat Parisiis. Mus. D. Bosc.

45. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*lacteus.* 45. E. albo-villoſus, thorace punctis; elytris macu-
lis nigris.
Habitat in Sumatra. D. Daldorff. Muſ. de Sche-
ſtedt.
Statura et magnitudo E. faſciati. Corpus totum
albo-villoſum, thorace punctis, elytris laeui-
bus, maculis aliquot nigris. Elytra apice emar-
ginata.

*excauatus.* 46. E. thoracis dorſo foueato, nigro, cinereo hinc
inde pubeſcente.
Habitat in Guinea. Muſ. D. Lund.
Statura et magnitudo omnino E. faſciati. Anten-
nae ſerratae, nigrae. Caput nigrum, cinereo-
villoſum. Thorax niger, cinereo-villoſus: fo-
uea lata, quae tamen nec baſin, nec apicem at-
tingit. Elytra ſtriata, ſubcrenata, nigra, baſi
vtrinque ad ſcutellum retuſa. Corpus nigrum.

*aeneus.* 47. E. thorace elytrisque aeneis, antennis ſimplici-
bus nigris. Ent. ſyſt. 2. 223. 31.
*Elater aeneus.* Linn. ſyſt. Nat. 2. 655. 31. Fn.
Sv. 740.
Oliv. Inſ. 2. 31. tab. 8. fig. 83.
Degeer Inſ. 4. 149. 8.
Payk. Fn. Sv. 3. 16. 19.
β. *Elater germanus*, thorace elytrisque atro-
coeruleis, corpore toto pedibusque nigris.
Linn. ſyſt. Nat. 2. 655. 30. Fn. Sv. 730.
Oliv. Inſ. 2. 31. tab. 2. fig. 12.
Sulz. Hiſt. Inſ. tab. 6. fig. 8.
Habitat in Europae hortis.

*impreſſus.* 48. E. pubeſcens, niger, thorace linea dorſali pun
ctisque duobus impreſſis, elytris ſtriatis, aeneo
nitidulis. Ent. ſyſt. 2. 223. 32.
*Elater impreſſus.* Payk. Fn. Sv. 3. 18. 21.
Habitat in Suecia. D. Schneider.

49. E

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

49. E. thorace elytrisque aeneis, antennis mares pectinatis. Ent. syst. 2. 223. 33. *pectinicornis.*
*Elater pectinicornis.* Linn. syst. Nat. 2. 655. 42.
Fn. Sv. 741.
Geoff. Inf. 1. 133 7.
Oliv. Inf. 2. 31. tab. 1. fig. 4. tab. 6. fig. 4.
Sulz. Inf tab. 5. fig. 36.
Schaeff. Icon. tab. 2. fig. 4.
Payk. Fn. Sv. 3. 9. 11.
Degeer Inf. 4. 145. 2. tab. 5. fig. 3.
Habitat in Europae hortis frequens.

50. E. subpubescens, obscure cupreus, scutello pallido. Ent. syst. suppl. 139. 33. * *aeruginosus.*
*Elater aeruginosus.* Oliv. Inf. 2. 31. tab. 8. f. 76.
Habitat in Gallia. Mus. D. Bosc.
*Variat* elytris concoloribus et aeneis.

51. E. thorace ferrugineo: linea dorsali nigra, coleoptris testaceis attenuatis: striis quinque nigris, capite furcato. Ent. syst. 2. 224. 34. *furcatus.*
Habitat in India. Mus. Britt.

52. E. thorace vtrinque vnidentato ferrugineo: linea dorsali nigra, coleoptris testaceis: vittis tribus atris. Ent. syst. 2. 224. 35. * *futuralis.*
*Elater futuralis.* Oliv. Inf. 2. 31. tab. 1. fig. 3.
*Elater angulatus.* Drury Inf. 3. tab. 47. fig. 5.
Habitat in Senegallia. Mus. D. Olivier.

53. E. fuscus, elytrorum vitta pedibusque testaceis. Ent. syst. 2. 224. 36. * *vittatus.*
Habitat in Germania. D. Smidt.

54. E. cupreus, elytris dimidiato-flauis. Ent. syst. 2. 225. 37. * *cupreus.*
*Elater cupreus.* Oliv. Inf. 2. 31. tab. 5. fig. 50.
Schaeff. Icon. tab. 38. fig. 2.
Habitat in Anglia.

55. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*cruciatus.* 55. E. thorace nigro: lateribus ferrugineis, coleoptris flauis: margine cruceque nigris. Ent. ſyſt. 2. 225. 38.
*Elater cruciatus.* Linn. ſyſt. Nat. 2. 653. 12. Fn. Sv. 722.
Oliv. Inſ. 2. 31. tab. 4. fig. 40.
Geoff. Inſ. 1. 133. 6.
Degeer Inſ. 4. 149. 9.
Sulz. Hiſt. Inſ. tab. 6. fig. 10.
Voet. Col. 2. tab. 43. fig. 12.
Payk. Fn. Sv. 3. 36. 11.
Habitat in Europae hortis.

*Syriacus.* 56. E. thorace rufo, elytris faſcia duplici alba, corpore nigro. Ent. ſyſt. 2. 225. 39.
*Elater Syriacus.* Linn. ſyſt. Nat. 2. 653. 11. Muſ. Lud. Vlr. 83.
Habitat in Oriente.

*caſtaneus.* 57. E. thorace teſtaceo pubeſcente, elytris flauis: apice nigris, corpore nigro. Ent. ſyſt. 2. 225. 40.
*Elater caſtaneus.* Linn. ſyſt. Nat. 2. 654. 18. Fn. Sv. 727.
Geoff. Inſ. 1. 132. 4.
Oliv. Inſ. 2. 31. tab. 3. fig. 25. tab. 5. fig. 51.
Degeer Inſ. 4. 153. 15.
Schaeff. Icon. tab. 11. fig. 9. tab. 31. fig. 4.
Habitat in Europae campis.

*latus.* 58. E. ouatus, obſcure griſeus, elytris ſtriatis.
Habitat in Auſtria. D. Scheidler.
Reliquis omnibus latior et breuior. Corpus totum obſcure aeneum, pilis griſeis pubeſcens. Elytra ſtriata.

*liuens.* 59. E. niger, thorace glaberrimo rubro, elytris teſtaceis. Ent. ſyſt. 2. 225. 41. *

*Ela-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Elater liuens.* Linn. ſyſt. Nat. 2. 654. 19. Fn. Sv. 728.
*Elater diſpar.* Payk. Fn. Sv. 2. 37. 42.
Oliv. Inſ. 2. 31. tab. 5. fig. 52.
Degeer Inſ. 4. 154. 17.
Schaeff. Icon. tab. 11. fig. 8.
Habitat in Europae hortis.

10. E. thorace elytrorumque margine ferrugineis, corpore elytrisque nigris. Ent. ſyſt. 2. 226. 42. *meſomelas*
*Elater meſomelas.* Linn. ſyſt. Nat. 2. 653. 16. Fn. Sv. 725.
Oliv. Inſ. 2. 31. tab. 5. fig. 54.
Habitat in Europae Quercu.
*Mas* praecedentis teſte D. de Paykull.
*Larua* hexapoda, ſiliformis, duriuſcula, caudae ſcuto corneo: lateribus dentatis, ſubtus tuberculo conico, tubuloſo, pediformi.

11. E. linearis, ater, thorace elytrisque rufis. Ent. ſyſt. 4. App. 451. 42. * *denticollis.*
Habitat in Germaniae Quercu. D. Prof. Helwig.

12. E. thorace rufo: medio fuſco, elytris linearibus teſtaceis. Ent. ſyſt. 2. 226. 43. * *linearis.*
*Elater linearis.* Linn. ſyſt. Nat. 2. 655. 15. Fn. Sv. 723.
*Lepturoides linearis.* Herbſt. Arch. 4. 103.
Oliv. Inſ. 2. 31. tab. 7. fig. 67.
Degeer. Inſ. 4. 154. 17.
Payk. Fn. Sv. 3. 3. 4.
Habitat in Europa boreali.
*Variat* elytrorum ſutura nigra.

13. E. piceus, thorace elytrisque obſcure nigris. Ent. ſyſt. 2. 226. 44. * *obſcurus.*
*Elater obſcurus.* Linn. ſyſt. Nat. 2. 655. 25. Fn. Sv. 735.

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 2. 31. tab. 8. fig. 76.
Schaeff. Icon. tab. 19. fig. 2.
Habitat in Europae hortis.

*bicolor.* 64. E. ferrugineus, elytris ſtriatis nigris.
Habitat in Tanger. D. Schousboe. Muſ. D. de Seheſtedt.
Paullo maior E. obſcuro. Caput fuſcum, antennis ferrugineis. Thorax punctatus, ferrugineus, immaculatus. Elytra ſtriata, nigra. Corpus ferrugineum.

*punctico-lis.* 65. E. niger, thorace punctato, elytris crenato-ſtriatis, antennis tibiisque teſtaceis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura omnino et magnitudo E. flauipedis, at diſtinctus. Caput et thorax punctata, nigra. Elytra crenato-ſtriata, parum pubeſcentia, concolora. Antennae valde ſerratae et pedes ferruginea.

*flauipes.* 66. E. thorace obſcure nigro, elytris ſtriatis fuſcis, pedibus teſtaceis. Ent. ſyſt. 2. 226. 45.
*Elater flauipes.* Oliv. Inſ. 2. 31. tab. 2. fig. 17.
Habitat in America. D. v. Rohr.

*rufidens.* 67. E. thorace fuſco: angulis poſticis rufis, elytris pubeſcentibus ſtriatis, obſcure rufis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo E. caſtanipedis. Caput et thorax fuſca, obſcura, angulis poſticis acutis, rufis. Elytra pubeſcentia, ſtriata, obſcure rufa. Corpus fuſcum, pedibus rufis.

*caſtanipes.* 68. E. fuſcus, cinereo-ſubuilloſus, antennis tibiisque teſtaceis.

*Ela-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Elater caſtaneus.* Ent. ſyſt. 2. 226. 46. *
Habitat in Inſula Guadeloupe. Muſ. D. de Seheſtedt.

69. E. ferrugineus, elytris nigro-punctatis. Ent. ſyſt. 2. 227. 47. * *maculatus*
*Elater maculatus.* Oliv. Inſ. 2. 31. tab. 2. f. 18.
Habitat in America meridionali.

70. E. ater, thorace rufo, antennarum articulo tertio baſi ſpinoſo. *ſpinicornis*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Animalculum ſingulare, E. ruficolli paullo maius. Caput nigrum, antennis ſerratis: articulo primo longiori, ſubincuruo, ſecundo breui, rotundato, tertio paullo longiori; baſi dente valido exſerto. Thorax gibbus, rufus. Elytra ſtriata, atra. Corpus fuſcum.

71. E. fuſcus, obſcurus, elytris ſtriatis, piceis, abdominis margine ferrugineo. *haemorrhoidalis.*
Habitat in Auſtria. D. Scheidler.
Statura et magnitudo E. filiformis. Caput et thorax ſubpunctata, fuſca, obſcura. Elytra ſtriata, picea. Abdomen piceum, margine ſegmentorumque marginibus ferrugineis. Pedes nigri.

72. E. villoſus, cinereo-fuſcus, immaculatus. Ent. ſyſt. 2. 227. 48. * *filiformis.*
*Elater filiformis.* Oliv. Inſ. 2. 31. tab. 4. f. 41.
Habitat in Italia. Dr. Allioni.

73. E. ferrugineus, thorace lineis duabus, elytris margine nigris. *2lineatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus medium, ferrugineum. Caput immaculatum. Thorax punctatus: lineis duabus, marginem anticum haud attingentibus, nigris. Scutellum

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tellum nigrum. Elytra ſtriata, ferruginea: margine exteriore nigro. Sutura in medio parum nigricat.

*Alopex.* 74. E. fuſco-ferrugineus, fuluo-tomentoſus, elytris ſtriatis: macula baſeos obſoleta, rufa.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura praecedentis, paullo minor. Corpus totum obſcure ferrugineum, tomento fuluo tectum. Elytra ſtriata, baſi parum rufa.

*triſtis.* 75. E. thorace atro-nitido, elytris baſi margineque exteriori liuidis. Ent. ſyſt. 2. 227. 49.
*Elater triſtis.* Linn. ſyſt. Nat. 2. 655. 26. Fn. Sv. 736.
Oliv. Inſ. 2. 31. tab. 4. fig. 39.
Schaeff. Icon. tab. 194. fig. 1.
Payk. Fn. Sv. 3. 29. 33.
Degeer Inſ. 4. 158. 20.
Habitat in Europae ligno putreſcente.

*marginatus.* 76. E. thorace fuſco, elytris teſtaceis: margine vndique nigro. Ent. ſyſt. 2. 227. 50.
*Elater marginatus.* Linn. ſyſt. Nat. 2. 655. 26. Fn. Sv. 736.
Oliv. Inſ. 2. 31. tab. 8. fig. 29.
Act. Nidroſ. 3. 16. tab. 6. fig. 6.
Payk. Fn. Sv. 3. 5. 6.
Habitat in Europae plantis.

*thoracicus* 77. E. niger, thorace rufo. Ent. ſyſt. 2. 227. 51.
Geoff. Inſ. 1. 132. 5.
Oliv. Inſ. 2. 31. tab. 3. fig. 24.
Voet. Col. 1. tab. 42. fig. 6.
*N. 29.* Schaeff. Icon. tab. 31. fig. 3.
Habitat in Anglia et Gallia.

78. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

78. E. niger, thorace poſtice rubro, nitido. Ent. ſyſt. 2. 227. 52. *ruficollis.*
*Elater ruficollis.* Linn. ſyſt. Nat. 2. 655. 14.
Fn. Sv. 724.
Oliv. Inſ. 2. 31. tab. 6. fig. 61.
Degeer Inſ. 4. 153. 16.
Payk. Fn. Sv. 3. 38. 43.
Habitat in Europae graminoſis.

79. E. thorace rufo: medio nigro, elytris corporeque ferrugineis. Ent. ſyſt. 2. 228. 53. *brunneus.*
*Elater brunneus.* Linn. ſyſt. Nat. 2. 653. 10.
Fn. Sv. 721.
Oliv. Inſ. 2. 31. tab. 3. fig. 30.
Schaeff. Icon. tab. 151. fig. 5. 6.
Herbſt. Arch. tab. 27. fig. 6.
Degeer Inſ. 4. 158. 19.
Payk. Fn. Sv. 3. 28. 32.
Habitat in Europae plantis.

80. E. capite thoraceque nigris, fugaci-aureis, elytris ſtriatis caſtaneis. *fugax.*
Habitat in Auſtria. D. Scheidler.
Statura et magnitudo E. brunnei. Caput et thorax nigra, tomento fugaci, aureo, tecta. Elytra ſtriata, teſtacea. Pedes teſtacei, femoribus nigris.

81. E. niger, thorace pubeſcente fuluo, elytris ſtriatis ſanguineis. Ent. ſyſt. 2. 228. 54. * *haematodes.*
*Elater haematodes.* Oliv. Inſ. 2. 31. tab. 1. f. 6.
*Elater purpureus.* Herbſt. Arch. tab. 27. fig. 10.
*Elater ſanguineus.* Sulz. Hiſt. Inſ. tab. 6. fig. 9.
Schaeff. Icon. tab. 2. fig. 6.
Habitat in Germania.

82. E. niger, thorace elytrisque fuluis. *pyrrhopterus.*
*Elater pyrrhopterus.* Oliv. Inſ. 2. 31. tab. 7. f. 66.
Habi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Gallia. Muſ. D. Boſc.

Statura et magnitudo omnino E. haematodae. Caput atrum. Thorax punctatus, canaliculatus, fuluus. Scutellum nigrum. Elytra ſtriata, fulua. Corpus nigrum.

*ſanguineus* 83. E. ater, elytris ſtriatis, ſanguineis, immaculatis. Ent. ſyſt. 2. 228. 53.

*Elater ſanguineus*. Linn. ſyſt. Nat. 2. 654. 21. Fn. Sv. 731.

Geoff. Inſ. 1. 131. 2.

Oliv. Inſ. 2. 31. tab. 1. fig. 7. tab. 5. fig. 48.

Degeer Inſ. 4. 151. 13.

IV. 39.31 Schaeff. Icon. tab. 31. fig. 7.

Panz. Fn. Germ. 5. tab. 13.

Payk. Fn. Sv. 3. 33. 37.

Habitat in Europae hortis.

*Varietas* maior, thorace magis fuluo pubeſcente, elytris magis coloratis lituraque obſoleta baſeos nigra; vix differt.

*ephippium.* 84. E. ater, elytris ſtriatis ſanguineis: macula communi dorſali nigra. Ent. ſyſt. 2. 228. 54. *

Oliv. Inſ. 2. 31. tab. 5. fig. 48. b.

Herbſt Arch. tab. 27. fig. 9.

IV 39 31 Schaeff. Icon. tab. 31. fig. 5.

Voet. Col. tab. 43. fig. 11.

Panz. Fn. Germ. 5. tab. 14.

Habitat in Germaniae Pinis putreſcentibus.

*Varietatem* praecedentis credit D. de Paykull.

*praeuſtus.* 85. E. ater, elytris ſtriatis ſanguineis, apice nigris. Ent. ſyſt. 2. 229. 57. *

VIII 30 IV 31 *Elater ſanguineus*. Payk. Fn. Sv. 3. 33. 37. β.

Habitat in Germaniae Pinis putreſcentibus. Dom. Smidt.

*teſtaceus.* 86. E. niger, elytris ſtriatis pedibusque teſtaceis. Ent. ſyſt. 2. 229. 58. *

*Ela-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Elater teſtaceus.* Payk. Fn. Sv. 3. 15. 16.
Habitat in Germania. D. Schneider.

87. E. ater, capite thoracisque lateribus aureis. *pennatus.*
Habitat in Carolina. Muſ. D. Beſc.
Statura et magnitudo praecedentium. Caput aureum, nitidum, antennis ſerratis, nigris. Thorax canaliculatus, ater, lateribus late aureis. Elytra crenato-ſtriata, atra, immaculata.

88. E. elytris antice dimidiato ruſis, corpore nigro. *balteatus.*
Ent. ſyſt. 2. 229. 59.
*Elater balteatus.* Linn. ſyſt. Nat. 2. 654. 22.
Fn. Sv. 732.
Oliv. Inſ. 2. 31. tab. 8. fig. 77.
Degeer Inſ. 4. 152. 14.
Schaeff. Icon. tab. 77. fig. 2.
Voet. Col. 1. tab. 43. fig. 10.
Payk. Fn. Sv. 2. 31. 35.
Habitat in Europae hortis.

89. E. niger, obſcurus, elytris ſtriatis: macula oblonga marginali baſeos flauicante. Ent. ſyſt. 4. App. 452. 59. * *lateralis.*
Habitat in Germania. D. Schneider.

90. E. ater, elytris teſtaceis, apice nigris. Ent. ſyſt. 2. 229. 60. * *elongatulus.*
*Elater elongatulus.* Oliv. Inſ. 2. 31. tab. 6. f. 58.
Habitat in Auſtria. D. Saldoner.

91. E. ſanguineus, capite thoracis linea dorſali elytrorumque apicibus nigris. Ent. ſyſt. 2. 229. 61. * *melanocephalus.*
*Elater melanocephalus.* Oliv. Inſ. 2. 31. tab. 4. fig. 36.
Thunb. nou. ſp. Inſ. 3. 63.
Habitat in Coromandel. Muſ. D. Banks.

92. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*veſpertinus* 92. E. teſtaceus, thorace vittis duabus, elytris margine ſuturaque nigris.
Habitat in Carolina. Muſ. D. Boſc.
Magnitudo E. ſputatoris. Caput fuſcum, antennis rufis. Thorax vix punctatus, teſtaceus: vittis duabus dorſalibus, latis, nigris. Elytra ſtriata, teſtacea, margine ſuturaque nunc latius, nunc anguſtius nigris. Corpus teſtaceum.

*diſcoideus.* 93. E. ater, coleoptris ſtriatis: limbo albo.
Habitat in Carolina. Muſ. D. Boſc.
Medius. Caput et thorax punctata, atra, immaculata. Elytra ſtriata, atra, margine baſeos lateralique late albo. Sutura atra. Corpus atrum.

*ſputator.* 94. E. thorace fuſco, nitido, elytris cinereis, corpore nigro. Ent. ſyſt. 2. 230. 62.
*Elater ſputator.* Linn. ſyſt. Nat. 2. 654. 24. Fn. Sv. 734.
Oliv. Inſ. 2. 31. tab. 4. fig. 31.
Degeer Inſ. 4. 151. 11.
Schaeff. Icon. tab. 19. fig. 2.
Payk. Fn. Sv. 3. 20. 23.
Habitat in Europae hortis.

*analis.* 95. E. niger, elytris, ano pedibusque teſtaceis. Ent. ſyſt. 2. 230. 63. *
Habitat in Germania.

*elegans.* 96. E. niger, thoracis marginibus, elytrorum vitta pedibusque teſtaceis. Ent. ſyſt. 2. 230. 64. *
Habitat in Americae meridionalis Inſulis. Dom. Smidt.

*filum.* 97. E. filiformis, niger, antennis pedibusque teſtaceis.
Habitat in Auſtria. D. de Meyerle.
Statura et magnitudo E. elegantis, at omnino filiformis. Thorax vix canaliculatus, niger, obſcurus. Elytra ſtriata, nigra, immaculata.

98. E.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

98. E. thorace fuſco-obſcuro, elytris ſtriatis teſta- *variabilis.*
ceis. Ent. ſyſt. 2. 230. 65. *
*Elater variabilis* Payk. Fn. Sv. 3. 26. 30.
*Elater obtuſus.* Degeer Inſ. 4. 147. 5. tab. 5. fig. 19.
Habitat in Europae truncis emortuis. D. Smidt.

99. E. fuſcus, cinereo-ſubuilloſus, elytris ſtriatis. *piloſus.*
Ent. ſyſt. 2. 231. 66. *
*Elater piloſus.* Leske Itiner. 1. 11. 1. tab. A. fig. 1.
Payk. Fn. Sy. 3. 25. 29. IV. 92 ~~VIII. 90~~.
Habitat in Luſatiae montibus. D. Prof. Leske.

100. E. ater, elytris ſtriatis, obſcure aeneis. Ent. *melancho*
ſyſt. ſuppl. 139. 66. * IV. 92 *licus.*
*Elater melancholicus.* Payk. Fn. Sv. 3. 17. 20.
Habitat in Dalekarliae Alpibus. D. de Paykull.

101. E. tomentoſus, liuens, thorace elytrisque ob- *longicollis.*
ſcurioribus. Ent. ſyſt. 2. 231. 67. *
*Elater longicollis.* Oliv. Inſ. 2. 31. tab. 8. f. 81. ~~VIII. 91.~~ IV. 93
Habitat in Gallia.

102. E. fuſcus, elytris ſtriatis, antennis pedibusque *pallipes.*
pallidis. Ent. ſyſt. 2. 231. 68. *
Habitat Tranquebariae. Dom. Hybner.

103. E. niger, elytris fuſcis. Ent. ſyſt. 2. 231. 69. * *ſtriatus.*
*Elater ſtriatus.* Payk. Fn. Sv. 3. 27. 31.
*Elater ſecalis.* Act. Holm. 1774. 4. 254. f. 1-3. ~~VIII. 91.~~ IV. 93
*Elater lineatus.* Oliv. Inſ. 2. 31. tab. 3. fig. 32.
Habitat larua in radicibus ſegetis, agros deua-ſtans. D. de Paykull.

104. E. nigricans, fulùo-tomentoſus, capite ely- *erythrocc-*
trorumque maculis duabus ferrugineis. *phalus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Medius. Caput nigrum, antennis fuſcis. Thorax punctatus, fuluo-tomentoſus, nigricans. Elytra ſubſtriata, tomentoſa, nigricantia: maculis duabus magnis, ferrugineis. Pedes ferruginei.

*rufipes.* 105. E. ater, thorace nitido, elytris ſtriatis, pedibus rufis. Ent. ſyſt. 2. 231. 70. *

*Elater rufipes.* Oliv. Inſ. 2. 31. tab. 7. fig. 72.

Geoff. Inſ. 1. 136. 14.

Payk. Fn. Sv. 3. 39. 45.

Habitat Halae Saxonum. D. Hybner.

*minutus.* 106. E. thorace atro nitido, elytris pedibusque nigris. Ent. ſyſt. 2. 231. 71. *

*Elater minutus.* Linn. ſyſt. Nat. 2. 656. 34. Fn. Sv. 744.

Oliv. Inſ. 2. 31. tab. 6. fig. 62.

Payk. Fn. Sv. 3. 40. 46.

Degeer. Inſ. 4. 159. 21.

Habitat in Europa.

*tridentatus.* 107. E. cinereus, thorace poſtice tridentato. Ent. ſyſt. ſuppl. 139. 71. *

Habitat Tranquebariae. D. Daldorff. Muſ. D. de Seheſtedt.

*pallens.* 108. E. ater, elytris ſtriatis pedibusque teſtaceis. Ent. ſyſt. 2. 233. 72. *

Geoff. Inſ. 1. 139. 19.

Habitat in Gallia. Muſ. D. Boſc.

*limbatus.* 109. E. thorace atro-nitido, elytris teſtaceis: limbo nigro. Ent. ſyſt. 2. 232. 73. *

*Elater limbatus.* Thunb. nou. ſp. Inſ. 3. 64.

Oliv. Inſ. 2. 31. tab. 7. fig. 73.

Payk. Fn. Sv. 3. 43. 49.

Habitat Kiliae. Muſ. D. Lund.

110.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

110. E. nigro-aeneus, elytris ſtriatis, pedibus rufis. *riparius.*
Ent. ſyſt. 2. 232. 74. *
*Elater politus*, ater, nitidus, elytris ſtriatis, pedibus anticis pallidis. Ent. ſyſt. 2. 234. 82. * vix differt.
*Elater riparius.* Payk. Fn. Sv. 3. 41. 47.
Panz. Fn. Germ. 34. tab. 12.
Habitat ad ripas Sueciae boreales. D. de Paykull.

111. E. nigro-aeneus, thorace nitido, antennis pedibusque nigris. *Bructeri.*
Panz. Fn. Germ. 34. tab. 13.
Habitat in Auſtriae montibus. D. de Meyerle.
Magnitudo E. riparii, at magis oblongus. Thorax punctatus, nitidulus. Elytra ſtriata, obſcuriora. Pedes nigri.

112. E. ater, elytris ſtriatis pedibusque obſcure cinereis. Ent. ſyſt. 2. 232. 75. * *aduena.*
Habitat ad Bizertam in Hiſpania. D. Prof. Vahl.

113. E. elytris ſtriatis: maculis duabus, antennis pedibusque flauis. Ent. ſyſt. 2. 232. 76. *nitidulus.*
Habitat in Cayenna. D. v. Rohr.

114. E. capite thoraceque atris, elytris nigris: maculis flaueſcentibus, pedibus flauis. Ent. ſyſt. 2. 233. 77. * *pulchellus.*
*Elater pulchellus.* Linn. ſyſt. Nat. 2. 656. 35. Fn. Sv. 745.
Oliv. Inſ. 2. 31. tab. 4. fig. 38.
Herbſt. Arch. tab. 27. fig. 7.
Payk. Fn. Sv. 3. 22. 25.
Habitat in Europae hortis.

115. E. ferrugineus, thoracis dorſo, elytrorum macula faſciaque nigris. *angulatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paruus. Caput nigrum. Thorax vix punctatus niger, margine laterali et poſtico teſtaceis. Elytra ſtriata, ferruginea. Macula ante medium oblonga faſciaque poſtica vtrinque dilatata nigris. Corpus ferrugineum.

*ſcriptus.* 116. E. ferrugineus, thoracis linea elytrorum faſciis tribus angulatis atris.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et magnitudo omnino praecedentis. Caput nigrum, antennis ferrugineis. Thorax vix punctatus, rufus margine tenuiſſimo lineaque dorſali latiore nigris. Elytra ſtriata, ferruginea: faſciis tribus nigris, anticis vngulatis, vltima apicis.

*V nigrum.* 117. E. ferrugineus, thoracis linea dorſali elytrorumque V nigris.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura praecedentium, at paullo minor. Caput ferrugineum. Thorax ferrugineus, nitidus, linea dorſali atra. Elytra ſtriata, ferruginea, pone medium V nigro-notata. Corpus ferrugineum.

*2guttatus.* 118. E. niger, elytris ſtriatis: macula media rufa. Ent. ſyſt. 2. 233. 78. *

*Elater 2guttatus.* Oliv. Inſ. 2. 31. tab. 6. fig. 59.

Habitat in Gallia meridionali. Muſ. D. Olivier.

*equeſtris.* 119. E. ater, nitens, elytris faſcia lata fulua. Ent. ſyſt. ſuppl. 140. 78. *

*Elater merdelloides.* Jaqu. Miſcellan.

*Elater cinctus.* Panz. Fn. Germ.

Habitat in Germania. Dom. Mauerhoff.

120

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

120. E. niger, elytris teſtaceis: punctis duobus nigris. Ent. ſyſt. ſuppl. 140. 78. * *2punctatus*

Habitat in Tanger. Dom. Schousboe. Muſ. D. Lund.

121. E. niger, elytris teſtaceis, apice nigris: puncto albo. Ent. ſyſt. 2. 233. 79. * *2maculatus.*

*Elater maculatus.* Oliv. Inſ 2. 31. tab. 5. fig. 10.
Roſſ. Fn. Etr. 1. tab. 3. fig. 10.

Habitat in Gallia meridionali. Dom. Olivier.

122. E. ater, nitidus, elytris teſtaceis, apice atris. *melanurus*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura parua praecedentium. Caput et thorax atra, nitida, immaculata. Elytra laeuia, teſtacea: margine exteriori apiceque late atris. Corpus atrum.

123. E. niger, elytris ſtriatis, teſtaceis: maculis tribus nigris. Ent. ſyſt. 2. 233. 80. * *3maculatus.*

Habitat Halae Saxonum. Dom. Hybner.

124. E. ſupra niger, thoracis limbo maculisque duabus elytrorum teſtaceis. Ent. ſyſt. 2. 233. 81. * *4maculatus.*

Habitat in Italia. Dr. Allioni.

125. E. ferrugineus, nitidus, elytris ſtriatis: faſcia media nigra. *vnifaſciatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Corpus paruum, ferrugineum. Elytra ſtriata, faſcia media ad marginem exteriorem dilatata, nigra. Antennae et pedes ferruginea.

126. E. laeuis, ater, elytris punctis tribus albis. Ent. ſyſt. 2. 234. 83. * *6guttatus.*

*Elater ſexguttatus.* Thunb. nou. ſp. 3. 65. fig. 75.

Habitat ad Cap. Bon. Spei.

127.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Alni.* 127. E. ater, nitidus, elytris ſtriatis rufis: macula poſtica oblonga nigra.

Habitat in Suecia. Dom. de Gyldenhal. Muſ. D. Lund.

Paruus Caput et thorax atra, nitida, immaculata. Elytra ſtriata, rufa, pone medium macula oblonga, lineari, nigra. Corpus nigrum, antennis pedibusque rufis.

*puſillus.* 128. E. ater, elytris teſtaceis: ſutura atra.

Habitat in Germania. D. Prof. Helwig. Muſ. D. de Seheſtedt.

Paruus. Caput et thorax atra, nitida, laeuia. Elytra vix ſtriata, flaua: ſutura atra.

*pygmaeus.* 129. E. niger, obſcurus, tibiis piceis. Ent. ſyſt. 2. 234. 84. *

*Elater pygmaeus.* Payk. Fn. Sv. 3. 42. 48.

Habitat in Germaniae truncis putreſcentibus. Dom. Smidt.

*variegatus* 130. E. ater, thorace elytrisque cinereo-variegatis.

Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.

Corpus paruum, latiuſculum. Thorax lateribus lineaque dorſali duplicata cinereis. Elytra baſi parum, faſcia media lata: puncto nigro, apiceque cinereis. Pedes nigri.

*3faſciatus.* 131. E. cinereus, thoracis dorſo elytrorum ſutura faſciisque tribus nigris.

Habitat in America meridionali. D. Smidt. Muſ. de Seheſtedt.

Corpus minutum. Caput cinereum, immaculatum. Thorax fuſcus, margine cinereo. Elytra ſubſtriata, cinerea, ſutura faſciisque tribus nigris, quarum prima abbreuiata.

132.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

132. E. thorace rufo: nigro bimaculato, elytris baſi rufis, apice nigris. Ent. ſyſt. 2. 234. 86. * *notatus.*
*Elater notatus.* Oliv. Inſ. 2. 31. tab. 4. fig. 37.
Habitat in Coromandel. Muſ. Dom. Banks.

133. E. thorace ferrugineo, elytris atris: faſciis duabus flauis. Ent. ſyſt. 2. 235. 87. * *feſtiuus.*
*Elater feſtiuus.* Thunb. nou. ſp. 3. 64. fig. 74.
Habitat in Aegypto. Prof. Forſkahl.

134. E. niger, nitidus, elytris puncto baſeos rubro. Ent. ſyſt. 2. 235. 88. * *2puſtulatus*
*Elater 2puſtulatus.* Linn. ſyſt. Nat. 2. 642. 9.
Geoff. Inſ. 1. 136. 15.
Oliv. Inſ. 2. 31. tab. 2. fig. 13.
Schaeff. Icon. tab. 104. fig. 6.
Herbſt. Arch. tab. 27. fig. 8.
Voet. Coleopt. 1. tab. 44. fig. 22.
Payk. Fn. Sv. 3. 30. 34.
Habitat in Germaniae nemoribus.

135. E. niger, elytris ſtriatis: puncto albo. *2notatus.*
*Elater binotatus.* Roſſ. Fn. Etr. Mant. 59. 144.
Habitat in Italia. Muſ. D. Lund.
Statura parua E. 2 puſtulati. Caput et thorax fuſca, immaculata. Elytra ſtriata, puncto medio albo. Antennae et pedes ferruginea.
Variat puncto altero baſeos rufo.

136. E. thorace ferrugineo: dorſo atro, elytris ſtriatis nigris: maculis duabus poſticis rufis. *4punctatus*
Habitat in India. D. Daldorff. Muſ. D. Lund.
Statura parua praecedentium. Caput atrum, antennis ferrugineis. Thorax laeuis, rufus, dorſo late atro. Elytra ſtriata, nigra: maculis duabus pone medium rufis. Corpus nigrum, pedibus rufis.

137.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*4puſtulatus.* 137. E. niger, elytris ſtriatis: punctis duobus teſtaceis. Ent. ſyſt. 2. 235. 89. *

*Elater 4puſtulatus.* Payk. Fn. Sv. 3. 44. 51.

Habitat Halae Saxonum. Dom. Hybner.

Minutus, diſtinctus, maculis duobus elytrorum teſtaceis.

## 131. LVCANVS. *Ligula* bifida, penicillata. *Antennae* pectinato-fiſſiles.

*Alces.* 1. L. mandibulis exſertis apice quadridentatis. Ent. ſyſt. 2. 236. 1. *

Oliv. Inſ. 1. 8. 1. tab. 2. fig. 3.

Petiv. Gazoph. tab. 57. fig. 15.

Habitat in Aſia.

*Giraffa.* 2. L. mandibulis exſertis, depreſſis: dentibus plurimis diſformibus, labio rotundato. Ent. ſyſt. 4. App. 452. 1. *

*Lucanus Giraffa.* Oliv. Inſ. 1. 1. tab. 5. fig. 16.

Habitat in Aſia.

*Cervus.* 3. L. mandibulis exſertis, vnidentatis, apice bifurcatis, labio deflexo: ruga transuerſa, eleuata. Ent. ſyſt. 2. 236. 2.

*Lucanus Cervus.* Linn. ſyſt. Nat. 2. 559. 1. Fn. Sv. 405.

Geoff. Inſ. 1. 61. 1. tab. 1. fig. 1.

Roeſ. Inſ. 2. Scarab. 1. tab. 4. 5.

Degeer

---

*Lucani* corpus magnum, oblongum, depreſſum, glabrum, immarginatum, tardum; *capite* magno, plano, exſerto, fere quadrato; *mandibulis* ſaepius exſertis, corniformibus; *ocellis* rotundatis, marginalibus; *antennis* diſtantibus, ſubter apicem capitis prominulum inſertis, longitudine thoracis; *thoracc* transuerſo, vtrinque deflexo, ſcutello rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* validis, curſoriis; *tarſis* quinquearticulatis; *colore* nigro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 4. tab. 12. fig. 1. 2.
Schaeff. Icon. tab. 133. fig. 1.
Herbſt. Col. 3. tab. 33. fig. 1. 2.
β. *Lucanus Dorcas*. Myll. Zool. Dan. 444.
*Platycerus fuſcus*. Geoff. Inſ. 1. 60. 2.
Roeſ. Inſ. 2. tab. 5. fig. 8.
Schaeff. Icon. tab. 133. fig. 2.
Oliv. Inſ. 1. 8. 1. tab. 2. fig. 3. 6.
Habitat in Europae ligno putreſcente.
*Larua* obeſa, albida, capite pedibusque ferrugineis.

4. L. mandibulis exſertis, vnidentatis, apice bifurcatis, labio deflexo, conico. Ent. ſyſt. 2. 237. 3. * *Elaphus.*
Oliv. Inſ. 1. 12. 4. tab. 3. fig. 7.
Habitat in Virginia.

5. L. mandibulis exſertis: denticulis mediis diſſormibus, apice biſurcatis. Ent. ſyſt. 2. 237. 4. * *Capreolus.*
*Lucanus Capreolus*. Linn. ſyſt. Nat. 2. 560. 2. Amoen. acad. 6. 391. 4. Muſ. Lud. Vlr. 30.
*Lucanus Capra*. Oliv. Inſ. 1. 12. 3. tab. 1. fig. 1.
Voet. Coleopt. tab. 29. fig. 3.
Sulz. Hiſt. Inſ. tab. 2. fig. 1.
Herbſt. Col. tab. 39. fig. 4.
Habitat in Germania.

6. L. mandibulis exſertis, intus bidentatis, longitudine capitis. Ent. ſyſt. 2. 237. 5. * *Dama.*
Degeer Inſ. 4. tab. 19. fig. 11. 12.
Oliv. Inſ. 1. 15. 8. tab. 2. fig. 4.
Habitat in Virginia.

7. L. mandibulis exſertis, tridentatis, niger, femoribus ferrugineis. Ent. ſyſt. 2. 237. 6. * *femoratus.*
Oliv. Inſ. 1. 17. 10. tab. 4. fig. 12.
Habitat in Cajenna. Muſ. Dom. Banks.

8.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Biſon.* 8. L. mandibulis exſertis, multidentatis, thoracis elytrorumque margine rubro. Ent. ſyſt. 2. 238. 7. *

*Lucanus Biſon.* Oliv. Inſ. 1. 13. 5. tab. 3. fig. 6. Voet. Coleopt. tab. 30. fig. 6.
Habitat in America meridionali.

*Gazella.* 9 L. mandibulis intus bidentatis, ater, elytrorum margine teſtaceo. Ent. ſyſt. 2. 238. 8. *
*Lucanus Gazella.* Oliv. Inſ. 1. 13. tab. 4. fig. 13.
Habitat in Siam. Muſ. Dom. Banks.

*Lama.* 10. L. mandibulis exſertis, tridentatis, capite brevioribus, thorace angulato. Ent. ſyſt. 2. 238. 9. *
*Lucanus Lama.* Oliv. Inſ. 1. 14. 17. t. 3. f. 8.
Habitat in India orientali.

*Futuralis.* 11. L. mandibulis exſertis, baſi dentatis teſtaceis: linea dorſali atra. Ent. ſyſt. 2. 238. 10. *
*Lucanus dorſalis.* Oliv. Inſ. 1. 16. 9. tab. 5. fig. 12.
Habitat — —

*Saiga.* 12. L. mandibulis exſertis multidentatis, labio abbreuiato emarginato. Ent. ſyſt. 4. Append. 452. 3. *
*Lucanus Saiga.* Oliv. Inſ. 1. tab. 5. fig. 18.
*Lucanus Elaphus.* Herbſt. Col. tab. 33. fig. 6. Voet. Col. 1. tab. 30. fig. 5.
Habitat in America meridionali.

*Taurus.* 13. L. mandibulis exſertis, longitudine capitis, intus bidentatis, corpore ſubpunctato.
Habitat in Sumatra. D. Daldorff.
Magnitudo L. parallelepipedi. Mandibulae exſertae, intus dentibus duobus validis, approximatis, obtuſis. Corpus totum obſcurum punctis paruis impreſſum.

Alter

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Alter forte ſexus mandibulis capite longioribus, capiteque antice quadridentato, elytris ſtriatis.

14. L. mandibulis exſertis vnidentatis, capite acuminato. *acuminatus.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Minor L. Tauro. Caput planum, punctatum, antice acuminatum, margineque vtrinque ſinuato. Mandibulae porrectae, planae, longitudine capitis, medio interiori vnidentatae. Thorax breuis, transuerſus, punctatus. Elytra ſtriata.

15. L. mandibulis exſertis lunatis, planis, vnidentatis, corpore laeui atro. *Barbaroſſa*

Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

Statura et magnitudo L. Damae. Totus glaber, ater. Mandibulae longitudine capitis ſupra planae et canaliculatae, intus vnidentatae. Abdomen parum rufo-pubeſcens.

16. L. mandibulis dente laterali eleuato, corpore depreſſo. Ent. ſyſt. 2. 239. 11. *parallelepipedus.*

*Lucanus parallelepipedus.* Linn. ſyſt. Nat. 2. 561. 6.

Geoff. Inſ. 1. 62. 3.

Oliv. Inſ. 1. 17. 11. tab. 4. fig. 9.

Voet. Col. tab. 30. fig. 7.

Schaeff. Icon. tab. 63. fig. 7.

Degeer. Inſ. 4. tab. 12. fig. 9.

Payk. Fn. Sv. 3. 47. 2.

Herbſt. Col. 3. tab. 34. fig. 5.

Habitat in Europae ſyluis.

17. L. mandibulis exſertis inermibus, capitis thoracisque lateribus punctatis. *inermis.*

Habitat in Sumatra. D. Daldorff.

Statura et ſumma affinitas L. parallelepipedi, at mandibulae inermes. Caput punctatum. Thorax

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

rix dorso laeui, nitido, lateribus punctatis. Elytra subpunctata. Corpus nigrum.

*cancroides.* 18. L. mandibulis incuruis: intus dente inoraſſato difformi, elytris punctatis ſubpubeſcentibus. Ent. ſyſt. 2. 239. 12. *

*Lucanus cancroides.* Oliv. Inſ. 1. 18. 12. tab. 4. fig. 11.

Habitat in Terra Diemenii. Muſ. D. Banks.

*lunatus.* 19. L. mandibulis exſertis lunatis: dente baſeos acuto, elytris ſtriatis.

*Lucanus lunatus.* Weber Inſ. 83. 1.

Habitat in Sumatra. D. Daldorff.

Paruus in hoc genere. Mandibulae exſertae, capite paullo longiores, lunatae, haud forficatae, acutae, baſi interiori dente acuto. Caput et thorax plana, punctata, atra, nitida. Elytra ſtriata, atra.

*piceus.* 20. L. mandibulis exſertis vnidentatis, piceus, fronte retuſa.

*Lucanus piceus.* Weber Inſ. 84. 2.

Habitat in America. D. Weber.

Magnitudo L. lunati. Caput magnum, punctatum, mandibulis exſertis: dente medio validiſſimo. Frons valde retuſa. Thorax punctatus, piceus. Elytra ſulcata, picea.

*tenebroides.* 21. L. mandibulis lunatis vnidentatis, ater, thorace marginato, elytris ſubſtriatis. Ent. ſyſt. 2. 239. 13. *

*Lucanus piceus.* Bensdorff. Act. Holm. 1785. 222. 5. tab. 8. fig. a.

Panz. Fn. Germ. 62. tab. 1.

Payk. Fn. Sv. 3. 48. 3.

Herbſt. Col. 3. 314. 15.

Habitat in Europa boreali. D. Boeber.

22.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

22. L. mandibulis exſertis vnidentatis, ater, thorace canaliculato, elytris crenato-ſtriatis. *ſtriatus.*

*Lucanus ſtriatus.* Oliv. Inſ. 1. tab. 4. fig. 14.

Habitat in India orientali.

Statura depreſſa praecedentium, at L. parallelepipedo duplo minor. Caput atrum, mandibulis porrectis, longitudine capitis, intus vnidentatis. Thorax laeuis, nitidus, canaliculatus: canali lato, abbreuiato, margine antico in medio parum eleuato. Elytra crenato-ſtriata.

23. L. coeruleſcens, mandibulis lunatis, thorace marginato. Ent. ſyſt. 2. 239. 14. *caraboides*

*Lucanus caraboides.* Linn. ſyſt. Nat. 2. 561. 7.
Fn. Sv. 407.
Geoff. Inſ. 1. 63. 4.
Schaeff. Icon. tab. 6. fig. 8.
tab. 75. fig. 7.
Degeer Inſ. 4. tab. 12. fig. 11.
Oliv. Inſ. 1. 20. 14. tab. 2. fig. 2.
Panz. Fn. Germ. 58. tab. 13.
Voet. Col. tab. 30. fig. 8.
Payk. Fn. Sv. 3. 49. 4.
Herbſt. Col. 3. 310. 10. tab. 34. fig. 6. 7.
Habitat in Europae locis ſuffocatis.

IV. 439.

24. L. vireſcens, abdomine pedibusque rufis. Ent. ſyſt. ſuppl. 139. 16. *rufipes.*

Panz. Fn. Germ. 58. tab. 14.
Herbſt. Col. 3. tab. 34. fig. 8.
Schaeff. Icon. tab. 75. fig. 7.
Habitat in Germania.
Nimis praecedenti affinis.

IV. 439

25. L. mandibulis inermibus, niger, thorace punctato, elytris ſtriatis. *punctatus.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Paullo

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paullo minor L. rufipede. Mandibulae exſertae breues, planae, acutae, inermes. Caput et thorax punctata, nigra. Elytra ſtriata, nigra. Corpus nigrum, abdominis margine parum rufescente.

132. AESALVS. *Palpi* filiformes, inaequales
*Maxilla* breuiſſima, apice palpigera.
*Antennae* lamellatae.

*ſcarabaeoides.* 1. AESALVS.
*Lucanus ſcarabaeoides.* Panz. Fn. Germ. 26. tab. 15. 16.
Habitat in Auſtria. D. de Creutzer.
*Os* maxillis palpisque. *Palpi* quatuor inaequales, filiformes: *anteriores* longiores, quadriarticulati: articulis ſubaequalibus; vltimo longiori, ouato, obtuſo adhaerentes maxillae apici, *poſteriores* breuiores, triarticulati: articulis aequalibus; vltimo obtuſo, adnati ſub labii apice. *Mandibula* breuis, cornea, arcuata, acuta, edentula. *Maxilla* breuiſſima, cornea, obtuſa, apice palpigera. *Ligula* nulla? *Labium* breue, corneum, planum, truncatum, integrum.
*Genus* ſingulare ob maxillam breuiſſimam et vix vllam.
Statura breuis, parua, Scarabaei mobilicornis. Antennae nigrae, claua picea. Mandibulae maris exſertae, lunatae, ſuperne cornu magno, erecto, compreſſo, obtuſo, mandibula maiore, feminae breues, arcuatae, muticae. Elytra nigra, ſtriis eleuatis, holoſericeis, interruptis, atris.

133. PAS-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

## 33. PASSALVS. *Palpi* filiformes.

*Maxilla* cornea, arcuata, dentata, apice acuta.

*Antennae* arcuato-lamellatae.

*interruptus.*

P. ater, vertice tuberculis tribus eleuatis: intermedio maiori compreſſo. Ent. ſyſt. 2. 240. 1.

*Lucanus interruptus*. Linn. ſyſt. Nat. 2. 560. 4.

Muſ. Lud. Vlr. 53.

Oliv. Inſ. 1. tab. 3. fig. 5.

Degeer Inſ. 4. tab. 19. fig. 13.

Sulz. Inſ. tab. 2 fig. 2.

Voet. Coleopt. tab. 29. fig. 1.

Brown Jamaic. 429. tab. 44. fig. 7.

Petiv. Gazoph. tab. 27. fig. 7,

Weber Inſ. 78. 1.

Habitat in Americae meridionalis Inſulis ſub ſaccharo putreſcente.

Mandibulae tridentatae.

*emarginatus.*

P. capite inaequali, mandibulis emarginatis, thorace laeuiſſimo.

Weber Inſ. 81. 4.

Habitat in Sumatra. Dom. Daldorff.

Statura omnino praecedentis, at diſtinctus. Mandibulae magnae, apice truncatae, emarginatae. Caput inaequale, denticulis vix determinatis, lateris ſiniſtri maioribus. Thorax laeuis ſerius, ater,

---

*Paſſali* corpus elongatum, depreſſum, planum, immarginatum; *capite* magno, inaequali, transuerſo, angulo antico prominulo acuto; *oculis* paruis, globoſis, marginalibus, vix prominulis; *antennis* diſtantibus, ſub capitis apicem inſertis, thorace breuioribus; *thorace* plano, canaliculato, ſubtus tomentoſo, ab abdomine remoto; *ſcutello* nullo; *elytris* rigidis, planis, longitudine abdominis; *pedibus* validis; *tarſis* quinquearticulatis; colore nigro, nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ater, nitidus: linea media haud impreſſa. Elytra ſtriata, atra: margine inprimis baſeos ruſo barbato. Corpus piceum.

*cornutus.* 8. P. ater, verticis cornu eleuato incuruo, elytrorum ſtriis omnibus laeuibus.

Habitat in Carolina, Muſ. Dom. Boſc.

Affinis certe P. interrupto, cuius marem olim credidi, at diſtinctus videtur paullo minor. Mandibulae craſſae, dente laterali valido, apiceque obſoleti tridentatae. Capitis cornu diſtinctum eleuatum, incuruum. Thorax ater, in medio canaliculatus punctoque vtrinque impreſſo. Elytrorum ſtriae omnes laeues.

*barbatus.* 4. P. ater, vertice antice quadridentato, elytris margine baſeos rufo-hirtis.

Habitat in Guinea. Muſ. D. de Seheſtedt.

Nimis certe affinis P. dentato, at diſtinctus videtur maior. Caput inaequale, denticuli vtrinque duobus breuibus, incumbentibus. Thorax canaliculatus, ater, nitidus, lateribus punctatis, ſubtus rufo-hirtus. Elytra crenato-ſtriata, baſi margine rufo-hirta.

*dentatus.* 5. P. vertice antice multidentato, thorace ſulcato: lateribus punctatis, elytrorum ſtriis lateralibus punctatis. Ent. ſyſt. 2. 241. 2. *

Weber Inſ. 82. 5.

Habitat in India orientali.

*bicolor.* 6. P, depreſſus, niger. ſubtus rufus, vertice inaequali, thorace canaliculato.

Habitat in Sumatra. Dom. Daldorff.

Praecedente duplo minor et magis depreſſus. Caput inaequale, denticulis plurimis, eleuatis, breuibus. Thorax laeuis, in medio valde canaliculatus. Elytra nigra, piceo-nitidula, crenato-ſtriata. Corpus rufum.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

7. P. ferrugineus, elytris teſtaceis. Ent. ſyſt. 2. 241. 3. * *minutus.*

Habitat in Americae meridionalis Inſulis. Dom. Smidt.

Corpus vix pediculo maius.

## 134. PRIONVS.

*Palpi* quatuor, filiformes.
*Maxilla* cylindrica, integra.
*Ligula* breuiſſima, membranacea.
*Antennae* ſetaceae.

1. P. thorace crenato, marginato, vnidentato, ater, elytris apice crenato-dentatis, thorace canaliculato. Ent. ſyſt. 2. 242. 2. * *fuliginoſus*

Habitat in America. Muſ. Brittann.

2. P. thorace marginato, vnidentato, rufo: mandibulis inflexis, acutis, ſterno ſpinoſo. Ent. ſyſt. 2. 243. 3. * *roſtratus.*

*Prionus roſtratus.* Oliv. Inſ. 2. 66. t. 10. f. 37.

Habitat in Siam. Muſ. D. Banks.

3. P. thorace marginato, crenato, vnidentato, niger, albo-vittatus, elytris apice crenato-dentatis. Ent. ſyſt. 2. 243. 4. * *lineatus.*

*Cerambyx lineatus.* Linn. ſyſt. Nat. 2. 624. 14. Sloan. Jam. 209. tab. 237. fig. 34.

Brown.

---

*Prioni* corpus magnum, oblongum, depreſſum, glabrum, tardum; *capite* magno, transuerſo, plano, inſerto; *oculis* transuerſis, reniformibus, marginalibus; *antennis* diſtantibus, elongatis, ſupra mandibulas inſertis; *thorace* plano, marginato: margine ſaepius dentato; *ſcutello* rotundato; *elytris* rigidis, fornicatis, immarginatis, longitudine abdominis; *pedibus* longiuſculis, curſoriis; *tarſis* quadriarticulatis; *colore* ſaepius obſcuro



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Brown. Jam. 430. tab. 43. fig. 7.
Herbſt. Arch. tab. 32. fig. 1.
Habitat in America. Muſ. Brittann.

*nitidus.* 4. P. thorace marginato, crenato, vnidentato, cyaneo, nitidus, elytris cupreis. Ent. ſyſt. 2. 243. 5. *

*Prionus nitidus.* Oliv. Inſ. 3. 66. t. 12. f. 48.
Habitat in Braſilia. Muſ. Dr. Hunter.

*Faber.* 5. P. thorace marginato, vtrinque vnidentato, elytris piceis, antennis mediocribus. Ent. ſyſt. 2. 244. 6. *

*Cerambyx Faber.* Linn. ſyſt. Nat. 2. 622. 6. Fn. Sv. 646.
Panz. Fn. Germ. 9. tab. 5.
Friſh. Inſ. 13. 22. tab. 17.
Schaeff. Icon. tab. 72. fig. 3.
tab. 111. fig. 1.
Payk. Fn. Sv. 3. 50. 1.
Habitat in Europae ramis.

*ſcabricornis.* 6. P. thorace ſubmarginato, vnidentato, ſubuilloſus, nigricans, elytris fuſcis: lineis duabus elevatis, antennis mediocribus. Ent. ſyſt. 2. 244. 7. *

*Prionus ſcabricornis.* Scopol. Carn. 174.
*Cerambyx ſcabricornis.* Oliv. Inſ. 66. tab. . fig. .
Fuesly Inſ. Helv. 13. 241. tab. 1. fig. 3.
Geoff. Inſ. 1. 210. 6.
Habitat in Europa auſtraliori. Muſ. D. Olivier.

*depſarius.* 7. P. thorace ſubmarginato, vnidentato, pubeſcente, corpore nigricante, antennis brevibus rubris. Ent. ſyſt. 2. 244. 8.

*Cerambyx depſarius.* Linn. ſyſt. Nat. 2. 624. 12.
Panz. Fn. Germ. 9. tab. 7.
Schaeff. Elem. tab. 67. fig. 3.

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

PAYK. Fn. Sv. 3. 52. 3.
Habitat in Svecia. Muſ. Dom. Zſchuck.

8. E. thorace ſubmarginato, vnidentato, pubeſcente, corpore pallido, antennis mediocribus. Ent. ſyſt. ſuppl. 141. 8. * *pallens.*
Habitat in Africa. Muſ. D. Lund.

9. P. thorace marginato, bidentato, elytris mucronatis, rufis. Ent. ſyſt. 2. 244. 9. * *mucronatus.*
Habitat in America. Muſ. D. Drury.
Corpus tomento aureo villoſum.

10. P. thorace bidentato: dentibus arcuatis, niger, elytris obtuſis, teſtaceis. Ent. ſyſt. 2. 245. 10. *arcuatus.*
Habitat in Terra Diemenii. Muſ. Dom. Banks.

11. P. thorace marginato, tridentato, niger, elytris cinereo-maculatis. Ent. ſyſt. 2. 245. 11. * *maculatus.*
*Prionus maculatus.* Oliv. Inſ. 66. tab. 4. fig. 14.
Habitat in Senegallia. Muſ. Gall.

12. P. thorace marginato, vtrinque tridentato, mandibulis porrectis, extus vniſpinoſis, antennis breuibus. Ent. ſyſt. 2. 245. 12. * *ceruicornis*
*Prionus ceruicornis.* Oliv. Inſ. 66. tab. 2. fig. 8.
*Cerambyx ceruicornis.* Linn. ſyſt. Nat. 2. 622. 3. Muſ. Lud. Vlr. 65.
Sloan. Jam. 2. tab. 237. fig. 6.
Merian. Sur. tab. 48.
Roeſ. Inſ. 2. ſcarab. 2. tab. 1. fig. 8.
Brown. Jamaic. tab. 44. fig. 8.
Aubent. Miſcell. tab. 90. fig. 3.
Voet. Coleopt. 2. tab. 2. fig. 45.
Habitat in Americae ligno Bombacis.
*Larua* edulis in deliciis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bupbtalmus.* 13. P. thorace vtrinque tridentato, ater, nitidus, antennis breuibus: articulis vltimis, apice ſpinoſis. Ent. ſyſt. 2. 246. 14. *

*Prionus ſpinicornis.* Ent. ſyſt. 2. 245. 13. *

*Prionus tranquebaricus.* Ent. ſyſt. ſuppl. 141 10. *

*Huſſarus Ceylonenſis.* Voet. Coleopt. 2. tab. 3 fig. 11.

Habitat in India orientali. Muſ. D. Lund.

*forficatus.* 14. P. thorace vtrinque tridentato, ater, elytris ferrugineis, antennis compreſſis, breuibus, mandibus elongatis, forficatis, extus vnidentatis. Ent ſyſt. ſuppl. 141. 15. *

Habitat in Tanger. Dom. Schousboe. Muſ. Dom Lund.

*breuicornis.* 15. P. thorace marginato, tridentato, atro, nitido dente poſtico obtuſiſſimo, antennis breuibus

Habitat in America boreali.

Statura et ſumma affinitas P. coriarii. Diſtinctus tamen videtur, paullo maior, antennis aliquanto breuioribus, thorace latiore, nitido: dente poſtico obſoleto.

*coriarius.* 16. P. thorace marginato, tridentato, corpore piceo antennis breuibus. Ent. ſyſt. 2. 246. 15. *

*Prionus coriarius.* Oliv. Inſ. 66. tab. 1. fig. 1

Geoff. Inſ. 1. 198. tab. 3. fig. 9.

Linn ſyſt. Nat. 2. 622. 7. Fn. Sv. 647.

Degeer Inſ. 5. 591. tab. 3. fig. 5.

Roeſ. Inſ. 2. Scar. 2. tab. 1. fig. 1. 2.

Schaeff. Elem. tab. 102.

Icon. tab. 67. fig. 3.

tab. 9. fig. 1.

IV 4f

Panz. Fn. Germ. 9. tab. 8.

Friſh. Inſ. 13. tab. 9.

Payl

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 51. 2.
Habitat in Europae Betulis putridis.

17. P. thorace vtrinque bidentato, corpore nigro, elytris ferrugineis, antennis breuibus. Ent. syst. 2. 247. 16. * *giganteus.*
*Cerambyx giganteus.* Linn. Mant. 1. 531.
Oliv. Inf. 66. tab. 6. fig. 21.
Aubent. Miscell. 1. tab. 64. fig. 2.
Habitat in Cajenna.

18. P. thorace marginato, tridentato, obscurus, pectore abdomineque ferrugineis, antennis breuibus. Ent. syst. 2. 247. 17. * *cylindricus*
*Cerambyx vnicolor* Drury. Inf. 1. tab. 37. fig. 1.
Habitat in America boreali. Dom. Blackburn.

19. P. thorace marginato, vtrinque quadridentato, elytris ferrugineis, nigro-marginatis. Ent. syst. 2. 247. 18. * *armillatus*
*Prionus armillatus.* Oliv. Inf. 66. t.. 5. f. 17.
*Cerambyx armillatus.* Linn. syst. Nat. 2. 622. 4.
Petiv. Gazoph. 525. tab. 15. fig. 6.
Habitat in India.

20. P. thorace marginato, multispinoso, mandibulis porrectis, vnidentatis, apice bifidis, pedibus anticis elongatis. Ent. syst. 2. 247. 19. * *Luzonum.*
Petiv. Gazoph. tab. 74. fig. 1.
Habitat in Insulis Luzonum. Mus. Britt.

21. P. thorace marginato, multispinoso, mandibulis porrectis, apice tridentatis, pedibus spinoso-serratis. Ent. syst. 2. 248. 20. * *serripes.*
Habitat in Africa aequinoctiali. Mus. D. Banks.

22. P. thorace marginato, multidentato, mandibulis porrectis muticis. *mandibularis.*
Oliv. Inf. 66. tab. 8. fig. 31.

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. Dr. Pflug. Muſ. D. Lund.

Magnus in hoc genere. Mandibulae capite thoraceque longiores, lunatae, acutae, omnino inermes. Caput canaliculatum, punctatum. Thorax planus, inaequalis, margine multidentato. Elytra picea, apice mucronata.

*damicornis* 23. P. thorace marginato, denticulato, mandibulis porrectis, bidentatis, antennis breuibus. Ent. ſyſt. 2. 248. 21. ♂

*Cerambyx damicornis*. Linn. Mant. 1. 532.

Degeer Inſ. 5. 97. 2.

Drury Inſ. 1. tab. 38. fig. 1.

Habitat in America.

*bifaſciatus* 24. P. thorace marginato, denticulato, corpore nigro, elytris rubris: faſciis duabus, atris, antennis breuibus. Ent. ſyſt. 2. 248. 22. ♀

*Cerambyx bifaſciatus*. Linn. ſyſt. Nat. 2. 624. 16.

Gron. Zooph. 532. tab. 14. fig. 4.

Habitat in America.

*Thomae.* 25. P. thorace marginato, crenulato, corpore nigro, elytris glabris, ferrugineis: margine luteo. Ent. ſyſt. 2. 248. 23. ♂

*Cerambyx Thomae*. Act. Soc. Berol. 4. tab. 7. fig. 9.

Habitat in Inſula St. Thom. Americae.

*bilineatus.* 26. P. thorace marginato, albo, bilineato, elytris ferrugineis: atomis albis margineque luteo. Ent. ſyſt. 2. 249. 24. ♂

Habitat in America. Dom. v. Rohr.

Vix a praecedente ſatis diſtinctus.

*Amazonus* 27. P. thorace marginato, denticulato, cyaneus, elytris baſi faſciaque rufis, apice aeneis.

Voet.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Voet. Col. tab. 2. fig. 8.

Habitat in America meridionali. D. Smidt. Muſ. de Seheſtedt.

Statura parua P. Thomae. Caput canaliculatum, cyaneum. Mandibulae exſertae, apice bifidae. Antennae longitudine corporis, cyaneae. Thorax inaequalis punctatus, cyaneus, magis obſcurus margine multidentato. Elytra punctata, cyanea: margine exteriori baſeos vsque ad faſciam faſciaque lata pone medium rufis. Elytra pone faſciam aenea, nitida. Corpus cyaneum, nitidum.

28. P. thorace marginato, crenato, capite ſub mandibulis ſpinoſo, mandibulis tridentatis. Ent. ſyſt. 2. 249. 25. * *ſpinibarbis*

*Cerambyx ſpinibarbis*. Linn. ſyſt. Nat. 2. 624. 4. Muſ. Lud. Vlr. 67.

Merian. Surin. 24. tab. 24.

Habitat in America meridionali.

29. P. thoracis lateribus ſcabris, multidentatis: dente poſteriori palmato, antennis breuibus. Ent. ſyſt. 2. 249. 26. * *palmatus.*

Oliv. Inſ. 66. tab. 7. fig. 25.

Habitat in Guinea. Muſ. D. Banks.

30. P. thoracis lateribus ſcabris, multidentatis, mandibulis breuibus, hirtis, dentatis. *dentatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura P. maxilloſi, paullo minor. Mandibulae exſertae, capite multo breuiores, latae, intus dentatae, acutae. Caput punctatum, canaliculatum. Thorax minus planus, punctatus, lateribus ſcabris, multidentatis. Elytra laeuia, picea, apice rotundata.

31.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*maxillosus* 31. P. thorace marginato, crenato, mandibulis porrectis, intus hirsutis, quadridentatis. Ent. syst. 2. 249. 27. *

Oliv. Ins. 66. tab. 1. fig. 3.
Drury Ins. 1. tab. 38. fig. 3.
Habitat in America. Mus. Britt.

*canaliculatus.* 32. P. thorace marginato, crenato: canali dorsali albo, villoso, antennis brevibus. Ent. syst. 2. 250. 28. *

Habitat in Americae Insulis. Mus. Dr. Hunter.

*cinnamomeus.* 33. P. thorace marginato, denticulato, mandibulis porrectis; tridentatis. Ent. syst. 2. 250. 28. *

*Cerambyx cinnamomeus.* Linn. syst. Nat. 2. 623. 10. Mus. Lud. Vlr. 66.
Oliv. Ins. 66. tab. 5. fig. 18.
Drury Ins. 1. tab. 40. fig. 2.
Sulz. Ins. tab. 5. fig. 2.
Merian Surinam. tab. 24.
Habitat in America meridionali.

*melanopus* 34. P. thorace marginato, dentato, mandibulis porrectis, multidentatis, elytris mucronatis. Ent. syst. 2. 250. 30. *

*Cerambyx melanopus.* Linn. syst. Nat. 2. 623. 8.
*Cerambyx crenulatus.* Drury Ins. 1. tab. 38. fig. 2.
Merian Surinam. tab. 24. fig. 1.
Habitat in America meridionali.

*crenatus.* 35. P. thorace marginato, crenato, elytris ferrugineis, capite canaliculato nigro.

Habitat in India. D. Smidt.

Magnus. Antennae nigrae, longitudine corporis. Mandibulae breues, crassae, medio interiori vnidentatae. Caput valde canaliculatum, nigrum. Thorax planus, marginatus: margine prominulo,

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

lo, crenato, obſcurus, in medio maculis duabus eleuatis, atris. Elytra rugoſa, ferruginea. Corpus obſcurum.

36. P. thorace multidentato, nigro, elytris vniden- *ſpinoſus.*
tatis, teſtaceis. Ent. ſyſt. 2. 250. 31. *
Habitat Tranquebariae. Dom. Hybner.

37. P. thorace marginato, integro, mandibulis fer- ***barbatus.***
rugineo - hirſutiſſimis, antennis mediocribus. Ent. ſyſt. 2. 250. 32. *
Habitat in America meridionali. Muſ. D. Banks.

38. P. thorace ſubemarginato, mutico, teſtaceus, an- *pectinicor-*
tennis breuibus pectinatis. Ent. ſyſt. 2. 251. *nis.*
33. *
Oliv. Inſ. 66. tab. . fig. .
Habitat in Senegallia. Muſ. Dom. Geoffroy.

39. P. thorace ſubemarginato, mutico, piceus, man- ***muticus.***
dibulis exſertis, dentatis, antennis breuibus, ſimplicibus.
Habitat in America boreali. Muſ. Dom. Banks.
Affinis videtur P. pectinicorni. Corpus paruum, totum piceum. Caput punctatum, mandibulis exſertis, medio vnidentatis apiceque emarginatis. Antennae breues, fere moniliformes: articulo primo craſſiori, vltimis compreſſis. Thorax punctatus, vix marginatus, muticus. Elytra punctata.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

135. CERAMBYX. *Palpi* quatuor, filiformes.
*Maxilla* obtusa, vnidentata.
*Ligula* bifida.
*Antennae* setaceae.

*longimanus.* 1. C. thorace spinis mobilibus, elytris basi vnidentatis apiceque bidentatis, antennis longis.
*Prionus longimanus.* Ent. syst. 2. 242. 1.
Oliv. Inf. 66. tab. 3. fig. 12.
tab. 4. fig. 12.
Linn. syst. Nat. 2. 621. 1. Mus. Lud. Vlr. 64.
Degeer Inf. 5. 102. 6.
Sloan Jam. 2. 209.
Grew. Mus. 161. tab. 13.
Roes. Inf. 2. Scarab. 2. tab. 1. fig. 2.
Aubent. Miscell. tab. 64. fig. 1.
Habitat in America meridionali.

*moschatus.* 2. C. thorace spinoso, viridis, nitens, antennis mediocribus, cyaneis. Ent. syst. 2. 251. 1.
Linn. syst. Nat. 2. 627. 34. Fn. Sv. 652.
Oliv. Inf. 67. tab. 2. fig. 7.
Degeer Inf. 5. 64. 2.
Geoff. Inf. 1. 103. 5.
Frisch. Inf. 13. tab. 11.
Schaeff. Icon. tab. 11. fig. 7.
tab. 55. fig. 1.

Voet.

---

*Cerambycis* corpus oblongum, angustum, depressiusculum, glabrum, immarginatum, tardum; *capite* ouato, inserto; *ocellis* paruis, rotundatis, marginalibus; *antennis* elongatis, inter oculos insertis; *thorace* plano, rotundato, inaequali, vtrinque spinoso; *scutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* longiusculis, cursoriis; *femoribus* compressis; *tarsis* quadriarticulatis; *colore* vario, interdum nitido.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Voet. Coleopt. 2. tab. 6. fig. 14.
Habitat in Europae Salice Moſchum ſpirans.

. C. thorace rotundato, ſpinoſo, corpore viridi, fe- *virens.*
moribus rufis. Ent. ſyſt. 2. 252. 2.
Linn. ſyſt. Nat. 2. 627. 38.
Oliv. Inſ. 67. tab. 11. fig. 78.
Drury Inſ. 1. tab. 40. fig. 1.
Brown Jam. tab. 43. fig. 8.
Habitat Larua in truncis Amyris balſamiferae Americis.
Imago Eugeniam Jambos ſpirat Dr. Schwarz.
Variat antennis corpore longioribus et breuioribus, femoribus dentatis et muticis, corpore viridi et cyaneo.

. C. thorace rotundato, ſpinoſo, cyaneus, femoribus *albitarſus.*
dentatis, rufis, tarſis poſticis niueis.
Habitat in Isle de France. Dom. Daldorff.
Statura C. nitentis, paullo maior. Caput cyaneum, nitidum, antennis corpore longioribus, nigris. Thorax inaequalis, ſpinoſus et elytra cyanea, nitida, immaculata. Pedes nigri, femoribus dentatis, rufis. Tarſi poſtici niuei.

. C. thorace rotundato, ſubſpinoſo, viridi-nitens, *nitens.*
femoribus clauatis: claua anticorum rufa. Ent. ſyſt. 2. 252. 3.
Oliv. Inſ. 67. tab. 17. fig. 107.
Voet. Coleopt. 2. tab. 6. fig. 15.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

. C. thorace ſpinoſo, aureo, viridi-nitens, elytris *micans.*
cyaneis: ſutura viridi.
Habitat in Guinea. Dom. Meier. Muſ. D. Lund.
Statura et magnitudo C. feſtiui. Caput viride, nitens, antennis corpore longioribus, nigris. Thorax rotundatus, ſpinoſus, aureus, nitidus. Elytra

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tra cyanea, ſutura late viridi. Corpus aeneum, nitidum. Pedes nigri, femoribus quatuor, anticis clauatis, rufis, poſtici clauati, compreſſi.

*ater.* 7. C. thorace rotundato, ſpinoſo, corpore viridi, antennis pedibusque rufis. Ent. ſyſt. 2. 252. 4. *
Linn. Mant. 532.
Drury Inſ. 1. tab. 39. fig. 4.
Oliv. Inſ. 67. tab. 19. fig. 141.
Habitat in Africa. Dr. Solander.

*feſtiuus.* 8. C. thorace ſpinoſo, viridi, elytris violaceis baſi viridibus, femoribus ferrugineis, vnidentatis. Ent. ſyſt. 2. 253. 5. *
Oliv. Inſ. 67. tab. 7. fig. 44.
tab. 18. fig. 44.
Habitat ad fluuium Gaboon Africae. Muſ. Dom. Banks.

*vittatus.* 9. C. thorace ſpinoſo, viridis, nitens, thorace elytrisque nigro-lineatis. Ent. ſyſt. 2. 253. 6. *
Oliv. Inſ. 67. tab. 2. fig. 10.
Voet. Col. 2. tab. 10. fig. 41.
Habitat in Braſilia. Muſ. Dom. Banks.

*velutinus.* 10. C. thorace ſpinoſo, niger, elytris vitta atra. Ent. ſyſt. 2. 253. 7. *
Oliv. Inſ. 67. tab. 6. fig. 41.
Voet. Col. 2. tab. 10. fig. 40.
Habitat in America. Dr. Schulz.

*ſericeus.* 11. C. thorace ſpinoſo, holoſericeo, ater, elytris, ſutura vittaque viridibus, femoribus rufis. Ent. ſyſt. 2. 253. 8. *
Oliv. Inſ. 67. tab. 5. fig. 37.
Habitat in Braſilia. Muſ. Dom. Olivier.

*ſuturalis.* 12. C. thorace ſpinoſo, ater, elytris, ſutura vittaque media aureis. Ent. ſyſt. 2. 254. 9. *
Oliv. Inſ. 67. tab. 6. fig. 40.

Voet.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Voet. Col. 2. tab. 10. fig. 42. 43.
Habitat in America. Dr. Hunter.

3. C. thorace ſpinoſo capiteque viridi-aeneis, nitidis, elytris obſcuris, antennis longis pedibusque rufis. Ent. ſyſt. 2. 254. 10. * *elegans.*
Oliv. Inſ. 67. tab. 5. fig. 35.
Habitat in Americae meridionalis Inſulis. Dom. Olivier.

14. C. thorace ſubſpinoſo, atro: macula laterali aurea, elytris cyaneis, atro-lineatis, antennis pedibusque rufis. *regius.*
Habitat in Guinea. Dom. Heſſe. Muſ. D. Lund.
Statura et magnitudo C. ſuturalis. Caput holoſericeum, atrum, ore rufo, mandibulis exſertis, dentatis, apice parum nigricantibus. Antennae corpore breuiores, rufae. Thorax inaequalis, ſubſpinoſus: margine antico eleuato, holoſericeus, ater, maculá magna, laterali, aurea. Elytra cyanea, baſi vittis holoſericeis, atris, ſtriata. Pedes rufi.

15. C. thorace ſpinoſo depreſſo, vireſcens, elytris cupreis, tibiis compreſſo-dilatatis. Ent. ſyſt. 2. 254. 11. * *latipes.*
Oliv. Inſ. 67. tab. 16. fig. 116.
Degeer Inſ. 7. 655. bb. tab. 49. fig. 3.
Herbſt. Arch. 7. tab. 45. fig. 11.
Habitat ad Cap. Bon. Spei.

16. C. viridis, nitens, elytris cyaneis, antennis apice albis. *albicornis.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura C. longipedis paullo minor. Antennae nigrae, articulis quatuor vltimis albis. Caput et thorax viridi-aenea, nitida, immaculata. Elytra

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tra cyanea, apice acuminata. Corpus viridi-aeneum, pedibus cyaneis.

*longipes.* 17. C. thorace ſubſpinoſo cyaneo, viridis, antennis longioribus, femoribus clauatis. Ent. ſyſt. 2. 255. 12. *

Oliv. Inſ. 70. tab. 1. fig. 3.
Degeer Inſ. 7. tab. 49. fig. 7.
Habitat ad Cap. Bon. Spei.

*cyanicornis.* 18. C. thorace ſubſpinoſo, viridis, nitens, antennis breuibus coeruleis, femoribus anticis clauatis, ferrugineis. Ent. ſyſt. ſuppl. 142. 12. *
Habitat in Aquapim Guineae. Muſ. D. Lund.

*interruptus.* 19. C. thorace ſpinoſo, ater, elytris maculis tribus linearibus albis, antennis breuibus. Ent. ſyſt. 2. 255. 13. *
Oliv. Inſ. 67. tab. 17. fig. 133.
Habitat - - Muſ. D. Olivier.

*Cerdo.* 20. C. thorace ſpinoſo, rugoſo, niger, elytris rotundatis, antennis longis. Ent. ſyſt. 2. 255. 14.
*Cerambyx Cerdo.* Linn. ſyſt. Nat. 2. 629. 39.
Oliv. Inſ. 67. tab. 10. fig. 65.
Geoff. Inſ. 1. 201. 2.
Voet. Col. 2. tab. 4. fig. 5.
Habitat in Europa.

*heros.* 21. C. thorace ſpinoſo rugoſo, niger, elytris ſubſpinoſis piceis, antennis longis. Ent. ſyſt. 2. 255. 15.
Oliv. Inſ. 67. tab. 1. fig. 1.
Geoff. Inſ. 1. 200. 1.
Friſh. Inſ. 1. 13. tab. 8.
Drury Inſ. 1. tab. 49. fig. 1.
Voet. Col. 2. tab. 5. fig. 9.
Roſſ. Fn. Etr. 1. 141. 358.
Habitat in Europae Quercu.

22.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

22. C. thorace ſubſpinoſo, niger, elytris viridibus, apice ſpinoſis. Ent. ſyſt. 2. 256. 16. * *ſpinicornis*

*Cerambyx torridus*. Oliv. Inſ. 67. tab. 14. f. 95.

Habitat in Sierra Leon Africae. Muſ. D. Banks.

23. C. thorace ſpinoſo rugoſoque, nigricans, elytris bidentatis, antennis ſpinoſis mediocribus. *denticornis*

Habitat in Guinea. D. Heſſe. Muſ. D. Lund.

Statura et magnitudo C. Bati. Antennae longitudine corporis, nigrae: articulis apice ſpinoſis. Caput nigrum. Thorax ſpinoſus, rugoſus. Elytra bidentata, nigra, pube cineraſcente micantia.

24. C. thorace ſubſpinoſo, ater, antennis mediocribus, rufo nigroque annulatis. Ent. ſyſt. 2. 256. 17. * *ater.*

Oliv. Inſ. 70. tab. 3. fig. 32.

Habitat ad Cap. Bon. Spei. Muſ. D. Olivier.

25. C. thorace rugoſo, ater, elytris laeuibus bidentatis. *bidens.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura omnino C. Cerdonis, at duplo minor. Antennae corpore breuiores, nigrae, piceo colore nitentes. Caput atrum. Thorax rotundatus, muticus, punctato-rugoſus. Scutellum cinereum. Elytra laeuia, nigra, apice bidentata. Corpus nigrum, piceo micans.

26. C. niger, thorace ſpinoſo tuberculatoque, elytris baſi ſcabris apice emarginatis, antennis longis. Ent. ſyſt. ſuppl. 141. 17. * *tuberculatus.*

Habitat in Cayenna. D. Cuvier.

27. C. thorace inermi rugoſo, niger, elytris cinereis, antennis breuibus. *voluulus.*

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat Tranquebariae. Dom. Daldorff. Muſ. D. Lund.

Statura C. bidentis, at minor. Antennae breues, nigrae. Caput nigrum. Thorax inermis, rugoſus, niger. Elytra ſubholoſericea, nigra. Corpus nigrum, abdomine cinereo-tomentoſo.

*Batus.* 28. C. thorace rugoſo, ſubſpinoſo, elytris apice ſpinoſis, antennis longis aculeato-ſpinoſis. Ent. ſyſt. 2. 256. 18.
Linn. ſyſt. Nat. 2. 615. 20. Muſ. Lud. Vlr. 69.
Oliv. Inſ. 67. tab. 5. fig. 32.
Muſ. Petropol. 652. 154.
Habitat in Americae meridionalis Inſulis.

*ferrugineus.* 29. C. thorace acute ſpinoſo-rugoſo, niger, elytris ferrugineis, antennis longis. Ent. ſyſt. 2. 256. 19. *
Linn. ſyſt. Nat. 2. 626. 25. Muſ. Lud. Vlr. 70.
Oliv. Inſ. 67. tab. 18. fig. 134.
Habitat in India orientali. D. de Slanbuſch.

*alpinus.* 30. C. thorace ſpinoſo, coleoptris faſcia maculisque quatuor atris, antennis longis. Ent. ſyſt. 2. 257. 20.
Linn. ſyſt. Nat. 2. 628. 35. Fn. Sv. 654.
Oliv. Inſ. 67. tab. 9. fig. 58.
Geoff. Inſ. 1. 202. 4.
IV. 42. Scheuchz. Itiner. 1. tab. 1. fig. 5.
Sulz. Inſ. tab. 4. fig. D.
Drury Inſ. 2. tab. 31. fig. 5.
Voet. Col. 2. tab. 9. fig. 38.
Habitat in Europae montoſis.

*ocellatus.* 31. C. thorace ſpinoſo, ater: linea dorſali fulua. elytris punctis ſubocellaribus fuluis, ano bidentato.
Oliv. Inſ. 67. tab. 13. fig. 12. d.
tab. 2. fig. 12. a. b. Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et magnitudo C. ſcalaris. Antennae corpore triplo longiores, atrae. Caput atrum, macula verticali, fulua. Thorax ſpinoſus, ater, linea dorſali fulua, parum albo-marginata. Elytra atra, punctis plurimis, ſparſis, fuluis, annulo albo cinctis. Corpus cinereo-pubeſcens, linea laterali fulua. Anus bidentatus. Pedes nigri.

32. C. thorace ſpinoſo, fuſcus: linea longitudinali alba, antennis longis. Ent. ſyſt. 2. 257. 21. * *ſcalaris.*

Voet. Col. 2. tab. 6. fig. 12.

Habitat in America meridionali. Muſ. D. Banks.

33. C. thorace ſpinoſo, viridi-aeneus, elytris teſtaceis, antennis breuibus. Ent. ſyſt. 2. 257. 27. * *ebulinus.*

Habitat ad Cap. Bon. Spei. Muſ. Dr. Hunter.

34. C. thorace biſpinoſo, rugoſo, ater, antennis longioribus ferrugineis. Ent. ſyſt. 2. 257. 28. * *Morio.*

Habitat in Cayenna. Muſ. Dr. Hunter.

35. C. thorace ſpinoſo, niger, elytris ſanguineis: macula nigra. Ent. ſyſt. 2. 257. 29. * *Koehleri.*

Linn. ſyſt. Nat. 2. 631. 50.

Geoff. Inſ. 1. 204. 6.

Oliv. Inſ. 67. tab. 3. fig. 13.

Schaeff. Icon. tab. 1. fig. 1.
tab. 153. fig. 4.

Herbſt. Arch. 9. tab. 24. fig. 6.

β. *Cerambyx badenſis.* Panz. Taſchenb. 2. 48. 6.

*Lamia hungarica.* Herbſt. Arch. 4. t. 25. f. 6.

Naturf. 19. tab. 4. fig. 1-5.

Habitat in Europa auſtraliori.

36. C. thorace ſpinoſo, ſanguineus, antennis elytrorum apicibus, abdomine pedibusque nigris. Ent. ſyſt. 2. 257. 30. * *Lundii.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 67. tab. 20. fig. 149.
Habitat Tranquebariae. Muf. D. Lund.

*Desfontainii.* 37. C. thorace fpinofo, fanguineo, nigro, maculato, ater, elytris fanguineis, bafi apiceque atris, antennis longiffimis. Ent. fyft. 2. 257. 31. *
Habitat in Barbaria. Muf. Dom. Desfontaines.

*humeralis.* 38. C. thorace fpinofo, niger, elytris macula bafeos fanguinea. Ent. fyft. fuppl. 143. 26. *
Habitat in America boreali. Dom. Herfchel.

*4dens.* 39. C. thorace fpinofo, ater, thoracis margine elytrisque teftaceis, apice nigris.
Oliv. Inf. 67. tab. 19. fig. 144.
Habitat in China. Muf. D. de Seheftedt.
Statura praecedentium. Antennae corpore longiores, atrae. Caput atrum, vertice teftaceo, dentibusque quatuor inter antennas: lateralibus teftaceis. Thorax lineis tribus, dorfalibus eleuatis, niger, lateribus teftaceis. Elytra lineis tribus eleuatis, teftacea, apice atra. Corpus atrum.

*fuccinctus.* 40. C. thorace bifpinofo, rugofo, elytris fafcia flaua, antennis longioribus, compreffis. Ent. fyft. 2. 258. 27.
Linn. fyft. Nat. 2. 627. 32. Muf. Lud. Vlr. 72.
Degeer Inf. 5. 113. 14. tab. 14. fig. 5.
Oliv. Inf. 67. tab. 7. fig. 43.
Drury Inf. 1. tab. 39. fig. 2.
Voet. Coleopt. 2. tab. 7. fig. 17.
Habitat in America meridionali.
Fafcia elytrorum nunc minor, nunc obfoleta, nunc omnino nulla.

*bicornis.* 41. C. thorace bifpinofo, tuberculatoque, flauus, elytris nigro-lineatis.
Oliv. Inf. 67. tab. 5. fig. 31.

Voet.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Voet. Col. 2. tab. 6. fig. 11.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura omnino et magnitudo C. ſtriati. Antennae corpore duplo longiores, baſi flauae, articulis apice nigris. Caput flauum, dentibus duobus, eleuatis, acutis inter antennas. Thorax biſpinoſus, ſpina poſteriore minore, flauus, tuberculisque vtrinque duobus, eleuatis, acutis, nigris. Elytra coſtata, flaua, linea marginali mediaque nigris. Corpus holoſericeo-aureum, abdomine nigro: lateribus aureis. Pedes teſtacei, poſtici nigri.

2. C. thorace ſubſpinoſo, rugoſo, ferrugineus, elytris flauo-ſtriatis, antennis longis. Ent. ſyſt. 2. 259. 28. * *ſtriatus.*
Oliv. Inſ 67. tab. 10. fig. 71.
Voet. Coleopt. 2. tab. 16. fig. 68.
Habitat in Cayenna. Muſ. Dr. Hunter.

3. C. thorace inaequali, ſubſpinoſo, ater, elytris rufis, margine ſuturaque abbreuiatis, nigris, antennis mediocribus. *abbreuiatus.*
Habitat in noua Cambria. Dom. Billardiere.
Praecedentibus minor. Antennae villoſae, atrae. Caput et thorax inaequalia, atra, immaculata. Elytra rufa, lineis tribus, eleuatis, flaueſcentibus, margine ſuturaque nigris, qui tamen color abbreuiatus, vix vltra medium extenditur. Corpus atrum.

4. C. thorace biſpinoſo, rufo, elytris laeuibus, nigris, antennis longis. Ent. ſyſt. 2. 259. 29. * *ruſipes.*
Oliv. Inſ. 67. tab. 1. fig. 3.
tab. 13. fig. 89.
Habitat in America meridionali. Muſ. Dr. Hunter.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*dimidiatus* 45. C. thorace biſpinoſo, rugoſo, flauus, nigro-pu
etatus, elytris nigris, baſi flauis, antennis m
diocribus. Ent. ſyſt. 2. 259. 30. *
Oliv. Inſ. 67. tab. 14. fig. 96.
Habitat in Cayenna. Muſ. Dr. Hunter.

*bicolor.* 46. C. thorace biſpinoſo tuberculatoque, ferrugineu
elytris vltra dimidium abdomineque atris. En
ſyſt. 2. 260. 31. *
Oliv. Inſ. 67. tab. 9. fig. 61.
Voet. Coleopt. 2. tab. 8. fig. 24.
Habitat in Cayenna. Dom. v. Rohr.

*depreſſus.* 47. C. thorace multiſpinoſus, depreſſus, niger, cin
reo-variegatus, elytris acuminatis, antenn
longis. Ent. ſyſt. 2. 260. 32. *
Oliv. Inſ. 67. tab. 5. fig. 30.
Sulz. Hiſt. Inſ. tab. 6. fig. 5.
Voet. Coleopt. tab. 13. fig. 55. 56.
Habitat in Americae Inſulis. Muſ. Dom. Banks.

*faſciatus.* 48. C. thorace ſpinoſo, cyaneus, elytris faſcia flau
antennis mediocribus, ante apicem flauis. En
ſyſt. 2. 260. 33. *.
Oliv. Inſ. 67. tab. fig. 4.
Pall. Icon. Inſ. tab. F. fig. 4.
Voet. Coleopt. 2. tab. 11. fig. 49.
Herbſt. Arch. 5. tab. 25 fig. 3.
Habitat Tranquebariae. Dr. Avering.

*barbicornis.* 49. C. thorace ſpinoſo, antennarum quatuor prim
articulis nigro-barbatis, corpore teſtaceo, n
gro, variegato. Ent. ſyſt. 2. 260. 34. *
Linn. ſyſt. Nat. 2. 625. 18.
Oliv. Inſ. 67. tab. 7. fig. 48.
Voet. Coleopt. 2. tab. 9. fig. 37.
Habitat in India orientali.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

b. C. thorace mutico, griſeo-fuſcus, elytris acuminatis: faſciis duabus, punctisque niueis, *albofaſciatus.*
Habitat in Guinea. Dom. Meier. Muſ. D. Lund.
Corpus paruum. Antennae vix longitudine corporis. Thorax rotundatus, inermis, ſtriis tribus macularibus punctisque aliquot paruis, niueis. Elytra apice acuminata: faſciis duabus, latis, abbreuiatis, punctisque aliquot paruis, niueis.

1. C. thorace ſpinoſo, elytris punctis faſciisque nigris, antennis longis. Ent. ſyſt. 2. 261. 35. *nebuloſus.*
Linn. ſyſt. Nat. 2. 627. 29. Fn. Sv. 650.
Oliv. Inſ. 66. tab. 7. fig. 47.
Panz. Fn. Germ. 14. tab. 13.
Schaeff. Icon. tab. 14. fig. 9.
Voet. Coleopt. 2. tab. 4. fig. 4. IV. 37. 43
Geoff. Inſ. 1. 204. 7.
Degeer Inſ. 5. 71. 8.
Sulz. Inſ. tab. 4. fig. c.
Habitat in Abietum truncis.

2. C. thorace ſpinoſo, fuluo, villoſo, elytris nigris: macnlata, poſtica villoſa, fulua, antennis mediocribus. Ent. ſyſt. 2. 261. 36. * *obſcurus.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

3. C. thorace ſpinoſo, fuſco, elytris laeuibus: faſciis griſeis, nigro-punctatis, antennis longiſſimis. Ent. ſyſt. 2. 261. 37. * IV. 37. 48 *griſeus.*
Panz. Fn. Germ. 14. tab. 14.
Habitat in Germania. Dom. de Loewenſkiold.

4. C. thorace ſpinoſo, griſeus, elytris lineis, eleuatis, nigropunctatis, apice fuſcis; antennis longiſſimis. Ent. ſyſt. 2. 261. 38. * *coſtatus.*
Habitat in Germania Dom. de Loewenſkiold.

5. C. thorace ſpinoſo, elytris integris: punctis tribus hiſpidis, antennis mediocribus, hirtis. Ent. ſyſt. 2. 262. 39. * *faſciculatus.*
Panz. IV. 28. 49

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 14. tab. 15.
Naturf. 24. 26. 34. tab. 1. fig. 3 4.
Payk. Fn. Sv. 3. 58. 7.
Degeer Inſ. 5. tab. 3. fig. 17.
Habitat in Germania. Dom. Hybner.

*hiſpidus.* 56. thorace ſpinoſo, elytris baſi albidis, apice biden
tatis, antennis mediocribus, hirtis. Ent. ſyſt. 2
262. 40.
~~Linn. ſyſt. Nat.~~ 2. 627. 30. Fn. Sv. 651.
Geoff. Inſ. 1. 206. 9.
~~Panz. Fn. Germ.~~ 14. tab. 16.
Oliv. Inſ. 67. tab. 11. fig. 77.
Schaeff. Icon. tab. 176. fig. 5.
Friſh. Inſ. 13. tab. 16.
~~Degeer Inſ. 5.~~ 71. 9. tab. 3. fig. 17.
Habitat in Europa boreali.

*piloſus.* 57. C. thorace biſpinoſo, elytris griſeis, apice vni
dentatis, antennis mediocribus, hirtis. Ent
ſyſt. 2. 262. 41.
Oliv. Inſ. 67. tab. 9. fig. 60.
Habitat Halae Saxonum. Dom. Hybner.

*4puncta-*
*tus.* 58. C. thorace ſpinoſo, piloſus, fuſcus, capite thor
ceque punctis duobus albis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Statura parua praecedentium. Antennae longa
ferrugineae: articulis apice fuſcis. Caput fu
ſcum, punctis duobus verticalibus, niueis. Th
rax ſpinoſus, fuſcus, punctis duobus albis. El
tra piloſa, fuſca, cinereo-varia. Femora cl
uata.

*balteatus.* 59. C. thorace ſubſpinoſo, brunneus, elytris, faſ
fuſca. Ent. ſyſt. 2. 262. 42. *
*Cerambyx balteatus.* Linn. ſyſt. Nat. 2. 627. 3
Oli

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 67. tab. 17. fig. 124.
Habitat in Luſitania.

50. C. thorace inermi, rugoſiſſimo, niger, antennis mediocribus, pedibusque piceis. Ent. ſyſt. 2. 263. 43. * *rugicollis.*

Habitat Tranquebariae. Dom. Hybner.
Forte potius Gnoma.

61. C. thorace mutico, rufus, elytris teſtaceis: macula nigra, antennis breuibus. Ent. ſyſt. 2. 263. 44. * *2 maculatus.*

Habitat in India orientali. Muſ. Dom. Lund.

62. C. thorace ſubſpinoſo, rufus, elytris teſtaceis, baſi atro-cyaneis, antennis breuibus. Ent. ſyſt. 2. 263. 45. * *ſplendidus.*

Habitat Tranquebariae. Muſ. D. Lund.

63. C. thorace ſubſpinoſo, niger, antennis albidis, elytris teſtaceis, baſi apiceque cyaneis, nitidiſſimis. *Daldorfii.*

*Lamia Daldorfii.* Illig.
Habitat in Sumatra. Daldorff. Muſ. D. de Seheſtedt.
Statura et ſumma affinitas C. nitentis. Antennae villoſae, albidae: articulo primo ferrugineo, apice obſcurae, longitudine corporis. Caput nigrum, immaculatum. Thorax rotundatus, ſtriga media eleuata, niger. Elytra ſubſtriata, nitida, teſtacea, baſi parum, apice latius, cyanea, nitida. Corpus nigrum. Pedes teſtacei, femoribus nigris.

64. C. thorace ſubſpinoſo, niger, antennis pedibusque flaueſcentibus. Elytris teſtaceis, apice nigris. *ignitus*

Habitat in Sumatra. Dom. Daldorff.
Varietatem praecedentis credit Dom. Illiger ac forte rite. Antennae piloſae, flaueſcentes, longitu-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

gitudine fere corporis. Caput nigrum, la.. flaueſcente: faſcia media nigra. Thorax ſubſpinoſus, niger, ruga media eleuata. Elytra teſtacea, nitida, apice nigra. Corpus nigrum, pectore flaueſcente. Pedes obſcuri.

*fulgidus.* 65. C. thorace ſubſpinoſo, ferrugineus, elytris cyaneis, nitidiſſimis.

Habitat in Sumatra. Dom. Daldorff.

Statura omnino praecedentium. Antennae piloſae, flauae: articulo primo ferrugineo, apice acutae, longitudine corporis. Caput et thorax ferruginea, ruga thoracis media eleuata. Elytra cyanea, nitidiſſima. Corpus ferrugineum.

*longicornis* 66. C. thorace inermi: dorſo plano, corpore fuſco griſeoque vario, antennis longiſſimis. Ent. ſyſt. 2. 263. 46. *

Habitat in Coromandel. Muſ. D. Banks.

*marginalis* 67. C. thorace inermi, elytris ſubteſtaceis: margine omni nigro. Ent. ſyſt. 2. 264 47. *

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*breuicornis.* 68. C. thorace inermi, viridi, elytris obſcuris, antennis breuibus, nigris. Ent. ſyſt. 2. 264. 48. *

Oliv. Inf. 70. tab. 2. fig. 22.

Habitat in Sierra Leon Africae. Muſ. D. Banks.

*ſcriptus.* 69. C. thorace inermi, nigro; lineis duabus albis, elytris obſcure - cyaneis: lineolis faſciaque media abbreuiata, compreſſis, cinereis. Ent. ſyſt. ſuppl. 143. 49. *

Habitat in Isle de France. Muſ. Dom. Boſc.

*iuuencus.* 70. C. thorace inermi, rugoſo, elytris acuminatis, nigris lanuginoſo - caneſcentibus, antennis longiſſimis, Ent. ſyſt. 2. 264. 49. *

Linn. ſyſt. Nat. 2. 631. 53.

Habitat in America.

71.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

71. C. thorace inermi, rugoſo, griſeus, elytris vni- *holoſeri-*
dentatis, holoſericeis, fuſco cinereoque micantibus, antennis mediocribus. Ent. ſyſt. 2. 264. 50. * *ceus.*

Oliv. Inſ. 62. tab. 17. fig. 127.

Habitat in India orientali. Muſ. D. Lund.

72. C. thorace inermi, ſubrugoſo, cinereus, immaculatus, antennis longiſſimis. Ent. ſyſt. 2. 265. 31. * *cinereus.*

Habitat Tranquebariae. Muſ. D. Lund.

## 136. LAMIA.

*Palpi* filiformes.
*Maxilla* cornea, bifida.
*Ligula* cornea, bifida.
*Antennae* ſetaceae.

1. L. thorace acute ſpinoſo, rugoſo, elytris cinereis, *gigas.*
macula marginali nigra, baſi bituberculatis, ſcabris, antennis longis. Ent. ſyſt. 2. 265. 1. *

Oliv. Inſ. 67. tab. 23. fig. 91.

Habitat in Guinea. Muſ. Dom. Banks.

2. L. thorace quadriſpinoſo, ſcutello elytrisque ſpi- *tribulus.*
noſis, antennis longioribus. Ent. ſyſt. 2. 266. 2. *

Oliv. Inſ. 67. tab. 14. fig. 100.

Habitat ad fluuium Gaboon Africae. Muſ. D. Banks. *fronticor-*

3. L. thorace ſpinoſo, frontis cornu porrecto, apice *nis.*
emarginato incuruo, antennis longis. Ent. ſyſt. 2. 266. 3. *

---

*Lamia* corpus cylindricum, rotundatum, glabrum, immarginatum, tardum; *capite* magno, obtuſo, decliui, inter antennas canaliculato; *oculis* magnis, rotundatis, marginalibus, prominulis; *antennis* elongatis, approximatis, inter oculos inſertis; *thorace* cylindrico, ſpinoſo, inaequali; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* validis, curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cerambyx 2punctatus.* Drury Inſ. 2. tab. 31. fig. 2.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*pedicornis.* 4. L. thorace elytrisque ſpinoſis, femoribus antici cornutis. Ent. ſyſt. 2. 267. 4. *
Oliv. Inſ. 67. tab. 16. fig. 119.
Habitat in noua Hollandia. Muſ. D. Banks.

*hyſtrix.* 5. L. thorace quinqueſpinoſo, elytris faſciculato-pi loſis, antennis mediocribus, ſerratis. Ent. ſyſt. 2. 267. 5. *
Oliv. Inſ. 67. tab. 15. fig. 110.
Habitat ad Cap. Bon. Spei. Muſ. Dom. Banks.

*criſta.* 6. L. thorace acute ſpinoſo, griſea, elytris baſi tu berculo compreſſo, tridentato. Ent. ſyſt. 2. 268. 6. *
Oliv. Inſ. 67. tab. 14. fig. 101.
Habitat in noua Zelandia. Muſ. D. Banks.

*marmorata.* 7. L. thorace acute ſpinoſo, cinereo fuſcoque, varia elytris baſi tuberculatis, antennis longis. Ent. ſyſt. ſuppl. 143. 7. *
Habitat in Africa. Muſ. Dom. Boſc.

*ambulator* 8. L. thorace antice vtrinque biſpinoſo, corpore ne buloſo. Ent. ſyſt. 2. 268. 7. *
Petiv. Gazoph. tab. 37. fig. 6.
Habitat in Inſulis Luzonum. Muſ. Britann.

*ſpinipes.* 9. L. thorace vtrinque biſpinoſo, elytris punctatis variegatis, femoribus anticis baſi vniſpinoſis.
Habitat in Isle de France. Muſ. D. de Seheſted
Affinis videtur L. ambulatori. Antennae longitu
dine corporis nigrae, albo Caput villo
ſum, pallide cinereum tice biſpino
ſus: ſpina v fup eriore
re, acu var
tra que

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

baſeos mediaque albis, teſtaceo-variegatis. Pedes cinerei, anticis ſpina baſeos erecta, acuta.

10. L. thorace ſpinoſo, bimaculato, elytris baſi ſcabris, apiceque mucronatis. Ent. ſyſt. 2. 290. 89. * *Rubus.*

*Cerambyx Rubus.* Linn. ſyſt. Nat. 2. 625. 21.
Oliv. Inſ. 67. tab. 7. fig. 57.
Degeer Inſ. 5. 106. 7. tab. 13. fig. 16.
Habitat in India.

11. L. thorace ſpinoſo, bimaculato, elytris albo-maculatis, baſi ſcabris, mucronatis, apice bidentatis. Ent. ſyſt. 2. 290. 90. * *8maculata*

Habitat Tranquebariae. Muſ. D. Lund.

12. L. thorace ſpinoſo, rugoſo, elytris truncatis, griſeis, antennis compreſſis: articulis apice ſpinoſis. Ent. ſyſt. 2. 290. 91. * *ſpinicornis*

Oliv. Inſ. 67. tab. 5. fig. 33.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

13. L. thorace ſpinoſo, capiteque albo-lineatis, elytris albo-variis, antennis longis. *lineator.*

Oliv. Inſ. 67. tab. 19. fig. 143.
Habitat in India.
Magnitudo L. ſpinicornis. Caput nigrum, vertice lineis tribus albis inter antennas, in maculam coeuntibus. Thorax ſcaber, acute ſpinoſus, lineis tribus albis. Elytra baſi parum ſcabra, apice truncata, nigra, albo-nebuloſa.

14. L. thorace ſpinoſo, ferruginea, elytris maculis tribus niueis. *lactator.*

Habitat in Guinea. Dom. Meier. Muſ. D. de Seheſtedt.
Statura et magnitudo L. Seheſtedii. Antennae ferrugineae. Mandibulae apice atrae. Caput ferrugineum, macula frontali niuea. Thorax ſpi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſpinoſus, maculis duabus baſeos, niueis. Elytra laeuia, ferruginea, maculis tribus magnis, nitidis. Corpus ferrugineum, linea laterali lata, niuea, quae ab oculis vsque ad anum extenditur.

*globifera.* 15. L. thorace inermi, rugoſo, obſcura, elytris punctis eleuatis, globoſis, atris.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura L. ſcabratoris. Antennae corpore longiores, nigrae. Thorax rotundatus, muticus, rugoſus, tuberculis dorſalibus, plurimis, eleuatis, ſubſpinoſis. Elytra obſcura, punctis plurimis, eleuatis globoſis, atris. Corpus obſcurum.

*ſcabrator.* 16. L. thorace ſpinoſo, ſubteſtacea, elytris punctis baſeos atris, ſcabris. Ent. ſyſt. 2. 291. 92. *
Oliv. Inſ. 67. tab. 3. fig. 14.
Habitat in India orientali. Muſ. D. Banks.

*gladiator.* 17. L. thorace ſpinoſo, elytris teſtaceo cinereoque reticulatis, baſi punctis eleuatis, ſcabris.
*Lamia ſpinator.* Ent. ſyſt. ſuppl. 145. 92. *
Habitat in India orientali. D. Daldorff.

*Sebeſtedii.* 18. L. thorace ſpinoſo, elytrisque fuſcis: maculis holoſericeis, niueis. Ent. ſyſt. ſuppl. 146. 94.*
Habitat in India orientali. Dom. Daldorff.

*fuſcator.* 19. L. thorace biſpinoſo, tuberculatoque, elytris apice tridentatis, griſeis: faſcia cinerea. Ent. ſyſt. 2. 291. 94. *
Habitat Tranquebariae. Muſ. Dom. Lund.

*barbicornis.* 20. L. thorace ſpinoſo elytrisque fuſco nigroque variis albomaculatis: articulo tertio faſciculo pilorum. Ent. ſyſt. ſuppl. 145. 94. *
Habitat in Guinea. Muſ. D. de Seheſtedt.

*tridentata* 21. L. thorace ſpinoſo criſtaque duplici griſea, elytris apice tridentatis.

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Guinea. Dom. Heſſe. Muſ. D. Lund.
Statura L barbicornis. Antennae longitudine corporis piloſae, cinereae. Caput canaliculatum, cinereum. Thorax acute ſpinoſus, griſeus criſta duplici eleuata, inaequali, obtuſa, fuſca. Elytra punctata, griſea, fuſco-maculata, apice tridentata. Corpus griſeum.

22. L. thorace ſpinoſo, elytris conuexis, atris, antennis mediocribus. Ent. ſyſt. 2. 268. 8. * *textor.*
*Cerambyx textor.* Linn. ſyſt. Nat. 2. 629. 41.
Fn. Sv. 656.
Geoff. Inſ. 1. 201. 3.
Oliv. Inſ. 67. tab. 6. fig. 39.
Degeer Inſ. 5. 64. 3.
Panz. Fn. Germ. 19. tab. 1.
Schaeff. Icon. tab. 10. fig. 1.
Payk. Fn. Sv. 3. 61. 1.
Habitat in truncis arborum Europae.

23. L. thorace ſpinoſo, rugoſo, nigra, elytris emarginatis, ferrugineis, antennis mediocribus. Ent. ſyſt. 2. 268. 9. *vaginator.*
Habitat in India orientali. Muſ. Dom. Lund.

24. L. thorace ſpinoſo, nigra, elytris vitta ſemicirculari punctoque marginali cinereis, antennis longis. Ent. ſyſt. 2. 269. 10. * *crucifera.*
Oliv. Inſ. 67. tab. 10. fig. 68.
Habitat - - Muſ. Dom. Olivier.

25. L. thorace ſpinoſo, nigra, antice faſciis, poſtice punctis viridibus, elytris baſi mucronatis. Ent. ſyſt. 2. 269. 11. * *pulchra.*
*Cerambyx pulchra.* Drury Inſ. 1. tab. 32. fig. 6.
Oliv. Inſ. 67. tab. 1. fig. 6.
Habitat in Africa. Dom. Drury.

26.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*regalis.* 26. L. thorace ſpinoſo, viridi-faſciato, elytris viridi-irroratis, maculisque tribus fuluis. Ent. ſyſt. 2. 269. 12. *
Oliv. Inſ. 67. tab. 4. fig. 24. a.
Brown. Illuſtr. tab. 50. fig. 1.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

*imperialis.* 27. L. thorace ſpinoſo, ferrugineo, villoſa, viridi-faſciata, antennis longis, atris.
Oliv. Inſ. 67. tab. 4. fig. 24. c.
Habitat in Guinea. Dom. Meier. Muſ. D. de Seheſtedt.
Nimis affinis L. regali, cuius forte ſpecimen nuper declaratum. Antennae longae, nigrae. Caput villoſo-ferrugineum, obſcure lineatum orbitaque oculorum viridi. Thorax ſpinoſus, ſtriga media impreſſa faſciaque poſtica viridi. Elytra villoſo-ferruginea, faſciis duabus, anteriore recta, media e maculis tribus quadratis, concatenatis punctoque poſtico viridibus. Pedes ferruginei.

*vndata.* 28. L. thorace mutico, cinerea, thorace faſcia, elytris duabus vndatis, atris. Ent. ſyſt. 2. 270. 13. *
Oliv. Inſ. 67. tab. 4. fig. 26.
Habitat - - -

*irrorata.* 29. L. thorace ſpinoſo, fuſco, ferrugineoque, varia, elytris nigris, ferrugineo-irroratis, antennis mediocribus. Ent. ſyſt. 2. 270. 14. *
Voet. Coleopt. 2. tab. 7. fig 18.
Habitat in India orientali. Muſ. Dom. Lund.

*nigricornis.* 30. L. thorace mutico, griſea, elytris dorſo planis, antennis breuibus nigris. Ent. ſyſt. 2. 270. 15. *
Habitat in Coromandel. Muſ. D. Banks.

*bifidator.* 31. L. thorace ſpinoſo tuberculatoque cinereo: margine lineolisque atris, elytris cinereis: maculis oblongis, atris.
Habitat in America boreali. Muſ. Dom. Boſc.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo omnino L. aedilis. Antennae corpore duplo longiores, cinereae, articulis apice fuſcis. Caput nigrum. Thorax vtrinque ſpina valida tuberculisque dorſalibus; cinereus, margine ſub ſpina, lineolis quinque abbreuiatis: intermedia ſolitaria, atris. Elytra baſi punctata, cinerea: macula lineari baſeos, magna marginali, magnaque oblonga pone medium, antice bifida, poſtice puncto paruo albido. Corpus griſeum, femoribus clauatis.

32. L. thorace ſpinoſo: punctis quatuor luteis, elytris obtuſis, nebuloſis, antennis longis. Ent. ſyſt. 2. 270. 16. * *aedilis.*

*Cerambyx aedilis.* Linn. ſyſt. Nat. 2. 628. 37. Fn. Sv 653.
Oliv. Inſ. 67. tab. 9. fig. 59.
Degeer Inſ. 5. 66. 5. tab. 4. fig. 1. 2.
Friſh. Inſ. 13. tab. 12.
Sulz. Inſ. tab. 4. fig. 27.
Schaeff. Icon. tab. 14. fig. 7.
Payk. Fn. Sv. 3. 62. 3.
Habitat in Europae borealis truncis.

33. L. thorace ſubſpinoſo: vtrinque macula ocellari atra, elytris variegatis, antennis longis. *ocellator.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura L. aedilis, at duplo minor. Antennae longae, ferrugineae, articulis baſi pallidis. Thorax ſpina parua poſtica; obſcurus, antice lineolis duabus paruis, atris et ad latera macula magna atra, annulo albo cincta. Elytra punctata, variegata, pone medium angulo vno alteroue atro punctoque paruo niueo. Femora clauata.

34. L. thorace ſpinoſo tuberculatoque, cinerea, elytris fuſco-variis, punctis eleuatis, nigris, ſca- *atomaria.* IV. 61.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

scabris, antennis longis. Ent. syst. 2. 271. 17. *

Habitat in Germania. Dom. Smidt.

*varia.* 35. L. thorace spinoso tuberculatoque, corpore nigro cinereoque vario, femoribus clauatis, antennis mediocribus. Ent. syst. 2. 271. 18. *

Oliv. Inf. 67. tab. 3. fig. 16.

Roffi Fn. Etr. 1. 142. 360.

Habitat in Europa australiori. Dom. Zschuck.

*antennator* 36. L. thorace quadrispinoso, antennis longissimis, tenuibus, ferrugineis, albo-subannulatis.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Statura praecedentium, at parua. Antennae longissimae, tenues, ferrugineae, annulo vno alteroue albido. Caput fuscum. Thorax tuberculis quatuor, in strigam dispositis; fuscus, linea dorsali albida. Elytra variegata, lituris variis albis. Pedes variegati, femoribus clauatis.

*araneiformis.* 37. L. thorace spinoso tuberculatoque, elytris porosis; macula marginali fusca, antennis longis. Ent. syst. 2. 271. 19. *

*Cerambyx araneiformis.* Linn. syst. Nat. 2. 625. 22.

Oliv. Inf. 67. tab. 5. fig. 34.

Drury Inf. 2. tab. 35. fig. 4.

Sloan. Jam. 2. tab. 237. fig. 24.

Voet. Col. 2. tab. 9. fig. 33. 34.

Habitat in America meridionali.

*punctata.* 38. L. thorace spinoso, fusca, albo-punctata. Ent. syst. 2. 272. 20. *

Oliv. Inf. 67. tab. 6. fig. 42.

Habitat in Cayenna. Mus. Dom. Olivier.

39.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

39. L. thorace ſpinoſo, elytris cinereis, nigro-maculatis, antennis longiſſimis: articulo tertio apice nodoſo. Ent. ſyſt. 2. 272. 21. * *nodoſa.*
Oliv. Inſ. 67. tab. 14. fig. 103.
Habitat in Marylandia. Muſ. Britt.

40. L. thorace multidentato: dorſo plano, elytris tibiisque anticis vnidentatis. Ent. ſyſt. 2. 272. 22. * *cancriformis.*
*Cerambyx puſtulatus.* Drury Inſ. 2. t. 35. f. 1.
Habitat in Jamaica.

41. L. thorace ſpinoſo tuberculatoque, elytris punctatis ſpinoſisque, antennis longis. Ent. ſyſt. 2. 273. 23. * *tuberculata.*
Oliv. Inſ. 67. tab. 16. fig. 114.
Habitat in Jamaica. Muſ. Dr. Hunter.

42. L. thorace acute ſpinoſo, cinerea, elytris ſpinoſis, antennis longis. Ent. ſyſt. 2. 273. 25. * *horrida.*
*Cerambyx horridus.* Oliv. Inſ. 67. t. 4. f. 29.
Habitat in Cayenna, Muſ. D. Olivier.

43. L. thorace quadriſpinoſo, elytris tuberculato-granulatis, tibiis anticis apice dilatatis. Ent. ſyſt. 2. 273. 26. *Scorpio.*
Oliv. Inſ. 67. tab. 3. fig 19.
Habitat in America meridionali. Muſ. Dr. Hunter.

44. L. thorace ſubſpinoſo, flauo, lineato, elytris baſi ſcabris, brunneis, flauo-punctatis, apice fuſcis, rufo-maculatis. *decorator.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura praecedentium, at minor. Antennae vix longitudine corporis, atrae, articulis tribus prioribus longioribus, clauatis. Caput obſcurum, lineis vtrinque duabus, obliquis, ſub oculis. Thorax ſubſpinoſus, obſcurus: lineis tribus flauis,



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

lateralibus abbreuiatis. Elytra basi scabra, obscura: punctis sparsis, flauis, pone medium nigra: maculis sparsis rufis. Thorax niger, rufo-maculatus. Abdomen nigrum, lineis duabus punctorum alborum. Femora clauata. Tibiae ante apicem vnidentatae.

*mixta.* 45. L. thorace spinoso, cinereo fuscoque variegata, antennis longis. Ent. syst. suppl. 144. 26. *
Habitat in America boreali. Dom. v. Rohr.

*obscura.* 46. L. thorace spinoso, fusca, elytris striatis, apice pallidis, antennis longis. Ent. syst. suppl. 144. 26. *
Habitat in America boreali. Dom. v. Rohr.

*glauca.* 47. L. thorace quinquespinoso, glauca, elytris muricatis, latere fasciaque nigris, antennis longis. Ent. syst. 2. 274. 27. *
*Lamia Spengleri.* Ent. syst. 2. 291. 93. *
*Cerambyx glaucus.* Linn. syst. Nat. 2. 626. 28.
Degeer Inf. 5. 7. 13. tab. 14. fig. 4.
Habitat in America.

*bidentata.* 48. L. thorace subspinoso, elytris bidentatis, scabris cinereo-fuscis. Ent. syst. 2. 274. 28. *.
Voet. Coleopt. 2. tab. 15. fig. 62.
Habitat in America. Dom. v. Rohr.

*scabra.* 49. L. thorace spinoso tuberculatoque, elytris basi scabris, apice bidentatis, antennis longissimis. Ent. syst. 2. 274. 29. *
Oliv. Inf. 67. tab. 17. fig. 128.
Habitat - - Mus Britt.

*compressa.* 50. L. thorace spinoso, grisea, femoribus compressis clauatis, antennis longis.
Habitat in Guinea. Dom. Hesse. Mus. D. Lund.
Parua in hoc genere. Antennae corpore longiores, nigrae: articulis basi cinereis; primo clauato

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nato. Caput cinereo-villoſum, canaliculatum. Thorax acute ſpinoſus, cinereus, lineis duabus fuſcis. Elytra cinereo fuſcoque varia. Pedes breues, femoribus inermibus, clauatis: claua compreſſa.

51. L. thorace ſubſpinoſo, nigro, elytris variegatis, antennis mediocribus. *clauator.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Parua, Antennae corpore vix longiores, nigrae, articulis baſi albis. Caput et thorax nigra, punctata, nigro alboque varia. Pedes pallidi, femoribus clauatis.

52. L. thorace tuberculato, elytris punctatis, cinereis; apice bidentatis, fuſcis, antennis longis. Ent. ſyſt. 2. 275. 30. * *praemorſa.*

Habitat in Americae Inſulis. Dom. Badier.

53. L. thorace bituberculato, elytris baſi tuberculo compreſſo, antennis breuibus baſi vnidentatis. *inaequalis.*

Habitat Tranquebariae. Dom. Daldorff. Muſ. D. de Seheſtedt.

Minuta in hoc genere. Antennae breues, nigrae, baſi dente eleuato, acuto. Thorax inermis, rotundatus, niger: dorſo albido; tuberculis duobus dorſalibus, compreſſis. Elytra punctata, baſi tuberculo compreſſo, quod in lineam eleuatam excurrit. Abdomen nigrum, ſegmentorum marginibus albis.

54. L. thorace biſpinoſo tuberculatoque griſea, elytris tuberculis duobus eleuatis, acutis. *obſcurator*

Habitat in Guinea. Dom. Meier. Muſ. D. Lund.

Minor L. griſatore. Antennae vix longitudine corporis. Thorax biſpinoſus, punctato-tuberculatus. Elytra griſea, baſi tuberculis duobus eleuatis, acutis, nigris, vno pone alterum.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſpinifera.* 55. L. thorace ſpinoſo, cinereo, villoſa, elytris bidentatis, teſtaceis, antennis longis. Ent. ſyſt. 2. 275. 31.*

Habitat in America meridionali. Dr. Pflug.

*griſator.* 56. L. thorace mutico, ſubſtriato, griſea, elytris ſubſpinoſis, baſi tuberculis duobus, antennis breuibus, villoſis. Ent. ſyſt. 2. 275. 32. *

Habitat Tranquebariae. Dom. de Schlanbuſch.

Elytra griſea, lineis ſpinarum breuiſſimarum, baſi tuberculis duobus compreſſis, apice litura atra.

*cylindrator.* 57. L. thorace mutico, villoſo, griſea, elytris ſubſpinoſis, baſi tuberculis duobus, apice obſcuris.

Habitat in Guinea. Dom. Meier. Muſ. D. de Seheſtedt.

Statura et ſumma affinitas L. griſatoris. Antennae breues, villoſae, nigrae, albo-punctatae. Caput griſeum. Thorax inermis, rotundatus, villoſus, cinereus, immaculatus. Elytra griſea, lineis e ſpinis eleuatis, acutis, baſique tuberculis duobus paruis, eleuatis, apice obſcura: litura alba.

*aſpinator.* 58. L. thorace ſpinoſo, griſea, elytris obſcuris, baſi ſcabris, apice bidentatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura L. griſator. Antennae corpore vix longiores, glabrae, baſi denticulo paruo, acuto. Caput griſeum, macula magna, verticali, ferrugínea. Thorax ſpinoſus, griſeus: linea dorſali obſcuriore. Elytra ſcabra, obſcura, apice cinereo atroque varia, terminantur dentibus duobus validis, obtuſis.

*aedificator* 59. L. thorace ſpinoſo tuberculatoque, griſea, elytris baſi bituberculatis, antennis mediocribus. Ent. ſyſt. 2. 275. 33. *

Habitat in India orientali. Muſ. D. Lund.

60.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

60. L. thorace ſpinoſo, elytris cinereis, nigro-irroratis maculisque numeroſis teſtaceis. Ent. ſyſt. 2. 276. 34. * *amputator*

Habitat in Americae meridionalis Inſulis; ramos etiam craſſiores abſcindere dicitur. Muſ. Dom. Banks.

61. L. thorace ſpinoſo, elytris baſi poroſis, antennis mediocribus. Ent. ſyſt. 2. 276. 35. * *ſternutator*

Habitat in Inſula Barbados. Dom. Drury.

62. L. thorace ſpinoſo, elytrisque baſi daedaleis apice griſeis, antennis mediocribus. Ent. ſyſt. 2. 276. 36. * *ſpinator.*

Habitat ad Cap. Bon. Spei. Muſ. Dom. Banks.

*Variat* ex Guinea elytris magis griſeis: maculis nigris nunc rarioribus, nunc crebrioribus.

63. L. thorace ſpinoſo, elytris teſtaceis, albo-irroratis: baſi punctis eleuatis, glabris. Ent. ſyſt. 2. 276. 37. * *gutturator*

Oliv. Inſ. 67. tab. 13. fig. 92.

Habitat in Africa. Muſ. Britt.

64. L. thorace inermi, ferrugineo, nigro, lineato, elytris fuſco ferrugineoque variis: macula marginali cinerea, antennis mediocribus. Ent. ſyſt. 2. 277. 38. * *nebuloſa.*

Oliv. Inſ. 67. tab. 3. fig. 15.

Schaeff. Icon. tab. 55. fig. 4.

Habitat in Germania. Muſ. D. Banks.

65. L. thorace inermi, cinerea, elytris macula apicis fuſca, antennis longis. *inermis.*

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

L. nebuloſa minor. Antennae corpore longiores, cinereae, articulis apice fuſcis. Thorax rotundatus, inermis, cinereus. Elytra cinerea, punctis

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ctis aliquot eleuatis indiſtinctis, macula magna poſtica, fuſca. Corpus cinereum, femoribus clauatis.

*repandator* 66. L. thorace ſpinoſo, obſcura, elytris faſcia poſtica repanda, albida: baſi punctis eleuatis, atris. Ent. ſyſt. 2. 277. 39. *
Habitat in Cayenna. Dom. v. Rohr.

*variolator.* 67. L. thorace ſpinoſo, lineato, elytris fuſcis: linea ſemicirculari, alba. Ent. ſyſt. 2. 277. 40. *
Habitat in Indiis.

*ſutor.* 68. L. thorace ſpinoſo, elytris atris, ferrugineo-maculatis, ſcutello luteo, antennis longiſſimis. Ent. ſyſt. 2. 277. 41. *
*Cerambyx ſutor.* Linn. ſyſt. Nat. 2. 628. 38.
Fn. Sv. 655.
Oliv. Inſ. 67. tab. 3. fig. 20.
Schaeff. Icon. tab. 65. fig. 1.
Degeer Inſ. 5. 65. 4.
Panz. Fn. Germ. 19. tab. 2.
Sulz. Inſ. tab. 5. fig. 4.
Payk. Fn. Sv. 3. 62. 2.
Habitat in Europae ſyluis.

*ſartor.* 69. L. thorace ſpinoſo, niger, ſcutello flauo, elytris immaculatis, antennis longiſſimis. Ent. ſyſt. 2. 278. 42. *
Panz. Fn. Germ. 19. tab. 3.
Habitat Dresdae in Aeſculo Hippocaſtano. Dom. Zſchuck.

*dentator.* 70. L. thorace ſpinoſo, fuſco cinereoque varia. Ent. ſyſt. 2. 278. 43. *
Habitat in Carolina.

*ruſticator.* 71. L. thorace acute ſpinoſo, cinereo, ſcutello albo, elytris punctatis, antennis longis.

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Sumatra. D. Daldorff. Muſ. D. Lund.
Minor L. ſutore. Antennae corpore parum longiores, cinereae, immaculatae. Caput, thorax, elytra ſubholoſericea, cinerea. Scutellum album. Elytra punctata. Corpus cinereum.

72. L. thorace ſpinoſo, nigra, thorace elytrisque fuluis, thorace nigro-lineato, elytris reticulatis. Ent. ſyſt. 2. 278. 44. * — *reticulator*
Oliv. Inſ. 67. tab. 12. fig. 83.
Habitat in India orientali. Muſ. D. Banks.

73. L. thorace acute ſpinoſo, atra, elytris ſtrigis numerosis interruptis albis, antennis mediocribus. Ent. ſyſt. 2. 278. 45. * — *ſcalator.*
Habitat in ~~Luſitania.~~ Muſ. D. Boſc.
Louisiana

74. L. thorace ſubſpinoſo, obſcure cinerea, fuſco-maculata, elytris apice rufis. Ent. ſyſt. 2. 279. 46. * — *molator.*
Oliv. Inſ. 67. tab. 14. fig. 99.
Habitat in Cayenna. Muſ. Dom. Hunter.

75. L. thorace ſpinoſo, elytris cinereo, fuſco glaucoque nebuloſis, antennis longiſſimis rufis. Ent. ſyſt. 2. 279. 47. * — *titillator.*
Oliv. Inſ. 61. tab. 15. fig. 109.
Habitat in Carolina. Muſ. Britt.

76. L. thorace ſpinoſo, elytris cinereo-puluerulentis: vittis duabus flauis, nigro-marginatis. Ent. ſyſt. 2. 279. 48. * — *vittator.*
*Cerambyx incanus.* Forſt. nou. ſp. Cent. 1. 38.
Oliv. Inſ. 67. tab. 15. fig. 104.
Petiv. Gazoph. tab. 24. fig. 6.
Habitat ad littora ſinus Campechiani. Muſ. Britt.

77. L. thorace ſpinoſo, nigra, elytris maculis quatuor flauis, ſubocellaribus, antennis longis. Ent. ſyſt. 2. 280. 49. * — *oculator.*

Ce-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cerambyx oculatus.* Degeer Inſ. 7. 654. t. 49. f. 1.
Oliv. Inſ. 67. tab. 2. fig. 2.
Voet. Col. 2. tab. 7. fig. 21.
Habitat ad Cap. Bon. Spei.

*Capenſis.* 78. L. thorace biſpinoſo, nigra, elytris faſciis quatuor rufis, antennis mediocribus. Ent. ſyſt. 2. 280. 50.
*Cerambyx Capenſis.* Linn. ſyſt. Nat. 2. 628. 32.
Oliv. Inſ. 67. tab. 8. fig. 51.
Drury Inſ. 1. tab. 39. fig. 3.
Petiv. Gazoph. tab. 3. fig. 8.
Voet. ſcarab. 2. tab. 8. fig. 25.
Habitat ad Cap Bon. Spei.

*hottentotta.* 79. L. thorace ſpinoſo rufo, elytris nigris: faſcia maculisque duabus marginalibus ſanguineis. Ent. ſyſt. 2. 280. 51. *
*Cerambyx hottentotta.* Brown Illuſtr. 118. tab. 49 fig. 1.
Oliv. Inſ. 67. tab. 4. fig. 27.
Voet. ſcarab. 2. tab. 8. fig. 26.
Habitat ad Cap. Bon. Spei. Dom. Yeats.

*2 cincta.* 80. L. thorace ſpinoſo, nigra, elytris faſciis duabus continuis rufis. Ent. ſyſt. ſuppl. 146. 51.
Habitat in India orientali. D. Daldorff.

*ferrugator* 81. L. thorace ſpinoſo, obſcure ferruginea, elytris obſcure vireſcentibus: ſtriga interrupta ſanguinea, antennis longis. Ent. ſyſt. 2. 280. 52. *
Habitat ad Cap. Bon. Spei. D. Spengler.

*humeralis.* 82. L. thorace ſpinoſo, elytris flauis nigro-faſciatis antice mucronatis. Ent. ſyſt. 2. 281. 53. *
Habitat - - D. Drury.

83.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

83. L. thorace biſpinoſo, nigra, elytris faſciis quinque rubris, antennis mediocribus. Ent. ſyſt. 2. 281. 54. * *5faſciata.*

Habitat in Guinea. D. Iſert.

84. L. thorace ſubſpinoſo, atra, elytris faſciis tribus flauis, antennis mediocribus. Ent. ſyſt. 2. 281. 55. * *3faſciata.*

Oliv. Inſ. 67. tab. 16. fig. 121.
Herbſt. Arch. tab. 45. fig. 10.
Habitat in Sierra Leon Africae. Dr. Solander.

85. L. thorace ſpinoſo, poſtice ſinuato, elytris flauis: faſciis duabus maculaque rhombea cyaneis. Ent. ſyſt. 2. 281. 55. * *2faſciata.*

Habitat in Iamaica. Muſ. Brittann.

86. L. thorace ſpinoſo atro: margine flauo, elytris atris: faſciis tribus flauis, punctisque duobus albis. Ent. ſyſt. 2. 282. 57. * *nobilis.*

Habitat in Cayenna. Muſ. Dr. Hunter.

87. L. thorace ſpinoſo, atra, elytris faſciis duabus punctoque apicis luteis. antennis mediocribus. Ent. ſyſt. 2. 282. 58. * *Aethiops.*

Oliv. Inſ. 67. tab. 1. fig. 2.
Habitat ad Cap. Bon. Spei. Muſ. Brittann.

88. L. thorace ſpinoſo, cinerea, thorace elytrisque fuſco variis, antennis longis. Ent. ſyſt. 2. 282. 59. * *variegator*

Oliv. Inſ. 67. tab. 16. fig. 112.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

89. L. thorace mutico, fuſca, thorace coleoptrisque ocellis quatuor atris. Ent. ſyſt. 2. 282. 60. * *curculionoides.*

*Cerambyx curculionoides.* Linn. ſyſt. Nat. 2. 634. 48.
Oliv. Inſ. 67. tab. 10. fig. 69.

Geoff.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 210. 5.
Schaeff. Icon. tab. 39. fig. 1.
Payk. Fn. Sv. 3. 63. 4.
Habitat in Europa auſtraliori.

*luſca.* 90. L. thorace ſpinoſo, rugoſo, nigra, ferrugineo irrorata, elytris baſi macula ocellari atra, antennis longiſſimis. Ent. ſyſt. 2. 283. 61. *
Habitat in Siam. Muſ. D. Banks.

*notata.* 91. L. thorace ſpinoſo, obſcura, elytris punctis duobus approximatis atris. Ent. ſyſt. 2. 283. 62. *
Habitat in Cayenna. D. v. Rohr.

*lugubris.* 92. L. thorace acute ſpinoſo, nigra, elytris ſcabris, ferrugineo obſolete punctatis, antennis longiſſimis. Ent. ſyſt. 2. 283. 63. *
Habitat in Italia. D. de Slanbuſch.

*triſtis.* 93. L. thorace ſpinoſo, fuſca, elytris ſcabris: maculis duabus atris, antennis medioeribus. Ent. ſyſt. 2. 283. 64. *
*Cerambyx triſtis.* Linn. ſyſt. Nat. 2. 629. 42.
Oliv. Inſ. 67. tab. 9. fig. 62.
Herbſt. Arch. tab. 25. fig. 7.
Habitat in ligno Cupreſſi Galliae auſtralioris.
*Variat* pro ſexu, antennis longitudine corporis et duplo longioribus.

*funeſta.* 94. L. thorace ſpinoſo, fuſca, elytris laeuibus: maculis duabus atris, antennis breuibus. Ent. ſyſt. 2. 285. 65. *
Oliv. Inſ. 67. tab. 9. fig. 63.
Panz. Naturf. 24. 26. 36. tab. 1. fig. 36.
Roſſ. Fn. Etr. 1. 145. 366.
Habitat in Galliae Sambuco. D. Brouſſonet.

*punctator.* 95. L. thorace ſpinoſo, atra, elytris albo-punctatis, antennis longis. Ent. ſyſt. 2. 285. 66. *

Ce-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cerambyx punctatus.* Oliv. Inſ. 67. tab. 7. f. 49. ſ. v.
*Cerambyx farinoſus.* Drury Inſ. 2. tab. 31. f. 4.
*Cerambyx Chinenſis.* Forſt. nou. Inſ. ſpec. Cent. 1. 39.
Habitat in China. Dr. Fothergill.

96. L. thorace ſpinoſo, elytris lineis tribus eleuatis ſerrulatis, corpore cinereo, antennis mediocribus. Ent. ſyſt. 2. 284. 67. * *criſpa.*
Oliv. Inſ. 67. tab. 3. fig. 18.
Habitat ad Cap: Bon. Spei. Muſ. D. Lee.

97. L. thorace ſpinoſo hirtoque, elytris punctis piloſo-criſtatis. Ent. ſyſt. 2. 284. 68. * *faſciculata*
Oliv. Inſ. 67. tab. 17. fig. 131.
Habitat in Cayenna. Muſ. D. Olivier.
Elytra punctata, nigro ferrugineoque varia, pilis rarioribus, longioribus, albis.

98. L. thorace ſpinoſo, cinerea, elytris ſubfaſciatis, antennis mediocribus. Ent. ſyſt. 2. 285. 69. * *rotator.*
Oliv. Inſ. 67. tab. 17. fig. 126.
Habitat in India. Muſ. D. Banks.

99. L. thorace ſpinoſo, nigra, elytris bicarinatis albo-lineatis, pedibus ferrugineis, antennis breuibus. Ent. ſyſt. 2. 285. 70. * *glycyrrhizae.*
*Cerambyx ouatus.* Sulz. Hiſt. Inſ. tab. 5. fig. 9.
Oliv. Inſ. 67. tab. 16. fig. 115.
Pallas Iter 2. 723. 60.
Habitat in Sibiria. Muſ. D. Banks.

100. L. thorace ſpinoſo, nigra, coleoptris cruce media alba. Ent. ſyſt. 2. 285. 71. * *cruciata.*
Pall. Icon. tab. F. fig. 5.
Habitat in Ruſſia auſtraliori. D. Boeber.

101. L. thorace ſubſpinoſo, nigra, elytris cinerascentibus, antennis breuibus. Ent. ſyſt. 2. 286. 72. Linn. *fulginator*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. syst. Nat. 2. 539. 43.
Geoff. Inf. 1. 205. 8.
Oliv. Inf. 67. tab. 4. fig. 21.
Frisch. Inf. 13. tab. 19.
Degeer Inf. 5. 70. 7.
Voet. scarab. tab. 8. fig. 30.
Habitat in Europa australiori.
*Variat* aetate tota nigra.

*cineraria.* 102. L. thorace spinoso, cinerascens, antennis breuibus. Ent. syst. 2. 286. 73. *
Pall. Icon. tab. F. fig. 11.
Habitat in Russia meridionali. Dom. Hybner.

*carinata.* 103. L. thorace spinoso, nigra, elytris piceis: carina laterali eleuata albida, antennis breuibus. Ent. syst. 2. 286. 74. *
Habitat in Sibiria. Mus. D. Banks.

*annulata.* 104. L. thorace spinoso ferrugineo, fusca, antennis albo-annulatis.
Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.
Minuta in hoc genere. Antennae longitudine corporis, fuscae: articulo secundo toto, tertio, quarto, quinto sextoque basi albis. Corpus totum ferrugineo fuscoque varium.

*coquus.* 105. L. thorace subspinoso, hirsuta, elytris obtusis sulcatis atris antice ferrugineis, antennis mediocribus. Ent. syst. 2. 286. 75. *
*Cerambyx coquus.* Linn. syst. Nat. 2. 630. 44.
Habitat in America boreali.

*ruspator.* 106. L. thorace spinoso fusco: maculis duabus ferrugineis, elytris fuscis cinereo-variegatis. Ent. syst. 2. 286. 76. *
Oliv. Inf. 67. tab. 17. fig. 129.
Habitat in Africa aequinoctiali. Mus. D. Banks.

107.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

107. L. thorace ſpinoſo quadripunctato, elytris ruſis: maculis quatuor nigris, antennis breuibus. Ent. ſyſt. 2. 287. 77. * *tornator.*
*Cerambyx tetrophthalmus.* Forſt. Cent. 1. 41.
Oliv. Inſ. 67. tab. 8. fig. 52.
Herbſt. Arch. tab. 32. fig. 4.
Habitat in America boreali.

108. L. thorace ſpinoſo, griſea, elytris ferrugineis irroratis faſciisque duabus cinereis. Ent. ſyſt. 2. 287. 78. * *Bankii.*
Oliv. Inſ. 67. tab. 15. fig. 111.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

109. L. thorace inermi, griſea, elytris faſciis duabus macularibus abbreuiatis ſtrigaque poſtica punctata niueis, antennis breuibus. Ent. ſyſt. 2. 287. 79. * *ſaltator.*
Habitat - - -

110. L. thorace inermi, griſea, elytris maculis duabus niueis, antennis breuibus. *ſtigma.*
Habitat in Inſulis maris pacifici. D. Billardiere.
Parua. Antennae corpore breuiores, cum capite griſeae. Thorax omnino inermis, griſeus, litura dorſali niueus. Elytra ſtriato-punctata, griſea, maculis duabus magnis niueis. Corpus griſeum.

111. L. thorace inermi, griſea: ſtrigis tribus maculaque punctata marginali, albis, antennis breuibus. ~~*varia.*~~ *erector.*
Habitat in Auſtralaſiae Inſulis. D. Billardiere.
Statura praecedentis. Antennae griſeae, corpore breuiores. Caput canaliculatum, griſeum. Thorax rotundatus, inermis, griſeus, ſtrigis aliquot obſcurioribus, lineolaque vtrinque obliqua, alba. Elytra punctato-ſtriata, griſea, ante medium ſtri-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſtrigis duabus, margine fere coëuntibus, pone medium macula magna, marginali, e punctis plurimis conflata, ſtrigaque poſtica albis. Corpus griſeum.

*hiſtrio.* 112. L. thorace inermi, flaueſcens: punctis numeroſis niueis ſparſis, antennis breuibus. Ent. ſyſt. 2. 288. 80. *

Habitat Tranquebariae. Muſ. D. Lund.

*villoſa.* 113. L. thorace ſubſpinoſo, villoſa, teſtacea, nitida elytris baſi cyaneis, antennis mediocribus.

Habitat in Auſtralaſiae Inſulis. D. Billardiere.

Magnitudo L. pedeſtris. Antennae villoſae, teſtaceae, longitudine corporis. Thorax villoſus, ſubſpinoſus, teſtaceus, immaculatus. Elytra villoſa, teſtacea, nitida, baſi cyanea, ſutura tamen teſtacea. Corpus teſtaceum.

*teſtator.* 114. L. thorace ſpinoſo cornuque breui recumbente, corpore teſtaceo, antennis breuibus apice nigris. Ent. ſyſt. 2. 288. 81.

Oliv. Inſ. 67. tab. 16. fig. 122.

Habitat in Africa aequinoctiali. Muſ. D. Banks.

*pedeſtris.* 115. L. thorace ſpinoſo, atra: vitta integra alba, antennis mediocribus. Ent. ſyſt. 2. 288. 82. *

Linn. ſyſt. Nat. 2. 631. 51.

~~Oliv. Inſ. 67. tab.~~ 4. fig. 23.

Panz. Fn. Germ. 66. tab. 9.

Roſſ. Fn. Etr. 1. 146. 368.

Habitat in Europa auſtraliori.

*rufipes.* 116. L. thorace ſpinoſo, nigra, elytrorum ſutura alba, antennarum baſi pedibusque rufis. Ent. ſyſt. 2. 288. 83. *

Habitat in Hungaria. Dom. Hybner.

*morio.* 117. L. thorace ſpinoſo, nigra, elytris vnicoloribus, antennis breuibus. Ent. ſyſt. 2. 288. 84.

Oliv.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 67. tab. 10. fig. 67.
Herbſt. Arch. tab. 25. fig. 10.
Scopoli Ent. carn. 169.
Habitat in Europa auſtraliori. Dom. Zſchuck.

118. L. thorace ſpinoſo, atra, elytris marginibus lineisque duabus vtrinque coëuntibus albis. Ent. ſyſt. 2. 289. 85. * *lineata.*
*Cerambyx Scopoli.* Herbſt. Arch. t. 25. f. 11.
Habitat in Germania. Dom. Schneider.

119. L. thorace ſpinoſo, atra, elytris fuſcis: margine vittaque abbreuiata, albis. *vittigera.*
*Lamia vittigera.* Panz. Fn. Germ. . tab. .
Habitat in Germania auſtraliori.
Nimis L. lineatae affinis. Caput, antennae, thorax, corpus, pedes atra, immaculata. Elytra minus atra, ſiue fuſca margine vel potius vitta ante marginem aliaque media abbreuiata albis. Sutura albicat.

120. L. rhorace ſpinoſo, fuſca: vittis tribus integris albis, antennis mediocribus. Ent. ſyſt. 2. 289. 86. * *molitor.*
Oliv. Inſ. 67. tab. 12. fig. 84.
Roff. Fn. Etr. 1. 147. 369.
Habitat in India. Muſ. Britt.

121. L. thorace mutico, griſea, coleoptris baſi bituberculatis maculisque duabus communibus, albis, antennis breuibus. Ent. ſyſt. 2. 289. 87. * *tuberculator.*
Oliv. Inſ. 67. tab. 16. fig. 120.
Habitat ad Cap. Bon. Spei. Muſ. Dom. Banks.

122. L. thorace ſubſpinoſo, nigra, elytris ſtriga obliqua, alba. *obliquator.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo L. tuberculatoris. Antennae longitudine corporis, nigrae, articulis basi pallidis. Caput et thorax fusca. Elytra punctata humero prominulo, nigra, striga media, obliqua, albida. Femora subclauata.

*mutator.* 123. L. thorace mutico, lineato, cinereo, villosa, antennis pedibusque testaceis. Ent. syst. 2. 290. 88. *

Oliv. Ins. 67. tab. 14. fig. 102.

Habitat in Otaheite. Mus. D. Banks.

*bidens.* 124. L. thorace acute spinoso, grisea, elytris apice bidentatis. Ent. syst. 2. 291. 95. *

Habitat in noua Hollandia. Mus. D. Banks.

*Cantator.* 125. L. thorace mutico, cinereo, nigro-punctato, elytris pallide testaceis, apice bidentatis, cinereis nigro-maculatis. Ent. syst. 2. 291. 96. *

Habitat in China. D. de Slanbusch.

*Solandri.* 126. L. thorace subspinoso, nigra, elytris bidentatis, fuscis cinereo-irroratis. Ent. syst. 2. 292. 97. *

Oliv. Ins. 67. tab. 16. fig. 118.

Habitat in noua Hollandia. Mus. D. Banks.

*cornutor.* 127. L. thorace obtuse spinoso, mandibulis basi cornutis, antennis longissimis. Ent. syst. 2. 292. 98. *

Oliv. Ins. 67. tab. 17. fig. 132.

Habitat in America. Mus. Brittann.

*leprosa.* 128. L. thorace spinoso, elytris basi variolosis, cinereis: macula laterali nigra, antennis longis. Ent. syst. 2. 292. 99. *

Habitat in America. D. Drury.

137. STE-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

57. STENOCORVS. *Palpi* quatuor:
*anteriores* filiformes;
*posteriores* clauati.
*Antennae* filiformes.

St. thorace subspinoso, cyaneus, elytris basi flauis. Ent. syst. 2. 293. 1. ♀ — *cyaneus.*
*Cerambyx palliatus.* Forst. nou. Inf. spec. Cent. 1. 40.
Habitat in India.

St. thorace spinoso, elytris bidentatis griseis: macula flaua, antennis longissimis. Ent. syst. 2. 293. 3. ♂ — *gargani-cus.*
Oliv. Inf. 67. tab. 13. fig. 105.
Drury Inf. 1. tab. 37. fig. 6.
Degeer Inf. 5. tab. 14. fig. 3.
Habitat in Marylandia. Mus. Brittan.

St. thorace vtrinque bidentato, elytris bidentatis viridibus: linea laterali lutea. Ent. syst. 2. 293. 4. ♀ — *festiuus.*
*Cerambyx festiuus.* Linn. syst. Nat. 2. 623. 11.
Oliv. Inf. 67. tab. 16. fig. 113.
Drury Inf. 1. tab. 37. fig. 5.
Voet. Coleopt. 2. tab. 16. fig. 66.
Herbst. Arch. tab. 25. fig. 12.
Sulz. Hist. Inf. tab. 5. fig. 6.
Degeer Inf. 5. 100. 4. tab. 13. fig. 14.
Habitat in America.

4.

---

Stenocori corpus subcylindricum, glabrum, immarginatum, tardum; *capite* rotundato; *oculis* magnis reniformibus, marginalibus; *antennis* elongatis: articulis apice saepe spinosis in serie oculorum insertis; *thorace* rotundato; *scutello* orbiculato; *elytris* rigidis, fornicatis, longitudine abdominis, apice saepius bidentatis; *pedibus* cursoriis; *tarsis* quadriarticulatis; *colore* vario.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*marylandius.* 4. St. thorace depreſſo, tuberculato, mutico, elytris bidentatis, fuſco cinereoque nebuloſis, antennis mediocribus Ent. ſyſt. 2. 294. 5.
*Callidium marylandicum.* Oliv. Inſ. 70. tab. 1 fig. 5.
Habitat in Marylandia. Muſ. Britt.

*ſpinicornis* 5. St. thorace inermi, tuberculato, elytris bidentati antennarum articulis biſpinoſis, corpore variegato. Ent. ſyſt. 2. 294. 6. *
*Cerambyx ſpinicornis.* Oliv. Inſ. 67. tab. 1 fig. 130.
Drury Inſ. 1. tab. 41. fig. 4.
Voet. Coleopt. 2. tab. 24. fig. 131.
Habitat in America.

*pictus.* 6. St. thorace mutico, atro, nineo, lineato, elytris bidentatis, atris, cyaneo-variegatis, niueoque maculatis.
Habitat in Sumatra. Dom. Daldorff.
Magnus in hoc genere. Antennae breues. fuſcae. Caput atrum, niueo-lineatum. Thorax rotundatus, muticus, ater: lineis tribus niueis, lateribusque cyaneis. Scutellum niueum. Elytra atra, cyaneo-lineata, maculisque variis, niueis apice bidentata. Corpus niueum, cyaneo-nitidum. Pedes cyanei.

*bidens.* 7. St. thorace inermi, ſubtuberculato, antennarum articulis biſpinoſis, corpore teſtaceo. Ent. ſyſt. 2. 294. 7. *
*Cerambyx bidens.* Oliv. Inſ. 67. tab. 7. fig. 12
Habitat in America meridionali. Dom. de Paykul

*ſemipunctatus.* 8. St. thorace ſpinoſo, inaequali, elytris bidentatis baſi punctatis, apice glabris. Ent. ſyſt. 2. 29 8. *

Oli

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 67. tab. 2. fig. 19.
Habitat in noua Hollandia. Muſ. D. Banks.

9. St. thorace mutico, inaequali, elytris apice biden- *irroratus.*
tatis, albo-irroratis, antennis longis, aculeatis.
Ent. ſyſt. 2. 295. 9. *
*Cerambyx irroratus.* Linn. ſyſt. Nat. 2. 633. 62.
Oliv. Inſ. 67. tab. 19. fig. 145.
Drury Inſ. 1. tab. 41. fig. 3.
Habitat in America.

10. St. thorace ſpinoſo, obſcurus, elytris villoſis, *obſcurus.*
punctato-ſtriatis, apice vniſpinoſis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Paullo minor St. irrorato. Antennae longitudine corporis, cinereae, articulis apice nigris. Caput obſcurum, cinereo-villoſum. Thorax punctatus, niger, lineis villoſo-cinereis. Elytra villoſa, punctato-ſtriata, nigra, apice ſpina vnica, valida. Pedes ferruginei.

11. St. thorace mutico, cinereo: linea glabra, fuſca, *glabratus.*
elytris bidentatis. Ent. ſyſt. 2. 295. 10. *
Habitat in Americae meridionalis Inſulis. Dom. v. Rohr.

12. St. thorace ſpinoſo, piceus, elytris punctis ſpar- *farinoſus.*
ſis farinoſis, antennis longis. Ent. ſyſt. 2. 295. 11. *
*Cerambyx farinoſus.* Linn. ſyſt. Nat. 2. 626. 24.
Oliv. Inſ. 67. tab. 7. fig. 46.
Degeer Inſ. 5. 108. 9. tab. 13. fig. 17.
Habitat in radice Argemones mexicanae.

13. St. thorace ſpinoſo, ferrugineo, elytris acumina- *6 maculatus.*
tis: maculis tribus flauis. Ent. ſyſt. 2. 295. 11. *



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cerambyx 6maculatus.* Oliv. Inſ. 67. tab. 15. fig. 108.

Habitat in Cayenna. Muſ. Dom. Olivier.

*4dens.* 14. St. thorace quadriſpinoſo, elytris baſi punctatis atris: maculis tribus ferrugineis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura omnino et magnitudo St. 6maculati. Antennae corpore longiores, ferrugineae, pube aurea micantes. Thorax ater, ſpina laterali duabusque dorſalibus eleuatis, validis, acutis, armatus. Scutellum pube aurea tectum. Elytra atra, baſi punctata: macula glabra, ferruginea, apice glabra, maculis duabus approximatis, ferrugineis; exteriore longiore, terminata ſpina elongata, valida, acuta.

*5maculatus.* 15. St. thorace ſubſpinoſo, rufo: maculis quatuor flauis, elytris bidentatis, rufis: punctatis duobus lineaque apicis albis, antennis longiſſimis. Ent. ſyſt. 2. 296. 13. *

Habitat in Inſula Guadeloupa. Dom. Iſert.

*4maculatus.* 16. St. thorace ſpinoſo, ſcabro, elytris bidentatis: maculis binis glabris. Ent. ſyſt. 2. 296. 14. *

*Cerambyx 4maculatus.* Linn. ſyſt. Nat. 2. 627. 27.

Sloan Jamaic. 2. tab. 237. fig. 3.

Drury Inſ. 1. tab. 37. fig. 3.

Voet. Coleopt. 2. tab. 15. fig. 65.

Habitat in America meridionali.

*maculoſus.* 17. St. thorace ſpinoſo, fuſcus, elytris bidentatis: maculis binis glabris, antennis mediocribus, nigris. Ent. ſyſt. 2. 296. 15. *

*Cerambyx maculoſus.* Oliv. Inſ. 67. t. . f. .

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. Muſ. Dom. Olivier.

8. St. thorace mutico, nigro: macula vtrinque rufa, elytris nigris: maculis binis glabris, antennis longiſſimis. Ent. ſyſt. 2. 297. 16. ♀ *geminatus.*

Habitat in Sierra Leon Africae. Dr. Pflug.

9. St. thorace mutico, cylindrico, cinereo, elytris ferrugineis: maculis duabus flaueſcentibus, obſoletis. Ent. ſyſt. ſuppl. 146. 16. ♀ *cylindricollis.*

Habitat in Americae Inſulis. Dom. v. Rohr.

10. St. thorace mutico, pallidus, elytris vniſpinoſis: punctis tribus nigris, antennis longis. Ent. ſyſt. 2. 297. 17. ♀ *pallens.*

Habitat in Americae meridionalis Inſulis. Dom. Iſert.

11. St. thorace mutico, antice attenuato, ferrugineus, elytris pallidis. Ent. ſyſt. 2. 297. 18. ♂ *ſtrepens.*

*Cerambyx ſtrepens.* Oliv. Inſ. 67. tab. . fig .

Habitat in Gallo prouincia noctu volitans cum ſtrepitu. Muſ. Dom. Olivier.

12. St. thorace mutico, ſubrotundo, rufus, elytris teſtaceis: margine vittaque abbreuiata, cyaneis. *vittatus.*

Habitat in Braſilia. Dom. Weber.

Statura et magnitudo praecedentis. Antennae corpore longiores, ferrugineae. Caput ferruginenm. Thorax rotundatus, muticus, ſcaber, ferrugineus, annulo magno, cyaneo. Elytra teſtacea: margine exteriori vittaque media abbreuiata, cyaneis. Corpus et pedes ferruginea.

13. St. thorace mutico, atro, elytris bidentatis, baſi teſtaceis, apice atris: faſcia pallida. Ent. ſyſt. 2. 297. 19. ♀ *faſciatus.*

Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 67. tab. 14. fig. 98.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

*circumflexus*, 24. St. thorace mutico, ferrugineus, elytris vnidentatis, teſtaceis: punctis duobus ſtrigaque vndata, fuſcis. Ent. ſyſt. 2. 300. 20. *
Habitat in Cayenna. Dom. v. Rohr.

*nanus*, 25. St. thorace rotundato, mutico, griſeus, antennis pedibusque teſtaceis. Ent. ſyſt. 2. 300. 21. *
Habitat - - Muſ. Dom. Lund.
Forte potius Callidium. Elytra apice emarginata.

*annulatus*. 26. St. thorace lineato, elytris vnidentatis, antennis longis: annulis tribus albis. Ent. ſyſt. 2. 300. 22. *
*Cerambyx annulatus*. Oliv. Inſ. 67. tab. 16. fig. 117.
*Cerambyx hirtipes*. Degeer Inſ. 5. 116. 19. tab. 14. fig. 10.
Habitat in America meridionali. Muſ. D. Banks.

*tomaculatus*. 27. St. thorace ſubſpinoſo, quadrimaculato, elytris bidentatis, ferrugineis, flauo cinereoque maculatis. Ent. ſyſt. 2. 300. 23. *
Oliv. Inſ. 67. tab. 12. fig. 86.
Habitat in America. Muſ. D. Banks.

*lineola* 28. St. thorace ſpinoſo, ferrugineo, elytris acuminatis, teſtaceis: lineolis tribus glabris, flauis. Ent. ſyſt. 2. 301. 24. *
Oliv. Inſ. 67. tab. 2. fig. 19.
Habitat in Braſilia. Muſ. D. Banks.

*hemipterus* 29. St. thorace ſpinoſo, ater, elytris abbreuiatis.
Habitat in Jaua. Muſ. Dom. Boſc.
Magnus. Antennae atrae, altero latere ſpinoſae, longitudine corporis. Caput atrum, mandibulis exſertis, acutis. Thorax villoſus, ater vtrin-

que

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

que ſpina obtuſa. Elytra valde abbreuiàta, vix ſtriata, atra, apice acutiuſcula. Pedes atri, tibiis poſticis compreſſis.

30. St. thorace ſubſpinoſo, elytris bidentatis, cinereo fuſco flauoque variis: angulo baſeos compreſſo. Ent. ſyſt. 2. 301. 25. *varius.*
Habitat in Cayenna. Dom. v. Rohr.

31. St. thorace ſpinoſo, ater, elytris apice vniſpinoſis, baſi teſtaceis, antennis longis. Ent. ſyſt. 2. 301. 26. * *pubeſcens.*
*Cerambyx pubeſcens.* Oliv. Inſ. 67. tab. 18. fig. 135.
Habitat in Cayenna.

32. St. thorace mutico, obſcurus, cinereo-villoſus, elytris bidentatis. Ent. ſyſt. 2. 302. 27. * *villoſus.*
Habitat in Carolina.

33. St. thorace ſubſpinoſo, elytris truncatis, bidentatis, ferrugineis, cinereo-faſciatis, antennis breuibus. Ent. ſyſt. 2. 302. 28. * *Druryi.*
Habitat in America. Dom. Drury.

34. St. thorace ſpinoſo, cinereus, elytris bidentatis: faſciis duabus vndatis, nigris. Ent. ſyſt. 2. 302. 29. * *vndatus.*
Habitat in America. Dom. Drury.

35. St. thorace ſpinoſo fuſco cinereoque, griſeus, elytris punctato-ſtriatis. Ent. ſyſt. ſuppl. 147. 29.* *plebejus.*
Habitat in America. Dom. Hybner.

36. St. thorace ſpinoſo, ferrugineo, fuſcus, elytris bidentatis, pallidioribus, antennis longis. Ent. ſyſt. 2. 302. 30. * *ruſticus.*
Oliv. Inſ. 69. tab. 2. fig. 16.
Habitat in India orientali. Muſ. Dom. Banks.

37.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*angulatus.* 37. St. thorace mutico, cinereus, elytris macula marginali fasciaque angulata, nigris.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Paruus in hoc genere. Antennae corpore duplo longiores: articulo primo longiori, incuruo. Caput et thorax cinerea. Elytra laeuia, cinerea: macula magna marginali, fasciaque poſtica angulata, fuſcis, apice parum acuminata. Femora clauata.

## 138. CALOPVS.

*Palpi* quatuor;
*anteriores* clauati,
*poſteriores* filiformes.
*Maxilla* bifida.
*Ligula* membranacea, bifida
*Antennae* filiformes.

*ſerraticornis.* 1. CALOPVS.

*Cerambyx ſerraticornis.* Linn. ſyſt. Nat. 2. 634. 65. Fn. Sv. 665.

Degeer Inſ. 5. 79. 16.

*IV. 46.* Payk. Fn. Sv. 3. 65. 1.

Habitat in Europa boreali.

Animalculum elongatum, cylindricum, obſcurum.

139. RHA-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

139. RHAGIVM. *Palpi* quatuor, capitati.
*Maxilla* vnidentata.
*Ligula* membranacea, bifida.
*Antennae* ſetaceae.

1. R. thorace ſpinoſo, griſeum, elytris nebuloſis, teſtaceo-ſubfaſciatis. Ent. ſyſt. 2. 303. 1. *mordax.*
Schaeff. Icon. tab. 8. fig. 2.
Habitat in Germania. Dom. Schneider.

2. R. thorace ſpinoſo, nigrum, elytris nebuloſis, teſtaceo-ſubfaſciatis. Ent. ſyſt. 2. 304. 2. *inquiſitor.*
*Cerambyx inquiſitor.* Linn. ſyſt. Nat. 2. 630. 49. Fn. Sv. 659.
Geoff. Inſ. 1. 223. 2.
Frifh. Inſ. 13. tab. 14.
Oliv. Inſ. 69. tab. 2. fig. 12.
Degeer Inſ. 5. 126. 2. tab. 4. fig. 1.
Payk. Fn. Sv. 3. 66. 1.
Habitat in Europae truncis arborum.
*Larua* hexapoda, nuda, alba, capite collarique corneis, fuſcis, dorſo canaliculato. Stroem. Act. Hafn. 1.

3. R. thorace ſpinoſo, cinereum, elytris lineis tribus, eleuatis: antennis faſciisque duabus nigris. Ent. ſyſt. 2. 304. 3. *indagator.*
Degeer Inſ. 5. tab. 4. fig. 7.
Payk. Fn. Sv. 3. 66. 2.
Habitat in Germania. Dr. Schulz.

4.

---

*Rhagii* corpus oblongum, glabrum, immarginatum, tardum; *capite* magno, ouato, prominulo; *oculis* rotundatis, prominulis; *antennis* elongatis, approximatis, inter oculos inſertis; *thorace* anguſto, cylindrico, vtrinque ſpinoſo; *ſcutello* rotundato; *elytris* rigidis, fornicatis, thorace multo latioribus, longitudine abdominis; *pedibus* curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*curſor.* 4. R. thorace ſpinoſo, elytris rufis: ſutura lineaque nigris. Ent. ſyſt. 2. 304. 4.
*Cerambyx curſor.* Linn. ſyſt. Nat. 2. 630. 45. Fn. Sv. 658.
Degeer Inſ. 5. 127. 3.
Sulz. Hiſt. Inſ. tab. 5. fig. 7.
Payk. Fn. Sv. 3. 67. 3.
Habitat in Europa boreali.

*cinctum.* 5. R. thorace ſpinoſo, nigrum, elytris ſubferrugineis: faſcia flaua, femoribus poſticis vnidentatis. Ent. ſyſt. 2. 305. 5. *
Habitat in Auſtria. Dom. Saldoner.

*Salicis.* 6. R. thorace tuberculato, ſpinoſoque, rufum, pectore elytrisque nigris.
*Callidium Salicis.* Ent. ſyſt. 2. 329. 51. *
*Rhagium etruſcum.* Roſſ. Fn. Etr. 1. 149. 373. tab. 1. fig. 13.
*Stenocorus ruficollis.* Herbſt. Arch. t. 25. f. 13.
Habitat in Europae Salicibus.

*Noctis.* 7. R. thorace ſpinoſo, nigrum, antennarum baſi ferruginea. Ent. ſyſt. 2. 305. 6.
*Cerambyx Noctis.* Linn. ſyſt. Nat. 2. 630. 48.
Schaeff. Icon. tab. 38. fig. 2.
Panz. Fn. Germ. 17. tab. 3.
Habitat in Europa boreali.

*biſaſciatum.* 8. R. thorace ſpinoſo, elytris faſciis duabus obliquis flauis. Ent. ſyſt. 2. 305. 7. *
Degeer Inſ 5. 124. 1. tab. 4. fig. 6.
Sulz. Hiſt. Inſ. tab. 5. fig. 8.
Oliv. Inſ. 69. tab. 1. fig. 6.
Schaeff. Elem. tab. 118. fig. 1.
~~Icon. tab. 8. fig.~~ 1-3.
Herbſt. Arch. 170. tab. 45. fig. 12.
Geoff. Inſ. 1. 222. 1.
Habitat in Europa boreali. 9.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

9. R. thorace ~~sub~~spinoso, nigrum, elytris flauo subreticulatis, pedibus rufis. Ent. syst. 2. 306. 8. * *clathratum.*

Habitat in Austria. Dom. Saldoner.

10. R. thorace spinoso, nigrum, elytris fascia lata, flaua. Ent. syst. 2. 306. 9. * *ornatum.*

Habitat in Pensylvania. Mus. Dom. Banks.

11. R. thorace spinoso, elytris eleuato-striatis, cinereis, nigro-vndatis. Ent. syst. 2. 306. 10. *minutum.*

Oliv. Inf. 69. tab. 1. fig. 3.

Habitat Kiliae.

140. GNOMA. *Palpi* quatuor: articulo vltimo setaceo.
*Maxilla* bifida: lacinia exteriore apice crassiore.
*Ligula* cornea, apice rotundata, subemarginata.
*Antennae* setaceae.

1. G. nigra, ferrugineo-irrorata, antennis longissimis. *longicollis.*

*Cerambyx longicollis.* Ent. syst. 2. 265. 52. *

Oliv. Inf. 67. tab. 11. fig. 73.

Habitat in India orientali. Mus. Dom. Lund.

*Os* maxillis palpisque. *Palpi* quatuor, inaequales, filiformes: *anteriores* paullo longiores, quadri-

---

*Gnomae* corpus subcylindricum, glabrum, immarginatum, tardum; *capite* magno, cylindrico, inserto; *oculis* magnis, reniformibus, lateralibus; *antennis* longis, approximatis, in sinu oculorum insertis; *thorace* elongato, cylindrico, rugoso, elytris multo angustiore; *scutello* rotundato; *elytris* rigidis, fornicatis, abdomine longioribus; *pedibus* longiusculis, compressis; *tarsis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

driarticulati, articulo primo breuiſſimo, ſecundo longiori, incuruo, tertio breuiori, quarto longiori, ſetaceo, adhaerentes maxillae dorſo, *poſteriores* paullo breuiores, triarticulati, articulo vltimo ſetaceo, adnati ligulae medio. *Mandibula* cornea, incurua, compreſſa, acuta, edentula. *Maxilla* cornea, biſida: lacinia interiore breuiore, compreſſa, ciliata, exteriore ſubcylindrica, in medio craſſa, ore incurua, parum piloſa. *Ligula* porrecta, cornea, baſi anguſtata, compreſſa, apice rotundata, dilutata, ſubintegra. *Labium* breue, corneum, transuerſum, integrum. *Antennae* longiſſimae. Genus Saperdis affine.

*cylindricollis.* 2. G. oliuacea, albolineata, antennis longiſſimis, atris.

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Minor G. longicolli. Antennae corpore duplo longiores, atrae, articulo tertio breui, quarto quintoque baſi parum albidis. Caput oliuaceum, orbita oculorum lineisque frontalibus albis. Thorax cylindricus, elongatus, oliuaceus, lineis tribus integris, albis. Elytra punctato-ſtriata, oliuacea, humeris albis: puncto magno atro in medio, lineola parua, poneque medium ſtriga albidis. Corpus albidum, pedibus oliuaceis.

*clauipes.* 3. G. cinerea, femoribus clauatis.

Habitat in Americae meridionalis Inſulis. Muſ. D. de Seheſtedt.

Praecedentibus multo minor. Antennae villoſae, corpore longiores: articulis baſi ferrugineis, apice fuſcis. Thorax cylindricus, elongatus, inermis, dorſo tuberculis aliquot obſoletis. Ely-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Elytra punctis eleuatis, minutis, ſcabra, apice rotundata, integra. Corpus cinereum, femoribus valde clauatis.

4. G. thorace rugoſo, nigra, elytris ſanguineis. *rugicollis.*
Habitat in Carolina. Muſ. Dom. Boſc.
Media, antennae nigrae in altero ſexu corpore longiores. Thorax cylindricus, rugoſus, ater, immaculatus. Elytra porcata, ſanguinea, immaculata. Corpus nigrum, abdomine ſanguineo. Pedes atri, femoribus valde clauatis.

## 141. SAPERDA.

*Palpi* filiformes.
*Maxilla* membranacea, bifida.
*Ligula* cordata, truncata.
*Antennae* ſetaceae.

1. S. corpore griſeo, nigro, punctato, antennis mediocribus. Ent. ſyſt. 2. 307. 1. *carcharias.*
*Cerambyx carcharius.* Linn. ſyſt. Nat. 2. 631. 52.
Fn. Sv. 660.
Oliv. Inſ. 68. tab. 2. fig. 22.
Geoff. Inſ. 1. 308. 1.
Drury Inſ. 1. tab. 41. fig. 5.
Schaeff. Icon. tab. 152. fig. 4.
Panz. Fn. Germ. 69. tab. 1.
Degeer Inſ. 5. 73. 10. tab. 3. fig. 19.
Payk.

---

*Saperdae* corpus elongatum, cylindricum, glabrum, immarginatum, agile; *capite* rotundato, transuerſo, inſerto; *oculis* reniformibus, lateralibus; *antennis* elongatis, in oculorum ſinu inſertis; *thorace* breui, rotundato, inermi; *ſcutello* parno, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 71. 1.
Habitat in Europae ſyluis.

*ſcalaris.* 2. S. coleoptris linea ſuturali dentata punctisque flauis, antennis mediocribus. Ent. ſyſt. 2. 307. 2.
*Cerambyx ſcalaris.* Linn. ſyſt. Nat. 2. 632. 55. Fn. Sv. 697.
Oliv. Inſ. 68. tab. 1. fig. 7.
Panz. Fn Germ. 64. tab. 3.
Voet. Coleopt. 2. tab. 17. fig. 78.
Friſh. Inſ. 12. 29. tab. 3. fig. 3.
Schaeff. Icon. tab. 38. fig. 5.
Degeer Inſ. 5. 77. 14.
Payk. Fn. Sv. 3. 72. 2.
Habitat in Europae Populo.

*vittigera.* 3. S. fuſco-griſea, elytris vitta lata, alba: punctis impreſſis, nigris.
Habitat in Barbaria. Dom. de Rehbinder.
Statura et magnitudo S. ſcalaris. Antennae corpore breuiores, fuſcae. Caput et thorax punctata, griſea. Elytra plana, punctata, griſea. vitta lata, dorſali, alba et in hac puncta impreſſa, nigra, magis diſtincta. Corpus obſcure griſeum.

*laeta.* 4. S. atra, thorace margine elytris maculis flauis punctisque duobus albis.
Oliv. Inſ. 68. tab. 9. fig. 76.
Habitat in Guinea. Meier. Muſ. D. Lund.
Statura et magnitudo S. ſcalaris. Antennae corpore breuiores, atrae. Caput atrum, macula magna frontali orbitaque oculorum laete flauis. Thorax lateribus late flauis. Elytra atra, maculis duabus magnis ad marginem exteriorem coëuntibus flauis. Pone has puncta duo alba et verſus apicem macula minor flaua. Sutura parum albicat.

5.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. S. alba, thorace coleoptrisque fuſcis: vittis duabus albis. Ent. ſyſt. 2. 307. 3. * *candida.*
Habitat - - Muſ. Dr. Hunter.

6. S. nigra, cinereo-villoſa, thorace linea dorſali, elytris vitta albis, antennis breuibus. Ent. ſyſt. 2. 308. 4. * *detrita.*
Habitat in Barbaria. Muſ. Dom. Desfontaines.

7. S. liuida, antennis elytrorumque punctis atris. Ent. ſyſt. ſuppl. 147. 4. * *morbilloſa.*
Habitat Tranquebariae. Muſ. D. Lund.

8. S. atra, elytris, albo-punctatis, antennis longis. Ent. ſyſt. 2. 308. 5. * *irrorata.*
Habitat in Africae plantis. Dom. Vahl.
*Variat* elytris immaculatis, forte aetate.

9. S. nigra, capite thorace anoque ferrugineis. Ent. ſyſt. 2. 308. 6. * *modeſta.*
Habitat in Africa aequinoctiali. Muſ. D. Banks.

10. S. ferruginea, antennis abdominisque apice ſupra nigris, elytris emarginatis. Ent. ſyſt. 2. 308. 7. * *atricornis.*
Habitat in China. Muſ. D. de Seheſtedt.

11. S. thorace luteo: punctis duobus atris, elytris nigris. Ent. ſyſt. 2. 308. 8. *oculata.*
*Cerambyx oculatus.* Linn. ſyſt. Nat. 2. 633. 60.
Fn. Sv. 664.
Oliv. Inſ. 68. tab. 1. fig. 4.
Panz. Fn. Germ. 1. tab. 18. *L. 44.*
Degeer Inſ. 5. 74. 11. tab. 3. fig. 20.
Schaeff. Icon. tab. 128. fig. 4.
Hoefnag. Inſ. tab. 13.
Payk. Fn. Sv. 3. 72. 2.
Habitat in Europa ſyluis.

12.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*tricolor.* 12. S. ferruginea, elytris bidentatis, punctato-ſtriatis, vireſcentibus, antennis mediocribus nigris. Ent. ſyſt. 2. 309. 9. *

Habitat in India orientali. Muſ. D. de Seheſtedt.

*villoſa.* 13. S. ferrugineo-griſeo-villoſa, ſcutello orbitaque oculorum fuluis.

*Saperda hirta.* Ent. ſyſt. 2. 309. 10. *

Habitat in noua Zelandia. Muſ. D. Banks.

*vnicolor.* 14. S. teſtacea, antennis pedibusque concoloribus, antennis longis. Ent. ſyſt. 2. 309. 11. *

Habitat in Inſula Amſterdam. Muſ. D. Banks.

*linearis.* 15. S. cylindrica, nigra, pedibus luteis, antennis mediocribus. Ent. ſyſt. 2. 309. 12.

*Cerambyx linearis.* Linn. ſyſt. Nat. 2. 632. 58. Fn. Sv. 663.

Oliv. Inſ. 68. tab. 2. fig. 13.

Schaeff. Icon. tab. 55. fig. 6.

I. 44. Herbſt. Arch. tab. 26. fig. 1.

Payk. Fn. Sv. 3. 73. 3.

Habitat in Europae Corylo.

*nigripes.* 16. S. cylindrica, nigra, thoracis lineis duabus ſcutelloque cinereis, pedibus nigris. Ent. ſyſt. 2. 310. 13. *

Habitat in Hungaria. Dom. Smidt.

*cylindrica.* 17. S. cylindrica, nigra, pedibus anticis luteis. Ent. ſyſt. 2. 310. 14.

*Cerambyx cylindricus.* Linn. ſyſt. Nat. 2. 633. 59. Fn. Sv. 662.

Oliv. Inſ. 68. tab. 2. fig. 11.

V. 22 Roeſ. Inſ. 2. ſcarab. 2. tab. 3.

Geoff. Inſ. 1. 208. 2.

Degeer Inſ. 5. 75. 12.

Panz Fn. Germ. 69. tab. 4.

Payk.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 73. 4.
Habitat in Pyri et Pruni ramis.

18. S. cylindrica, atra, thorace ſupra ferrugineo: punctis tribus nigris. Ent. ſyſt. 2. 310. 15. * *tripunctata.*
Habitat in Virginia.

19. S. cylindrica, thorace ferrugineo: punctis tribus atris, elytris atris: vitta abbreuiata teſtacea. *mandarina.*
Habitat in Carolina. Muſ. D. Boſc.
Affinis certe S. tripunctatae, vix diſtincta. Caput ferrugineum, antennis nigris. Thorax ferrugineus, punctis tribus atris, duobus in medio tertioque in medio baſeos. Elytra punctato-ſtriata, atra: vitta lata media pallida, quae tamen apicem haud attingit.

20. S. thorace rufo, corpore nigro, pedibus rufis. Ent. ſyſt. 2. 310. 16. * *melanocephala.*
Habitat in Africa. D. Prof. Vahl.

21. S. violacea, antennarum articulo primo ſecundoque clauato-piloſis. Ent. ſyſt. 2. 310. 17. * *pilicornis.*
Pall. Icon. tab. F. fig. 12.
Habitat in America meridionali. Dom. Hybner.

22. S. atra, elytris emarginatis baſi flauis, antennis barbatis. *humeralis.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura praecedentium. Antennae corpore paullo longiores, nigrae, pilis longioribus barbatae. Caput et thorax nigra. Elytra ſtriata, emarginata, nigra, baſi flaua. Corpus nigrum, cinereo-pubeſcens.

23. S. capite thoraceque rufis, elytris cyaneis, antennis mediocribus ante apicem barbatis. Ent. ſyſt. 2. 311. 18. * *barbicornis.*
Habitat in Brasilia.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*erythrocephala.* 24. S. thorace villoso, rufa, antennis, pectore elytrisque nigris. Ent. syst. 2. 311. 19. *
*Cerambyx erythrocephalus.* Schrank. Austr. 270.
Panz. Fn. Germ. 69. tab. 5.
Habitat in Germania. Dom. Hybner.

*discoidea.* 25. S. thorace villoso, virescens, coleoptrorum disco fusco: punctis duobus fasciaque communi lunata virescentibus. Ent. syst. suppl. 147. 19. *
Habitat in America boreali. D. v. Rohr.

*ruficollis.* 26. S. thorace villoso, rufo, antennis elytrisque fuscis. Ent. syst. 2. 311. 20. *
Habitat in Virginia.

*fasciata.* 27. S. thorace subspinoso, cyanea, elytris fasciis duabus flauis. Ent. syst. 2. 311. 21. *
Oliv. Inf. 68. tab. 3. fig. 24.
Habitat in Sibiria. Mus. D. Banks.

*clauicornis.* 28. S. viridis, elytris maculis tribus flauis. Ent. syst. 2. 311. 22. *
Oliv. Inf. 68. tab. 3. fig. 30.
Habitat ad Cap. Bon. Spei. Mus. D. Banks.

*latipes.* 29. S. nigra, femoribus clauatis apice violaceis, tibiis posticis compressis, antennis longis. Ent. syst. 2. 312. 23. *
Habitat ad Cap. Bon. Spei. Mus. Prof. Leske.

*vittata.* 30. S. grisea, vittis tribus dentatis albis, antennis longis. Ent. syst. 2. 312. 24. *
*Cerambyx trilineatus.* Linn. Mant. 352.
Drury Inf. 1. tab. 41. fig. 1.
Oliv. Inf. 67. tab. 19. fig. 142.
Habitat in America. D. Drury.

*6punctata.* 31. S. nigra, elytris punctis tribus flauis. Ent. syst. 2. 312. 25. *
Habitat ad Cap. Bon. Spei. D. de Paykull.

32. S

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

32. S. griſea, antennis breuibus atris, apice griſeis, elytris apice acuminatis nigris. *acuminata*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura parua S. lateralis. Antennae breues, craſſiores, articulis tribus vltimis griſeis. Thorax griſeus, linea dorſali lata lateralibusque nigris. Elytra laeuia, griſea, ſutura late apiceque acuminato nigris. Corpus cinereum.

33. S. nigra, capite thoracisque ſpinoſi lateribus rufis, antennis breuibus, apice griſeis. *rufi-cornis.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo praecedentis. Antennae breues, craſſae, nigrae, apice cineraſcentes. Caput rufum. Thorax ſpinoſus, niger, lateribus late rufis. Elytra laeuia, nigra, apice emarginata. Corpus nigrum.

34. S. nigra, thoracis lateribus elytrorumque vitta laterali ferrugineis. Ent. ſyſt. 2. 312. 26. * *lateralis.*

Habitat in America boreali. Muſ. D. Banks.

35. S. thorace atro, puncto vtrinque ferrugineo, elytris acuminatis griſeis. Ent. ſyſt. 2. 313. 27. * *lyncea.*

Habitat in noua Zelandia. Muſ. D. Banks.

36. S. cinerea, thorace elytrisque nigro-punctatis, antennis longis. *Boſcii.*

Habitat in Iaua. Muſ. D. Boſc.

Corpus medium. Antennae corpore duplo longiores, nigrae. Caput cinereum, macula verticali nigra. Thorax rotundatus, muticus, laete cinereus, punctis nigris dorſalibus tribus 2. 1. baſeos maiori, transuerſo lateralibusque vtrinque tribus. Elytra laeuia, laete cinerea,



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

punctis plurimis nigris: tribus communibus. Pedes elongati, nigri.

*griſea.* 37. S. griſea, ſcutello margine, elytris lineolis flaueſcentibus. Ent. ſyſt. 2. 213. 28. *

Habitat in noua Hollandia. Muſ. D. Banks.

*gracilis.* 38. S. ferruginea, elytris teſtaceis: vitta marginali antennisque nigris.

Habitat in Carolina. Muſ. D. Boſc.

Statura omnino S. analis. Caput ferrugineum, antennis totis nigris. Thorax cylindricus, rufus, immaculatus. Elytra plana, punctato-ſtriata, teſtacea, vitta marginali nigra. Corpus rufum.

*biloba.* 39. S. thorace rotundato rugoſo, cinerea, coleoptris maculis quinque niueis: communi biloba, antennis longis.

Habitat in Inſulis Oceani auſtralis. D. Billardiere.

Statura et magnitudo S. griſeae. Antennae longiſſimae, teſtaceae: articulo primo griſeo. Caput griſeum, ſub antennis niueum. Thorax cylindricus, rugoſus, griſeus, punctis duobus baſeos niueis. Scutellum niueum. Elytra laeuia, lineola marginali baſeos puncto pone medium macula maiori, communi biloba, baſeos niueis. Corpus niueum.

*virginea.* 40. S. cinerea, thorace ſubſpinoſo: maculis duabus nigris, elytris apice bidentatis flauis: faſcia griſea.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Parua. Caput flauum, antennis longitudine corporis nigris. Thorax ſubſpinoſus, cinereus, maculis duabus marginis antici fuſcis. Elytra cinerea margine laterali nigro, apice emarginata, flaua: faſcia fuſca. Corpus cinereum.

41. S.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

41. S. teſtacea, elytris apice anoque nigris. Ent. ſyſt. *analis.*
2. 313. 29. *

Habitat in Africa aequinoctiali. Muſ. D. Banks.

42. S. thorace rotundata, flaua, antennis elytrisque *bidentata.*
apice bidentatis, nigris. Ent. ſyſt. 2. 313. 30. *

Habitat in Aquapim Guineae. Dr. Iſert.

*Variat* capite nigro et flauo, thorace punctis quatuor duobus, aut nullis.

43. S. griſea, elytris apice rotundatis, atris: litura *liturata.*
flaua.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Parua. Antennae corpore longiores, nigrae. Caput griſeum, ſtriga verticali flaua. Thorax griſeus, linea dorſali flaueſcente. Elytra punctata, griſea, apice rotundata, atra: litura flaueſcente. Corpus griſeum, femoribus clauatis.

44. S. obſcura, elytris bidentatis, fuſcis: vitta gri- *varicornis*
ſea, antennis elongatis, atris, ante apicem albis.

Habitat in Sumatra. Dom. Daldorff. Muſ. D. de Seheſtedt.

Statura parua S. cylindricae. Antennae corpore longiores, atrae: articulo vltimo baſi late albo. Caput obſcurum. Thorax cylindricus, inermis, niger: linea laterali pallida. Scutellum pallidum. Elytra punctata, fuſca: vitta lata media palleſcente, apice bidentata. Corpus fuſcum: lateribus palleſcentibus.

45. S. fuſca, thorace lineato, ſcutello flauo, anten- *Cardui.*
nis longis.

*Saperda nigricornis.* Ent. ſyſt. 2. 314. 31. *
*Cerambyx Cardui.* Linn. ſyſt. Nat. 2. 632. 56. VI. 23
Oliv. Inſ. 68. tab. 1. fig. 5.
Panz. Fn. Germ. 69. tab. 6.

Herbſt.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Arch. tab. 26. fig. 2.
Habitat in Europae auſtralioris Carduis.

*birſutula.* 46. S. villoſa, cinerea, thorace lineato, elytris nigro-irroratis.
Habitat in Auſtria. Dom de Meyerle.
IX. Nimis affinis S. nigricorni, at magis villoſa et elytra magis nigro-irrorata. Forte mera varietas.

*nigricornis.* 47. S. cinerea, thorace lineato, elytris fuſcis vnicoloribus, antennis mediocribus.
IX. *Saperda futuralis.* Ent. ſyſt. 2. 314. 32. *
Habitat Lipſiae. Prof. Leske.

*futuralis.* 48. S. thorace lineato, vireſcens, elytris acuminatis: futura alba.
*Saperda annulata.* Ent. ſyſt. 2. 314. 33. *
Oliv. Inſ. 68. tab 2. fig. 16.
II. 41. Panz. Fn. Germ. 23. tab. 16.
Roſſ. Fn. Etr. 1. 152. 379.
Habitat in Africae plantis. Dom. Prof. Vahl.

*annulata.* 49. S. thorace rotundato, ſubſpinoſo, antennarum articulis baſi albis.
*Saperda lineata.* Ent. ſyſt. 2. 314. 34. *
Habitat in America meridionali. Muſ. D. Lund.

*lineata.* 50. S. thorace rotundato, nigro, albo-lineato, elytris bidentatis, teſtaceis, albo-lineatis.
*Saperda triſtis.* Ent. ſyſt. 2. 314. 35. *
Habitat Tranquebariae. Muſ. D. Lund.

*triſtis.* 51. S. thorace quadriſpinoſo, corpore fuſco. Mant. Inſ. 1. 220. 23. *
Habitat in Germania.

*ferrea.* 52. S. thorace rotundato, hirta, fuſca, elytris punctatis.
IX. *Saperda ferrea.* Schrank. Auſtr.
Habitat in Auſtria. Dom. de Meierle.
variet. nigripedis

Cor-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Corpus totum fuſcum, griſeo-hirtum. Thorax rotundatus, muticus. Elytra punctis impreſſis.

53. S. thorace fuſco, albolineato, elytris acuminatis, fuſcis: ſtriga abbreuiata lineolaque apicis albis. *acuta.*

Habitat in Sumatra. D. Daldorff.

Media. Antennae breues, fuſcae: articulo tertio elongato, apice albo. Caput fuſcum, lineis quatuor albis. Thorax rotundatus, inermis, fuſcus: lineis tribus albis. Elytra plana, punctata, fuſca, ſtriga media abbreuiata, externe angulum paruum formante, alba, apice acuminata, lineola parua, albida. Corpus fuſcum, vittis niueis.

54. S. thorace mutico, atro: lateribus niueis, elytris cinereis, apice bidentatis, fuſcis: macula fulua. *dimidiata.*

Habitat in Sumatra. Dom. Daldorff.

Parua. Antennae obſcure-ferrugineae longitudine corporis. Caput nigrum, orbita oculorum cinerea. Thorax rotundatus, muticus, ater, lateribus niueis. Elytra plana, baſi punctata, cinerea, apice bidentata, fuſca, macula laete fulua. Corpus niueum, ano nigro. Pedes teſtacei.

55. S. thorace flauo, lineato, elytris punctis quatuor flauis, antennis mediocribus. Ent. ſyſt. 2. 315. 37. *populnea.*

*Cerambyx populneus.* Linn. ſyſt. Nat. 2. 632. Fn. Sv. 661. VI. 23

Oliv. Inſ. 68. tab. 1. fig. 1.

Geoff. Inſ. 1. 208. 3.

Panz. Fn. Germ. 69. tab. 7.

Schaeff. Icon. tab. 48. fig. 5.

Habitat in Europae Americae Populetis.

56. S. viridis, thorace punctis duobus, elytris quatuor nigris. Ent. ſyſt. 2. 315. 38. *tremula.*

Panz. 5. 45.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 1. tab. 7.
Schaeff. Icon. tab. 101. fig. 1.
Habitat Halae Saxonum. Dr. Schreber.

*punctata.* 57. S. viridis, punctis nigris numeroſis, antennis mediocribus. Ent. ſyſt. 2. 315. 39.
*Cerambyx punctatus.* Linn. ſyſt. Nat. 2. 1067. 7.
Oliv. Inſ. 68. tab. 1. fig. 9.
Sulz. Hiſt. Inſ. tab. 5. fig. 10.
Habitat in Europa auſtrali.

*Scidlii.* 58. S. cinerea, thorace punctis octo, elytris quinque nigris.
~~*Saperda Scidlii.* Panz.~~ Nat. 24. 28. 36. tab. 1. fig. 36.
Habitat in Auſtria. Dom. de Meyerle.
Statura omnino et magnitudo S. tremulae. Antennae mediocres, cinereae, articulis apice nigris. Caput cinereum, fronte nigra. Thorax cinereus, vtrinque punctis quatuor nigris. Elytra planiuſcula, cinerea, punctis quinque nigris, praeterea linea lateralis a baſi vltra medium ducta et ſub hac punctum ſolitarium. Corpus cinereum.

*viresceus.* 59. S. thorace villoſo, cinereo, elytris ſubattenuatis, viridibus. Ent. ſyſt. 2. 315. 40. *
Oliv. Inſ. 68. tab. 2. fig. 11.
Panz. Naturf. 24. 28. 37. tab. 1. fig. 37.
Habitat in Italia, Dr. Allioni; in Galliae Sambuco, Dr. Brouſſonet.

*tenuicornis.* 60. S. nigra, thoracis lateribus, elytrorum macula baſeos faſciaque media ferrugineis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Media. Antennae corpore duplo longiores, tenuiſſimae, nigrae: articulo primo craſſiori, longiſſimo.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

fimo. Caput nigrum, ore lineaque inter antennas ferrugineis. Thorax holoſericeus, niger, lateribus rufis. Elytra nigra, macula magna baſeos cum thoracis margine coëunte faſciaque mediaque ferrugineis. Corpus nigrum, pedibus flaueſcentibus.

61. S. capite thoraceque rufis, nigro-punctatis, elytris griſeis, immaculatis. *Argus.*

Panz. Fn. Germ. 69. tab. 2.

Habitat in Hungaria. Dom. de Meyerle.

Paullo minor S. tremula. Antennae corpore breuiores, nigrae. Caput rufum, puncto medio frontali tribusque baſeos nigris. Thorax rotundatus, rufus: punctis ſeptem 4. 3. eleuatis, glabris, atris. Elytra pubeſcentia, griſea. Corpus griſeum, ano late rufo. Pedes rufi femoribus, puncto apicis nigro.

62. S. fuſca, pedibus teſtaceis, antennis longiſſimis, nigris: annulo albo. Ent. ſyſt. 2. 315. 41. *longicornis*

Habitat in Africa aequinoctiali. Muſ. D. Banks.

63. S. fuſca, antennis obſcuris, pedibus teſtaceis. *flauipes.*

Habitat in Tanger. Dr. Schousboe. Muſ. D. de Scheſtedt.

Statura et magnitudo praecedentium. Antennae obſcurae: articulo primo longiori, nigro. Caput, thorax, elytra ſubtruncata, nigra, immaculata. Pedes teſtacei.

64. S. fuſca, cinereo-trilineata, antennis longiſſimis. *antennata.*

Habitat in Guinea. Dom. Heſſe. Muſ. D. Lund.

Corpus tenue, elongatum. Antennae corpore duplo longiores, fuſcae. Caput et thorax fuſca, lineis tribus cinereis. Elytra piceo-fuſca, lineis obſoletis, cinereis, apice rotundata.

65. S. niuea, dorſo atomisque numeroſiſſimis atris. *marmorea irrorata.*

Habitat in Jaua. Muſ. Dom. Boſc. Paul-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paullo maior S. voluulo. Antennae corpore breuiores, nigrae: articulis basi albis. Caput villosum, album, punctis numerosis, atris. Thorax elytraque villosa, niuea: dorso atomisque numerosis, atris.

*femorata.* 66. S. thorace antice fusco, postice testaceo, elytris fasciis tribus nigris testaceisque, antennis longissimis. Ent. syst. 2. 316. 42. *

Habitat - - Mus. Britt.

*haemorrhoidalis.* 67. S. atra, thorace punctis tribus sanguineis, ano pedibusque testaceis. Ent. syst. 4. Append. 451.

Habitat in Italia. Dr. Allioni.

*voluulus.* 68. S. supra nigra, thoracis elytrorumque margine cinereo. Ent. syst. 2. 316. 43. *

Habitat in Cayenna. Dom. v. Rohr.

*lemniscata* 69. S. fusca, cinereo-sexlineata, antennis longissimis.

Habitat in Carolina. Mus. Dom. Bosc.

Media. Antennae corpore multo longiores, barbatae, fuscae: articulo primo longiori. Caput fuscum, lineis quatuor cinereis. Thorax cylindricus, punctatus, fuscus, lineis sex et elytra punctata, lineis tribus, quarum interiores apice coëunt. Corpus fuscum.

Affinis videtur S. antennatae.

*ferruginea* 70. S. thorace subspinoso, ferruginea, antennis pedibusque nigris. Ent. syst. 2. 316. 44. *

*Cerambyx cantharinus.* Linn. syst. Nat. 2. 637. 82.

III. 54

Panz. Fn. Germ. 34. tab. 14.

Habitat in Germania. Dom. de Hattorff.

*spissicornis.* 71. S. nigra. thoracis linea dorsali, elytrorum sutura maculisque albis, antennis hirtis.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt. Par-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Parua. Antennae longitudine corporis, hirtae, nigrae: articulo 1. 2. tertioque baſi incraſſatis. Thorax rotundatus, niger: linea dorſali lateralique albis. Elytra nigra ſutura, lineolis baſeos maculisque apicis albis. Apex rotundatus. Abdomen nigrum, margine albo.

72. S. thorace ſubſpinoſo, ferruginea, antennis pedibusque concoloribus. Ent. ſyſt. 2. 316. 45. * *brunnea.*
Panz. Fn. Germ 34. tab. 15.
Habitat in Germania. Dom. Prof. Helwig.

73 S. teſtacea, thorace ferrugineo, antennis fuſcis. *tuberculata.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus paruum, teſtaceum. Antennae nigrae: articulo primo teſtaceo. Thorax ſubtuberculatus, ruſeſcens. Elytra punctata, rotundata, teſtacea.

74. S. nigra, elytris teſtaceis. Ent. ſyſt. 2. 316. 46. * *teſtacea.*
Habitat in Germania. Dom. de Hattorff.

75. S. corpore violaceo, immaculato. Ent. ſyſt. 2. 316. 47. * *violacea.*
*Saperda micans.* Panz. Fn. Germ.
Oliv. Inſ. 68. tab. 2. fig. 12.
Habitat in Italia. Dr. Allioni.

76. S. glabra, atra, thoracis margine flauo. Ent. ſyſt. ſuppl. 148. 47. * *marginata*
Habitat in Carolina. Dom. Boſc.

77. S. nigra, elytris flauis, apice nigris. Ent. ſyſt. 2. 317. 48. *praeuſta.*
*Leptura praeuſta.* Linn. ſyſt. Nat. 2. 641. 24.
Geoff. Inſ. 1. 209. 4.
Schaeff. Icon. tab. 52. fig. 8.
Payk. Fn. Sv. 3. 79. 12.
Habitat in Europa auſtraliori.

78.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ephippium.* 78. S. nigra, thoracis linea dorſali ſcutelloque cinereis, femoribus ferrugineis. Ent. ſyſt. 2. 317. 49. *

*Il. 41.* Panz. Fn. Germ. 23. tab. 17.
Habitat in Hungaria. Dom. Hybner.

*picea.* 79. S. picea, antennis pedibusque ferrugineis. Ent. ſyſt. 2. 317. 50. *
Habitat in America ſeptentrionali. D. Lund.

*birticornis* 80. S. griſea, antennis longis, hirtis. Ent. ſyſt. ſuppl. 148. 50. *
Habitat in India orientali. Muſ. D. Lund.

*marginella* 81. S. griſea, thoracis linea dorſali, ſcutello elytrorumque ſutura albis, antennis longis.
Habitat in Auſtria. Dom. de Meyerle.
Statura parua, filiformis praecedentium. Antennae corpore longiores, nudae, nigrae. Caput et thorax cineraſcentia, linea dorſali albidiore. Scutellum albidum, puncto nigro. Elytra cineraſcentia, ſutura margineque albidis. Pedes nigri.

*birta.* 82. S. nigra, hirta, antennis longis. Ent. ſyſt. 2. 317. 51. *
*Saperda filum.* Roſſ. Fn. Etr. 1. 152. 380. tab. 5. fig. 10.
Habitat in Italia. Dr. Allioni.

*ſcutellata.* 83. S. thorace ſubſpinoſo, nigra, antennis, elytris pedibusque cineraſcentibus, ſcutello albo. Ent. ſyſt. 2. 317. 52. *
Habitat in Germania. Muſ. Dom. de Seheſtedt.

*necydalina* 84. S. obſcura, elytris abbreuiatis, pallidis, antennis longis, ferrugineis. Ent. ſyſt. ſuppl. 148. 52. *
Habitat in Americae meridionalis Inſulis. Muſ. D. Lund.

*rufimana.* 85. S. cyanea, antennis nigris, femoribus anticis rufis.

Sa-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Saperda rufimana.* Panz. Fn. Germ.
Habitat in Auſtria. Dom. de Meyerle. III. 55.
Parua. Antennae longitudine corporis, nigrae. Corpus totum pubeſcens, cyaneum, immaculatum, femoribus anticis ſolis rufis. Elytra punctata.

86. S. nigra, thoracis linea dorſali femoribusque apice rufis. Ent. ſyſt. 2. 318. 53. * *lineola.*
Panz. Fn. Germ. 23. tab. 18. II. 41
Herbſt. Arch. tab. 26. fig. 4.
Habitat in Italia. Dr. Allioni.

142. CALLIDIVM. *Palpi* quatuor, clauati.
*Maxilla* membranacea, bifida.
*Ligula* bifida: laciniis tenuiſſimis.
*Antennae* ſetaceae.

1. C. thorace ſubuilloſo, fuſco, elytris teſtaceo cinereoque variis, antennis mediocribus. Ent. ſyſt. 2. 348. 1. * *obſcurum.*
Habitat in Terra Diemenii. Muſ. Dom. Banks.

2. C. thorace villoſo: tuberculis duobus, corpore fuſco. Ent. ſyſt. 2. 318. 2. *bajulus.*
*Cerambyx bajulus.* Linn. ſyſt. Nat. 2. 636. 76. VI. 24
Fn. Sv. 672.
Oliv.

---

*Callidii* corpus planiuſculum, ſaepius pubeſcens, depreſſiuſculum, immarginatum, tardum; *capite* ouato, obtuſo, inſerto; *oculis* reniformibus, marginalibus; *antennis* thorace longioribus, in ſinu oculorum inſertis; *thorace* plano: margine rotundato, prominulo; *ſcutello* paruo, triangulo; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* elongatis, curſoriis, femoribus ſaepius clauatis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 70. tab. 3. fig. 30.
Friſh. Inſ. 13. tab. 10.
Panz. Fn. Germ. 70. tab. 1.
Sulz. Inſ. tab. 4. fig. 29.
Voet. Col. 2. tab. 22. fig. 111. 112.
Schaeff. Icon. tab. 68. fig. 1.
Elem. tab. 76. fig. 4.
Payk. Fn. Sv. 3. 85. 6.
Degeer Inſ. 5. 86. 22.
Habitat in Europae truncis.
Elytra in altero ſexu pallida.

*fennicum.* 3. C. thorace tuberculato, rufo, elytris violaceis, antennis mediocribus. Ent. ſyſt. 2. 319. 3.
*Cerambyx fennicus.* Linn. ſyſt. Nat. 2. 636. 77. Fn. Sv. 664.
Oliv. Inſ. 70. tab. 1. fig. 9.
Geoff. Inſ. 1. 219. 19.
Panz. Fn. Germ. 70. tab. 2.
Voet. Col. 2. tab. 20. fig. 97.
Schaeff. Icon. tab. 4. fig. 12.
Herbſt. Arch. tab. 26. fig. 8.
Payk. Fn. Sv. 3. 87. 9.
Habitat in Europae truncis.

*ruficolle.* 4. C. thorace ſubcylindrico, ſpinoſo, rufo, nigra, elytris violaceis, antennis mediocribus, piceis. Ent. ſyſt. 2. 319. 4. *
Oliv. Inſ. 70. tab. 2. fig. 27.
Habitat in Italia. Dr. Allioni.

*ebulinum.* 5. C. thorace tuberculato, nigrum, elytris violaceis. Ent. ſyſt. 2. 319. 5. *
*Cerambyx ebulinus.* Linn. ſyſt. Nat. 2. 627. 83. *
Habitat in Africae plantis. Dom. Prof. Vahl.

6.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. C. thorace laeui, fuſca, elytris pallidis: lineis duabus, poſtice coëuntibus fuſcis. *tenellum.*
Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.
Statura parua C. ruficollis. Antennae mediocres, ferrugineae. Caput et thorax rotundatus obſcura. Elytra punctata, integra, pallida: lineis duabus, ante apicem coëuntibus, fuſcis. Corpus flaueſcens, pedum geniculis nigris.

7. C. fuſcum, thorace elytrisque viridi-aeneis, femoribus ferrugineis. Ent. ſyſt. 2. 319. 6. * *aeneum.*
Oliv. Inſ. 70. tab. 4. fig. 46.
Habitat in India. Muſ. Dr. Hunter.

8. C. nigrum, opacum, femoribus omnibus clauatis, antennis longis. Ent. ſyſt. 2. 320. 7. * *clauipes.*
Oliv. Inſ. 70. tab. 4. fig. 33.
Panz. Fn. Germ. 70. tab. 3. VI-24
Herbſt. Arch. tab. 26. fig. 9.
Payk. Fn. Sv. 3. 81. 2.
Degeer Inſ. 5. 88. 23.
Habitat in Europae truncis. Dom. de Hattorff.

9. C. thorace ſpinoſo, nudo, atrum, antennis breuibus. Ent. ſyſt. 2. 320. 8. * *ſpinoſum.*
Habitat in Hungaria. Dom. Hybner.

10. C. thorace ſubpubeſcente, corpore violaceo, antennis breuibus. Ent. ſyſt. 2. 320. 9. *violaceum.*
*Cerambyx violaceus.* Linn. ſyſt. Nat. 2. 635. 70.
Fn. Sv. 667.
Friſch. Inſ. 12. tab. 3.
Oliv. Inſ. 70. tab. 1. fig. 2.
Herbſt. Arch. tab. 26. fig. 10. VI: 25
Panz. Fn. Germ. 70. tab. 4.
Voet. Col. 2. tab. 23. fig. 123.
Schaeff. Icon. tab. 4. fig. 13.
tab. 234. fig. 1.
Habitat in Europae ſyluis. II.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*punctatum* 11. C. thorace pubeſcente, nigrum, punctatum, femoribus rubris, antennis mediocribus. Ent. ſyſt. ſuppl. 149. 10. *

Habitat Halae Saxonum. Dom. Hybner.

*femoratum* 12. C. thorace nudo, corpore atro, opaco, femoribus rubris, antennis mediocribus. Ent. ſyſt. 2. 320. 10.

*Cerambyx femoratus.* Linn. ſyſt. Nat. 2. 634. 69.

Oliv. Inſ. 70. tab. 7. fig. 77.

Schaeff. Icon. tab. 55. fig. 7.

Degeer Inſ. 5. 93. 81.

Panz. Fn. Germ. 70. tab. 5.

Habitat in Germaniae nemoribus.

*falcratum.* 13. C. thorace nudo, nitido, corpore atro, femoribus rufis, antennis mediocribus. Ent. ſyſt. 2. 320. 11. *

*Callidium aulicum.* Payk. Fn. Sv. 3 83. 3. β.

Habitat Halae Saxonum. Dom. Hybner.

*amethyſtinum.* 14. C. pubeſcens, cyaneum, pedibus rufis: femoribus claua atra. Ent. ſyſt. 2. 321. 12. *

Habitat in ~~America~~ Germania. Prof. Helwig.

*2faſciatum* 15. C. thorace pubeſcente, atrum, elytris faſciis duabus rufis. Ent. ſyſt. 2. 321. 13. *

Oliv. Inſ. 70. tab. 4. fig. 42.

Habitat in America meridionali. Muſ. D. Banks.

*acuminatum.* 16. C. thorace verrucoſo, nigricante, elytris acuminatis viridibus: ſutura cyanea. Ent. ſyſt. 2. 321. 14. *

Habitat ad Cap. Bon. Spei. Muſ. Dom. Lee.

*ruſticum.* 17. C. thorace verrucoſo, nigrum, elytris teſtaceis: macula media apiceque nigris. Ent. ſyſt. 2. 321. 15. *

Oliv. Inſ. 70. tab. 4. fig. 49.

Habitat in Ruſſia. Dom. Lee.

18.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

8. C. thorace glabro, ſubſpinoſo, nigrum, elytris obſcuro-aeneis. Ent. ſyſt. ſuppl. 149. 15. * — *hungaricum.*

*Callidium hungaricum.* Herbſt. Arch. tab. 26. fig. 12.
Panz. Fn. Germ. 70. tab. 7.
Habitat in Germania. Dom. Hybner.

9. C. thorace glabro, corpore fuſco, aeneo, antennis pedibusque fuſcis. Ent. ſyſt. 2. 321. 16. * — *variabile.*

*Cerambyx variabilis.* Linn. ſyſt. Nat. 2. 635. 75. Fn. Sv. 669.
*Callidium dilatatum.* Payk. Fn. Sv. 3. 91. 12.
Degeer Inſ. 5. 89. 25.
Oliv. Inſ. 70. tab. 6. fig. 65.
Friſh. Inſ. 12. tab. 6. fig. 3. 4.
Schaeff. Icon. tab. 4. fig. 12.
Panz. Fn. Germ. 70. tab. 6.
Habitat in Europae nemoribus.

10. C. thorace holoſericeo, cinereo, elytris teſtaceis: punctis eleuatis, rubris, ſcutello niueo. Ent. ſyſt. 2. 322. 17. * — *ſericeum.*

*Callidium holoſericeum.* Roſſ. Fn. Etr. 1. 153. 382. tab. 1. fig. 6.
Habitat in Barbaria. Dom. Prof. Vahl.

11. C. thorace elytrisque cinereo-villoſis, antennarum articulis baſi ſcutelloque niueis. Ent. ſyſt. ſuppl. 149. 17. * — *concolor.*

Habitat - - -

12. C. thorace tuberculato, obſcure, rufo: linea dorſali nigra, elytris depreſſis, cineraſcentibus: lineis duabus eleuatis, atris. Ent. ſyſt. 2. 322. 18. * — *tenebroſum*

Oliv. Inſ. 67. tab. 18. fig. 139.
Habitat in Cayenna. Muſ. Dom. Olivier.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*variolosum* 23. C. thorace subtuberculato, nigrum, elytris variolosis: sutura acuminata. Ent. syst. suppl. 149. 18. *

Habitat Tranquebariae. Mus. D. Lund.

*rusticum.* 24. C. thorace nudo, corpore lurido, antennis breuibus. Ent. syst. 2. 322. 19.

Linn. syst. Nat. 2. 634. 67. Fn. Sv. 666.

Oliv. Ins. 70. tab. 3. fig. 39.

Schaeff. Icon. tab. 63. fig. 6.

Panz. Fn. Germ. 70. tab. 8.

Sulz. Ins. tab. 4. fig. 29.

Hist. Ins. tab. 13. fig. 1.

Payk. Fn. Sv. 3. 80. 1.

Degeer Ins. 5. 83. 20.

Habitat in Europae syluis.

*vittatum.* 25. C. thorace rotundato, nudo, elytris ferrugineis vitta media nigra, antennis corpore longioribus. Ent. syst. 2. 323. 20. ♀

Habitat in Guinea. Dom. Ilsert.

*scutellare.* 26. C. thorace tuberculato, nigrum, elytris punctatis, scutello albo, antennis mediocribus, ferrugineis.

Habitat in Amboina. Dom. Billardiere.

Duplo minus C. rustico. Caput nigrum, antennis ferrugineis, longitudine corporis. Thorax punctatus, subtuberculatus. Scutellum villosum album. Elytra punctata, nigra. Pedes subferruginei, femoribus clauatis.

*agreste.* 27. C. thorace nudo, nigrum, elytris striatis, fuscis antennis breuibus. Ent. syst. 2. 323. 21. *

var. striati

Habitat Halae Saxonum. Dom. Hybner.

*stigma.* 28. C. thorace punctato, corpore atro, elytris laeuibus: stigmate albo. Ent. syst. 2. 323. 22.

*Cerambyx stigma.* Linn. syst. Nat. 2. 625. 72.

O

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inſ. 70. tab. 2. fig. 21.
Voet. Coleopt. 2. tab. 25. fig. 134. 135.
Degeer Inſ. 5. 109. 22. tab. 14. fig. 13.
Habitat in America.

29. C. thorace hirto, fuſcum, antennarum articulis baſi rufis. Ent. ſyſt. 2. 323. 23. * *fugax.*
Oliv. Inſ. 70. tab. . fig. .
Habitat in Gallo prouincia. Muſ. Dom. Olivier.

30. C. thorace nudo, corpore fuſco, antennis longis flaueſcentibus. Ent. ſyſt. 2. 323. 24. * *pygmaeum*
Habitat in Anglia, Italia. *VI. 30*

31. C. thorace mutico, nudo, atrum, nitidum, coleoptris faſcia interrupta, rubra. Ent. ſyſt. 2. 324. 25. * *equeſtre.*
Habitat in Cayenna. Dom. v. Rohr.

32. C. thorace mutico, fuluo, nigrum, antennis longiſſimis. Ent. ſyſt. 2. 324. 26. * *fuluicolle.*
Habitat in Surinam. Muſ. Dom. Boſc.

33. C. thorace rotundato, hirto, elytris acuminatis, pallide-teſtaceis. Ent. ſyſt. 2. 324. 27. * *hirtum.*
Oliv. Inſ. 70. tab. 5. fig. 62.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

34. C. thorace rotundato, pubeſcente, teſtaceo, elytris vireſcentibus, baſi teſtaceis, Ent. ſyſt. 2. 324. 28. * *pubeſcens.*
Oliv. Inſ. 70. tab. 6. fig. 75.
Habitat ad Cap. Bon. Spei. Muſ. Dom. Olivier.

35. C. thorace rotundato, ſubtus, vtrinque macula tomentoſa, ferruginea, antennis longis, barbatis. Ent. ſyſt. 2. 324. 29. * *barbatum.*
Oliv. Inſ. 70. tab. 4. fig. 41.
Habitat Tranquebariae. Dr. Koenig.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*mixtum.* 36. C. thorace rotundato ferrugineo, cinereo-lineato, elytris teſtaceis: macula grifeo-fuſca. Ent. ſyſt. ſuppl. 149. 29. *

*Callidium griſeum.* Oliv. Inſ. 70. tab. fig.

Habitat in Gallia. Muſ. D. Boſc.

*griſeum* 37. C. thorace rotundato pubeſcente, corpore obſcure cinereo, antennis breuibus. Ent. ſyſt. 2. 325. 20. *

Habitat in Barbaria. Muſ. D. Desfontaines.

*compreſſum.* 38. C. thorace laeui, obſcure, nigrum, antennis longis pedibusque teſtaceis, femoribus compreſſis Ent. ſyſt. 2. 325. 31. *

Oliv. Inſ. 70. tab. 4. fig. 44.

Habitat in Siam. Muſ. Dom. Banks.

*variegatum.* 39. thoracis dorſo glabro, atro: lineis quatuor albis, elytris flauo-irroratis. Ent. ſyſt. 2. 325. 32.

Oliv. Inſ. 70. tab. 5. fig. 58.

Habitat in noua Zelandia. Muſ. D. Banks.

*lineatum.* 40. C. thorace lineis duabus, elytris quatuor albis: mediis coëuntibus abbreuiatis. Ent. ſyſt. 2. 325. 33. *

Oliv. Inſ. 70. tab. 4. fig. 50.

Habitat in noua Hollandia. Muſ. D. Banks.

*ſulcatum.* 41. C. thorace tomentoſo, cinereo, elytris albo nigroque ſtriatis. Ent. ſyſt. 2. 326. 34. *

Oliv. Inſ. 70. tab. 4. fig. 48.

Habitat in noua Zelandia. Muſ. D. Banks.

*ſanguineum.* 42. C. thorace ſubtuberculato elytrisque ſanguineis, antennis mediocribus. Ent. ſyſt. 2. 326. 35. *

*Cerambyx ſanguineus.* Linn. ſyſt. Nat. 2. 636. 80.

Oliv. Inſ. 70. tab. tab. 1. fig. 1.

Geoff. Inſ. 1. 270. 21.

Vol. 26 Schaeff. Icon. tab. 64. fig. 1.

Panz. Fn. Germ. 70. tab. 9.

Voet

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Voet. Coleopt. 2. tab. 23. fig. 122.
Payk. Fn. Germ. 3. 39. 10.
Degeer Inſ. 5. 92. 29.
Habitat in Europae ſyluis.

3. C. thorace ſubtuberculato, corpore teſtaceo, antennis mediocribus, fuſcis. Ent. ſyſt. 2. 326. 36. * *teſtaceum.*
*Cerambyx teſtaceus.* Linn. ſyſt. Nat. 2. 635. 75. Fn. Sv. 670.
Geoff. Inſ. 1. 218. 18.
Oliv. Inſ. 70. tab. 1. fig. 11.
Schaeff. Icon. tab. 64. fig. 6.
Habitat in Europae ſyluis.

4. C. thorace ſubtuberculato, ferrugineo, elytris teſtaceis, antennis mediocribus, villoſis, femoribus clauatis. Ent. ſyſt. 2. 327. 37. * *pilicorne.*
Habitat in Americae Inſulis. Dr. Iſert.

5. C. thorace ſubtuberculato, teſtaceum, elytris apice violaceis. Ent. ſyſt. 2. 327. 38. * *praeuſtum*
Habitat in Italia. Dr. Allioni.
Feminam C. fennici credit Dom. de Paykull, at vix rite.

6. C. thorace tuberculato, villoſo, nigrum, elytris rubris: macula apiceque violaceis. Ent. ſyſt. 2. 327. 89. * *lignerum.*
Habitat in Marylandia. Dr. Schulz.

7. C. thorace rotundato, ferrugineum, elytris pubeſcentibus, aeneo-micantibus. *micans.*
Habitat in Guinea. Dom. Heſſe. Muſ. D. de Seheſtedt.
Statura parua praecedentium. Antennae corpore longiores, ferrugineae. Caput et thorax ferruginea, nitida, immaculata. Elytra ſubſtriata, villoſa, aeneo valde micantia. Corpus ferrugineum, pedum geniculis nigris.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*flavipes.* 48. C. thorace rotundato, pubescente, ferrugineum, pedibus testaceis. Ent. syst. 2. 327. 41. *
Habitat ad Cap. Bon. Spei. Mus. D. Lund.

*luridum.* 49. C. thorace subtuberculato, nudo, nigrum, elytris laeuibus, testaceis.
*Cerambyx luridus.* Linn. syst. Nat. 2. 634. 68.
Panz. Fn. Germ. 70. tab. 10.
Oliv. Inf. 70. tab. 7. fig. 78.
Payk. Fn Sv. 3. 87. 8.
Habitat in Europae syluis.

*fuscum.* 50. thorace subtuberculato, canaliculato, nudo, nigrum, elytris striatis, obscure-testaceis, antennis mediocribus. Ent. syst. 2. 328. 42. *
Habitat Halae Saxonum. Dom. Hybner.

*4maculatum.* 51. C. thorace villoso, fuscum, elytris maculis duabus ferrugineis, antennis longioribus. Ent. syst. 2. 328. 43. *
Habitat in Insula Guadeloupe. Mus. D. Lund.

*melancholicum.* 52. C. thorace plano, nudo: macula dorsali, fulua fuscum, pedibus testaceis, antennis breuibus. Ent. syst. suppl. 149. 43. *
Habitat in Germaniae Quercu. Dom. Hybner.

*triste.* 53. C. thorace subtuberculato, nudo, elytris substriatis, corpore lurido, immaculato. Ent. syst. 2. 328. 44. *

var. rustica ~~*Callidium aruense*~~ Panz. Naturf. 24. 30. 21. tab. 1. fig. 41.
~~Panz. Fn. Germ. 70. tab. 11.~~
Habitat in Saxoniae Quercu. Dom. Hybner.

*vaticum.* 54. C. thorace laeui, obscure, ferrugineo, elytris testaceis. Ent. syst. 2. 328. 45. *
Habitat in Africa aequinoctiali. Mus. D. Banks.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

55. C. thorace ſubuilloſo, nigro: lineis quatuor albis, intermediis abbreuiatis, antennis breuibus. Ent. ſyſt. 2. 328. 46. * *hafniense.*
*Cerambyx liciatus.* Linn. ſyſt. Nat. 2. 636. 78.
Oliv. Inſ. 70. tab. 1. fig. 8.
Panz. Naturf. 24. tab. 1. fig. 38. VI. 27
Panz. Fn Germ. 70. tab. 12.
Herbſt. Arch. tab. 26. fig. 18.
Payk. Fn. Sv. 3. 93. 15.
Habitat in Europa boreali.

56. C. thorace laeui, nitido, elytris violaceis, tibiis ruſis, antennis breuibus. Ent. ſyſt. 2. 329. 47. * *ruſipes.*
Oliv. Inſ. 70. tab. 6. fig. 66.
Habitat in Germania. Dom. de Hattorff.

57. C. thorace glabro, corpore nigro, elytris ſtriatis, antennis breuibus. Ent. ſyſt. 2. 329. 48. * *ſtriatum.*
*Cerambyx ſtriatus.* Linn. ſyſt. Nat. 2. 635. 73.
Fn. Sv. 668.
Oliv. Inſ. 70. tab. 2. fig. 24.
Herbſt. Arch. tab. 26. fig. 13. VI. 28
Panz. Fn. Germ. 70. tab. 13.
Degeer Inſ. 5. 90. 26.
Payk. Fn. Sv. 3. 83. 4.
Habitat in Europae ſyluis.

58. C. nigrum, thorace atro, nitido, antennis breuibus. Ent. ſyſt. ſuppl. 151. 48. * *aulicum.*
Panz. Fn. Germ. 70. tab. 14. VI. 28
Payk. Fn. Sv. 3. 82. 3. var. ſimile
Habitat in Daniae nemoribus.

59. C. thorace rotundato, pubeſcente, ferrugineo, elytrorum ſutura acuminata, corpore teſtaceo, antennis longis. Ent. ſyſt. ſuppl. 150. 19. * *ſtrepens.*
Habitat in Tanger. Dom. Schousboe.

60.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bicolor.* 60. C. thorace glabro, flaueſcente, capite elytrisque vireſcentibus, antennis mediocribus, nigris. Ent. ſyſt. 2. 329. 49. *

Oliv. Inſ. 70. tab. 1. fig. 4.

Habitat in America meridionali.

*lynceum.* 61. C. thorace rotundato, ſubſpinoſo, villoſo, nigrum, elytris macula ſubdidyma, fulua. Ent. ſyſt. 2. 329. 50. *

*Cerambyx lynceus.* Oliv. Inſ. 67. tab. 12. fig. 97.

Habitat ad Cap. Bon. Spei. Dom. Yeats.

*cyaneum.* 62. C. thorace plano, tuberculato, villoſum, violaceum, abdominis penultimo ſegmento ferrugineo, antennis mediocribus. Ent. ſyſt. 2. 330. 52. *

Habitat in Italia.

*puſillum.* 63. C. thorace rotundato, nigrum, antennis tibiisque teſtaceis. Ent. ſyſt. 2. 330. 53. *

VII. 96. Habitat in Germania. Dom. Schneider.

Nimis affine videtur C. pygmaeo.

*2maculatum.* 64. C. thorace rotundato, villoſo, fuſcum, elytris maculis duabus ferrugineis. Ent. ſyſt. 2. 330. 54. *

Oliv. Inſ. 70. tab. 5. fig. 61.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*flauum.* 65. C. thorace ſubrotundo, corpore flauo, femoribus clauatis, antennis mediocribus. Ent. ſyſt. 2. 330. 55. *

Habitat in America. Muſ. Tottianum.

*vndatum.* 66. C. thorace tuberculato, elytris nigris: faſciis duabus vndatis albis, antennis breuibus. Ent. ſyſt. 2. 331. 56. *

VI. 28 *Cerambyx vndatus.* Linn. ſyſt. Nat. 2. 636. 79. Fn. Sv. 675.

Panz. Fn. Germ. 70. tab. 15.

Oliv.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 78. tab. 3. fig. 36.
Voet. Col. 2. tab. 19. fig. 87.
Degeer Inf. 5. 91. 28.
Payk. Fn. Sv. 3. 92. 14.
Habitat in Europae truncis.

67. C. thorace rotundato, nigrum, elytris faſciis tribus cinereis. Ent. ſyſt. ſuppl. 152. 56. * *terminans*
Habitat in America boreali. D. Herſchel.

68. C. thorace rotundato, elytris liuidis: faſciis tribus fuſcis, antennis breuibus. Ent. ſyſt. 2. 331. 57. * *coloretum.*
Oliv. Inf. 70. tab. 6. fig. 67.
Habitat in Carolina.

69. C. thorace rotundato rufo, elytris nigris baſi rufis: faſciis duabus albis. Ent. ſyſt. 2. 331. 58. * *varium.*
Oliv. Inf. 70. tab. 5. fig. 55.
Habitat in America boreali. D. Yeats.

## 143. CLYTVS.

*Palpi* quatuor breues, extrorſum craſſiores.
*Maxilla* bifida.
*Ligula* breuis, truncata, integra.
*Antennae* ſetaceae.

1. C. thorace rotundato flauo faſciato, elytris faſciis ſeptem *flexuoſus.*

---

Clyti corpus oblongum, cylindricum, ſaepius pubeſcens, immarginatum, tardum; *capite* ouato, decliui, inſerto; *oculis* magnis, reniformibus, marginalibus; *antennis* thorace longioribus, approximatis, in ſinu oculorum inſertis; *thorace* globoſo, latitudine elytrorum; *ſcutello* rotundato; elytris rigidis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, curſoriis; *femoribus* compreſſis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

septem flauis: anticis antrorsum posticis, retrorsum arcuatis.
*Callidium flexuosum.* Ent. syst. 2. 331. 59. *
*Leptura Robiniae.* Forst. Cent. 1. 43.
*Leptura picta.* Drury Inf. 1. tab. 41. fig. 2.
Oliv. Inf. 70. tab. 6. fig. 76.
Voet. Col. tab. 91. fig. 91.
Habitat in Americae Robinia.

*minutus.* 2. C. testaceus, elytris fascia abbreuiata alba. Ent. syst. 2. 332. 60. *
Oliv. Inf. 70. tab. 5. fig. 56.
Habitat in noua Zelandia. Mus. D. Banks.

*floralis.* 3. C. thorace globoso albo-fasciato, elytris nigris: fasciis quinque albis; secunda tertiaque lunatis. Ent. syst. 2. 332. 61.
*Callidium fasciatum.* Herbst. Arch. tab. 26 f. 17.
*Cerambyx floralis.* Pall. Iter. 2. 724. 63.
Oliv. Inf. 70. tab. 5. fig. 53.
Voet. Col. tab. 19. fig. 93.
Panz. Fn. Germ. 70. tab. 16.
Habitat in Italia. Dr. Allioni.

*fulminans.* 4. C. thorace globoso maculato, elytris nigris: fasciis vndato-angulatis albis. Ent. syst. 2. 332. 62. *
Oliv. Inf. 70. tab. 5. fig. 63.
Habitat in America ~~meridionali~~. Mus. D. Banks.

*semipunctatus.* 5. C. thorace globoso nigro: punctis tribus albis, elytris nigris: punctis duobus lunulisque tribus albis. Ent. syst. suppl. 151. 62. *
Habitat in Germania.

*atomarius* 6. C. thorace globoso cinereo maculato, elytris punctis strigaque postica flexuosa albidis. Ent. syst. 2. 332. 63. *
Habitat Halae Saxonum. D. Hybner.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

7. C. thorace globoſo virescente: maculis duabus ferrugineis, elytris bidentatis nigris: ſtrigis flauis baſi rufis. Ent. ſyſt. ſuppl. 152. 63. *luſcus.*

Habitat in America boreali. D. v. Rohr.

8. C. thorace rotundato, elytris faſciis quatuor flauis: prima interrupta, reliquis retrorſum arcuatis. Ent. ſyſt. 2. 333. 64. *arcuatus.*

*Leptura arcuata.* Linn. ſyſt. Nat. 2. 640. 21. Fn. Sv. 696.

Oliv. Inſ. 70. tab. 2. fig. 16.

Herbſt. Arch. tab. 26. fig. 14. I. 45.

Geoff. Inſ. 1. 212. 10.

Panz Fn. Germ. 4. tab. 14.

Schaeff. Icon. tab. 38. fig. 6.

~~tab. 107. fig. 2. 3.~~

Payk. Fn. Sv. 3. 95. 17.

Habitat in Europae hortis.

*Os* maxillis palpisque. *Palpi* quatuor, breues, aequales extrorſum craſſiores: *anteriores* vix longiores, quadriarticulati, articulis ſenſim craſſioribus, vltimo breui truncato, adhaerentes maxillae dorſo; *poſteriores* triarticulati: articulo vltimo breui, craſſiori, truncato, adnati ligulae baſi ſuperiori. *Mandibula* breuis, cornea, craſſa, edentula, apice acuta. *Maxilla* breuis, cornea, compreſſa, bifida: lacinia interiore breui, acuta, exteriore longiore, apice craſſiore, truncata. *Ligula* breuis, membranacea, truncata, integra. *Labium* transuerſum, anguſtum, corneum, late ſubemarginatum.

9. C. thorace rotundato, nigro, elytris nigris: faſciis flauis, ſecunda antrorſum arcuata, pedibus totis ferrugineis. Ent. ſyſt. 2. 333. 65. *arietis.*

~~*Leptura arietis.* Linn. ſyſt.~~ Nat. 2. 640. 23. Fn. ~~Sv. 695.~~

Oliv.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 70. tab. 2. fig. 20.
Geoff. Inf. 1. 214. 11.
Herbſt. Arch. tab. 26. fig. 16.
Petiv. Gazoph. tab. 36. fig. 6.
Schaeff. Icon. tab. 38. fig. 7.
tab. 107. fig. 3.
I. 45
Panz. Fn. Germ. 4. tab. 15.
Friſh. Inſ. 12. tab. 3. fig. 5.
Habitat in Europae hortis.

*gazella.* 10. C. thorace rotundato, nigro, elytris nigris: faſciis flauis, ſecunda antrorſum arcuata, pedibus ferrugineis: femoribus nigris. Ent. ſyſt. 2. 333. 66. *
I. 45
Habitat Kiliae.

*feſtiuus.* 11. C. thorace rotundato, niger, elytris maculis tribus flauis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et ſumma affinitas C. laeti. Antennae extrorſum fere incraſſatae, nigrae. Caput et thorax nigra, immaculata. Elytra truncata, ſcutello maculisque tribus laete flauis: baſeos duabus approximatis, maioribus ad ſuturam, poſteriore transuerſa. Corpus nigrum, pectore vtrinque macula, abdomine faſciis duabus, baſeos flauis. Pedes ſubelongati, nigri.

*laeſus.* 12. C. thorace rufo, elytris truncatis, atris: maculis tribus flauis, antennis pedibusque ferrugineis.
Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.
Minor C. arietis. Antennae breues, ferrugineae. Caput rufum, mandibulis, atris. Thorax rufus, baſi interdum flauus. Elytra laeuia, nigra, ſtriga abbreuiata, per ſcutellum ducta, macula lineari obliqua, altera approximata, triangulari ad

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ad basin strigaque pone medium flauis. Corpus villoso-flauum, segmentis abdominalibus apice atris. Pedes ferruginei.

13. C. ater, elytris medio coarctatis: maculis duabus albis. *coarctatus*

Oliv. Inf. 70. tab. 6. fig. 71.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

C. plebejo paullo maior. Antennae breues, compressae. Caput et thorax pilosa, atra. Elytra nitidiora, pilosa, laeuia, atra: maculis duabus albis, altera lineari obliqua ad basin, altera maiore in medio. Corpus atrum, abdomine basi pectoreque argenteo nitidulis.

14. C. thorace rufo-maculato, niger, elytris margaritaceis basi nigris. *margaritaceus.*

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura et magnitudo C. plebeii. Caput cum antennis atrum. Thorax globosus, ater, vtrinque maculis duabus magnis, rufis. Elytra plana, laeuissima, margaritacea, nitida, basi margineque exteriori atris. Corpus atrum, femoribus clauatis.

15. C. thorace immaculato, elytris nigris: strigis tribus punctoque humerali albis. Ent. syst. 2. 334. 67. * *plebejus.*

~~*Callidium pallidum.* Oliv. Inf. 70. tab. 6. f. 72.~~

~~*Leptura rustica.* Linn. syst. Nat. 2. 639. 17. Fn. Sv. 692.~~

~~Geoff. Inf. 1. 215. 12.~~ VII. 96.

Petiv. Gazoph. tab. 63. fig. 5.

Schaeff. Icon. tab. 2. fig. 7.

Payk. Fn. Sv. 3. 99. 21.

Habitat in Europae nemoribus.

16.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*massiliensis* 16. C. thorace immaculato, elytris nigris: strigis tribus albis; anteriore incurua. Ent. syst. 2. 334. 68. *

*Leptura massiliensis.* Linn. syst. Nat. 2. 640. 19.
VII. 36. Geoff. Inf. 1. 215. 12.
Oliv. Inf. 70. tab. 6. fig. 70.
Habitat in Germania.

*mucronatus.* 17. C. thorace maculato, elytris mucronatis nigris: strigis tribus arcuatis flauis basi ferrugineis. Ent. syst. 2. 334. 69. *
Oliv. Inf. 70. tab. 3. fig. 34.
Habitat in America. D. Solander.

*angulatus.* 18. C. thorace muricato, elytris mucronatis: maculis tribus dorsalibus punctoque medio flauis. Ent. syst. 2. 335. 70. *
Habitat - - Mus Britt.

*erythrocephalus.* 19. C. thorace subspinoso, elytris bidentatis fuscis: strigis quatuor flauis, femoribus clauatis compressis. Ent. syst. 2. 335. 71. *
Oliv. Inf. 70. tab. 5. fig. 60.
Habitat in America. Dr. Solander.

*similis.* 20. C. thorace nigro: tuberculis transuersis, elytris testaceis: basi margineque omni cinereis. Ent. syst. suppl. 152. 7. *
Habitat in Cayenna. D. Cuvier.

*detritus.* 21. C. thorace flauo fasciato, elytris nigris: fasciis quinque flauis, pedibus ferrugineis. Ent. syst. 2. 335. 72. *
*Leptura detrita.* Linn. syst. Nat. 2. 646. 20.
VIII. 31. Fn. Sv. 694.
Oliv. Inf. 70. tab. 2. fig. 17.
Schaeff. Icon. tab. 38. fig. 9.
tab. 64. fig. 3.

Geoff.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 216. 13.
Habitat in Europae hortis.

22. C. niger, elytris glaucis baſi nigro-maculatis. *glaucus.*
Ent. ſyſt. 2. 336. 73. *
Oliv. Inf. 70. tab. 6. fig. 68.
Habitat in India orientali. Muſ. D. Banks.

23. C. ater, thorace faſciis duabus, elytris quatuor *6faſciatus.*
flauis. Ent. ſyſt. 2. 336. 74. *
Habitat - - Muſ. Britt.

24. C. thorace ferrugineo, elytris nigris: faſciis *trifaſcia-*
tribus albis; prima annulari. Ent. ſyſt. 2. 336. *tus.*
75. *
Oliv. Inf. 70. tab. 5. fig. 59.
Habitat in Luſitania. Dr. Grey.

25. C. thorace nigro trimaculato, elytris rotundatis *Verbaſci.*
ſubuireſcentibus: faſciis tribus abbreuiatis nigris, prima lunari. Ent. ſyſt. 2. 336. 76. *
~~*Leptura Verbaſci.* Linn. ſyſt. Nat.~~ 2. 640. 22.
Oliv. Inf. 70. tab. 1. fig. 5.
~~tab. 6. fig. 15.~~
~~Sulz. Hiſt. Inf. tab. 5. fig. 12.~~
Panz. Fn. Germ. 75. tab 17.
Herbſt. Arch. tab. 26. fig. 19.
Habitat in Europae auſtralioris Verbaſco.

26. C. thorace rotundato: faſcia nigra, elytris vire- *ornatus.*
ſcentibus: faſciis tribus integris nigris, prima annulari. Ent. ſyſt. 2. 336. 77. *
*Callidium ornatum.* Herbſt. Arch. tab. 26. f. 16.
*Stenocorus duplex.* Scopol. Fl. Inf. tab. 20. fig. 5.
Panz. Fn. Germ. 70. tab. 18.
Sulz. Hiſt. Inf. tab. 5. fig. 12.
Habitat in Germania. D. Smidt.

27. C. thorace nigro-maculato, elytris bidentatis *bidens.*
vireſcentibus: annulo macula ſtrigaque nigris.

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Sumatra. D. Daldorff.

Statura omnino et magnitudo C. ornati. Antennae breues, fuſcae, baſi ferrugineae. Caput vireſcens. Thorax rotundatus, vireſcens: maculis nigris. Elytra apice bidentata, vireſcentia, annulo baſeos ſtriga media, cum annulo coëunte maculaque orbiculata apicis atris. Pedes nigri.

*annulus.* 28. C. vireſcens, elytris faſciis tribus nigris: tertia annulari.

Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

Statura omnino C. ornati. Caput vireſcens, antennis breuibus, ferrugineis. Thorax rotundatus, vireſcens, immaculatus. Elytra vireſcentia: faſciis tribus nigris; vltima annulari. Pedes pallide teſtacei.

*4punctatus.* 29. C. vireſcens, elytris punctis quatuor. Ent. ſyſt. 2. 337. 78. ♂

Geoff. Inf. 1. 213. 8.

Panz. Fn. Germ. 70. tab. 19.

VI. 30

Habitat in Gallia haud infrequens.

*annularis.* 30. C. thorace nigro-maculato, elytris bidentatis ſubuireſcentibus: faſciis tribus nigris, prima annulari. Ent. ſyſt. 2. 337. 79. ♂

*Callidium annulare.* Oliv. Inſ. 70. tab. 6. f. 74.

Habitat in Siam, Muſ. D. Banks. in India. Muſ. D. Lund.

*aegyptiacus.* 31. C. thorace ferrugineo, elytris cineraſcentibus: faſciis tribus fuſcis; prima annulari. Ent. ſyſt. 2. 337. 80. ♂

Habitat in Oriente. D. Prof. Forſkahl.

*myſticus.* 32. C. niger, elytris fuſcis: faſciis apiceque cinereis baſi rufis. Ent. ſyſt. 2. 337. 81.

*Leptura myſtica.* Linn. ſyſt. Nat. 2. 639. 18.

VII-37. Fn. Sv. 693.

Oliv

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 70. tab. 1. fig. 14.
Geoff. Inf. 1. 217. 15.
Schaeff. Icon. tab. 2. fig. 9.
Voet. Coleopt. 2. tab. 26. fig. 8.
Payk. Fn. Sv. 3. 98. 20.
Degeer Inf. 5. 82. 19.
Habitat in Europae hortis.

33. C. virefcens, antennis nigris, pedibus teftaceis. *virens.*
Ent. fyft. 2. 338. 82. *
Habitat in Barbaria. Muf. Dom. Desfontaines.

34. C. niger, elytris cinereo-faſciatis, baſi bituber- *gibboſus.*
culatis, apice acuminata. Ent. fyft. 2. 338. 83. *
Oliv. Inf. 70. tab. 2. fig. 18.
Habitat in Italia. D. Daldorff.

35. C. thorace brunneo, elytris nigris: faſcia media *unifaſcia-*
niuea, baſi brunneis. Ent. fyft. 2. 338. 84. * *tus.*
Oliv. Inf. 70. tab. 1. fig. 12.
Habitat in Gallo prouincia. Dom. Olivier.

36. C. ater, elytris ſtriga obliqua alba. Ent. fyft. 2. *picipes.*
338. 85. *
Oliv. Inf. 70. tab. 4. fig. 43.
Habitat in America. Muf. Dr. Hunter.

37. C. ferrugineus, elytris aureis: ſtrigis duabus *piniadeus.*
albis.
Habitat in Carolina. Muf. Dom. Boſc.
Statura et magnitudo C. Alni. Antennae corpore paullo longiores, ferrugineae. Caput et thorax globoſus ferruginea, immaculata. Elytra aurea, nitida, baſi linea impreſſa ſtrigisque duabus albis, quae tamen vix futuram attingunt. Corpus ferrugineum, femoribus clauatis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Alni.* 38. C. niger, elytris faſciis duabis albis, elytrorum baſi antennis tibiisque ferrugineis. Ent. ſyſt. 2. 338. 86. *

*Leptura Alni.* Linn. ſyſt. Nat. 2. 689. 19.
Oliv. Inſ. 70. tab. 3. fig. 7.
Herbſt. Arch. tab. 26. fig. 21.
Scopol. Flor. Inſ. tab. 20. fig. 3.
Panz. Fn. Germ. 70. tab. 20.
Payk. Fn. Sv. 3. 100. 22.
Habitat in Anglia, Germania.

## 144. LEPTVRA.

*Palpi* filiformes.
*Maxilla* vnidentata, apice ſetoſa.
*Ligula* membranacea, bifida.
*Antennae* ſetaceae.

*vnipunctata.* 1. L. nigra, elytris rufis: puncto medio nigro. Ent. ſyſt. 2. 339. 1.
Oliv. Inſ. 73. tab. 1. fig. 9.
Habitat Dresdae. Dom. Romanus.

*haſtata.* 2. L. atra, elytris rubris: apice ſuturaque media nigris. Ent. ſyſt. 2 339. 2. *
Oliv. Inſ. 73. tab. 1. fig. 5.
Geoff. Inſ. 1. 226. 6.
Sulz. Inſ. tab. 5. fig. 11.
Habitat in Europa auſtraliori.

*2punctata.* 3. L. nigra, villoſa, elytris luteis: ſutura puncto media apiceque nigris. Ent. ſyſt. 2. 340. 3.
Habitat in Sibiria. Muſ. D. Banks.

---

*Leptarae* corpus oblongum, antice poſticeque attenuatum, ſaepius tomentoſum, immarginatum, tardum; *capite* exſerto ouato; *ocelis* globoſis, prominulis, marginalibus; *antennis* thorace longioribus, approximatis, inter oculos inſertis *thorace* rotundato, anguſto, antice anguſtiore, inermi *ſcutello* paruo, triangulo; *elytris* rigidis, lorniatis, ſaepe emarginatis, longitudine abdominis; *pedibus* longiuſculis curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. L. thorace villoſo aureo, elytris teſtaceis apice ni- *tomentoſa.*
gris. Ent. ſyſt. 2. 340. 4. *
Geoff. Inſ. 1. 227. 8.
Payk. Fn. Sv. 3. 108. 9.
Degeer Inſ. 5. 136. 13.
Habitat in Gallia. Muſ. D. de Seheſtedt.

. L. thorace laeui, nigra, elytris teſtaceis, apice ni- *maculicor-*
gris, antennarum articulis, baſi flauis. Ent. ſyſt. *nis.*
ſuppl. 152. 5. *
Payk. Fn. Sv. 3. 111. 12.
Degeer Inſ. 5. 139. 16.
Habitat in Germania.

. L. nigra, elytris rubeſcentibus liuidisue: ſutura *melanura.*
apiceque nigris. Ent. ſyſt. 2. 340. 5.
Linn. ſyſt. Nat. 2. 637. 2.
Geoff. Inſ. 1. 226. 7. tab. 4. fig. 1.
Petiv. Gazoph. tab. 30. fig. 5.
~~Degeer Inſ. 5. 137. 14.~~ VI. 31.
~~Schaeff. Icon. tab. 39. fig.~~ 4.
Panz. Fn. Germ. 69. tab. 19.
Payk. Fn. Sv. 3. 110. 11.
Habitat in Europae floribus frequens.

. L. nigra, elytris pedibusque liuidis, elytris ſutura *laeuis.*
apiceque nigris. Ent. ſyſt. 2. 340. 6. *
Panz. Fn. Germ. 34. tab. 16.
Payk. Fn. Sv. 3. 128. 31. III. 56.
Degeer Inſ. 5. 139. 17.
Habitat Kiliae.

. L. nigra, elytris teſtaceis, immaculatis, pedibus *liuida.*
nigris. Ent. ſyſt. 2. 340. 7. *
Herbſt. Arch. tab. 26. fig. 23.
Habitat in floribus Kilonienſibus. Muſ. D. Lund.

. L. nigra, elytris teſtaceis: vitta nigricante. Ent. *ſtrigilata.*
ſyſt. 2. 341. 8. *



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 112. 13.
Habitat in Suecia. Dom. Com. de Souza.

*emargina-ta.* 10. L. atra, elytris, purpureis, apice emarginatis nigris. Ent. syst. 2. 341. 9. *
Habitat in Cayenna. Dr. Hunter.

*sanguino-lenta.* 11. L. nigra, elytris sanguineis. Ent. syst. 2. 341. 10
*Leptura sanguinolenta.* Linn. syst. Nat. 2. 638 4. Fn. Sv. 679.
*Leptura variabilis.* Payk. Fn. Sv. 3. 109. 10.
Panz. Fn. Germ. 69. tab. 8.
Vl. 91. Oliv. Inf. 73. tab. 2. fig. 13.
~~Schaeff. Icon. tab.~~ 39. fig. 9.
Degeer Inf. 5. 137. 14.
Habitat in Europae floribus.

*cincta.* 12. L. atra, elytris punctatis, testaceis: margine tot atro.
Habitat in Austria. Dom. de Meyerle.
Statura omnino L. sanguinolentae, at minor, cu
Vl. 91. 32 ius marem olim credidi. Caput, thorax, corpu atra, immaculata. Elytra punctata, testacea margine exteriori, late atro. Abdomen subtus pilis aureis, nitens.

*meridiana* 13. L. thorace subspinoso, elytris fastigiatis, pector nitente. Ent. syst. 2. 341. 11. *
~~*Stenocorus dispar.* Panz. Fn. Germ. 17. tab. 1. 2~~
IL. 40. IV. 64. *Cerambyx meridianus.* Linn. syst. Nat. 2. 63 47. Fn. Sv. 648.
Schaeff. Icon. tab. 82. fig. 4.
tab. 279. fig. 3.
Payk. Fn. Sv. 3. 104. 4.
Degeer Inf. 5. 130. 5.
Habitat in Europae floribus.
Mas nigricans, femina testacea.

*thoracica.* 14. L. atra, thorace rufo.

*Lept*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Leptura thoracica.* Creutz. Inſ. 1. 125. 16. tab. 3. fig. 28.
Payk. Fn. Sv. 3. 105. 5.
Habitat in Carolina. Dom. de Meyerle.
Statura L. meridianae, at paullo craſſior. Caput, elytra, corpus nigra, immaculata. Thorax ſolus rufus.

15. L. atra, elytris punctatis, baſi rubris. *canadenſis.*
*Leptura canadenſis.* Oliv. Inſ. 73. tab. fig.
Habitat in Canada. Muſ. Dom. Boſc.
Statura magna L. thoracicae. Caput atrum, articulis intermediis teſtaceis. Thorax rotundatus, punctatus, ater, immaculatus. Elytra punctata, atra, baſi late ruſa. Abdomen atrum, abdomine villis argenteis micante.

16. L. ferruginea, antennis elytris pectoreque fuſcis. *villica.*
Ent. ſyſt. 2. 341. 12.
*Leptura veneſtita.* Linn. ſyſt. Nat. 2. 638. 6.
Payk. Fn. Sv. 3. 113. 14.
Schaeff. Icon. tab. 69. fig. 1.
Habitat in Anglia. Muſ. Britt.

17. L. nigra, thorace elytris tibiisque purpureis. *rubra.*
Ent. ſyſt. 2. 342. 13.
Linn. ſyſt. Nat. 2. 638. 3. Fn. Sv. 500.
Friſh. Inſ. 12. tab. 6. fig. 6.
Payk. Fn. Sv. 3. 107. 8.
Panz. Fn. Germ. 69. tab. 11.
Degeer Inſ. 5. 132. 7.
Sulz. Inſ. tab. 5. fig. 30.
Schaeff. Icon. tab. 39. fig. 2.
Habitat in Europa.

18. L. nigra, elytris teſtaceis, tibiis rufis, thorace poſtice rotundato. Ent. ſyſt. 2. 342. 14. *teſtacea.*
*Leptura teſtacea.* Linn. ſyſt. Nat. 2. 638. 5. Fn. Sv. 680.

*Leptu-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Leptura diſpar.* Payk. Fn. Sv. 3. 107. 8.
Panz. Fn. Germ. 69. tab. 12.
Degeer Inſ. 5. 135. 11.
Schaeff. Icon tab. 39. fig. 3.
Habitat in Europae floribus.
Mas praecedentis videtur. Saepius in copula inueniuntur.

*pubeſcens.* 19. L. nigra, cinereo-villoſa: coſta alarum baſi teſtacea. Ent. ſyſt. 2. 342. 15. *
Payk. Fn. Sv. 3. 106. 7.
Habitat in Suecia. D. de Paykull.

*virens.* 20. L. ſericeo-vireſcens, antennis fuſco viridique variis. Ent. ſyſt. 2. 342. 16.
Linn. ſyſt. Nat. 2. 638. 7. Fn. Sv. 682.
Oliv. Inſ. 73. tab. 2. fig. 111.
Panz. Naturf. 24. 31. 42. tab. 1. fig. 42.
Payk. Fn. Sv. 3. 117. 19.
Panz. Fn. Germ. 69. tab. 13.
Degeer Inſ. 5. 131. 6.
Habitat in Europae arboribus.

*ſmaragdula.* 21. L. ſericeo-vireſcens, antennis pedibusque nigris. Ent. ſyſt. 2. 342. 17. *
Payk. Fn. Sv. 3. 118. 20.
Habitat in Suecia. Dom. de Paykull.

*holoſericea.* 22. L. nigra, cinereo-pubeſcens, elytris emarginatis, tibiis teſtaceis.
Habitat in Auſtria. Dom. de Meyerle.
Maior L. atra et omnino diſtincta. Corpus totum nigrum, cinereo-holoſericeum, capite tamen elytrisque emarginatis, magis nigris. Tibiae omnes baſi teſtaceae.

*ruſipes.* 23. L. villoſa, nigra, elytris rotundatis, pedibus ruſis.
Habitat in Auſtria. Dom. de Meyerle.
Affinis certe L. atrae. Differt tantum, elytris haud emarginatis et pedibus omnibus ruſis. 24.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. L. corpore toto nigro. Ent. ſyſt. 2. 342. 28. * *atra*
Geoff. Inſ. 1. 228. 10.
Oliv. Inſ. 73. tab. 1. fig. 4. *VI. 32.*
Panz. Fn. Germ. 69. tab. 14.
Voet. Col. 2. tab. 29. fig. 9.
Payk Fn. Sv. 3. 125. 26.
Habitat in Europae floribus.

5. L. nigra, abdomine humerisque ferrugineis. Ent. *humeralis.*
ſyſt. 2. 343. 29.
Habitat in Germania. Dom. Schneider. *IV. 65*

6. L. aureo-pubeſcens, elytris ferrugineis: macu- *fugax.*
la apicis flaua. Ent. ſyſt. ſuppl. 153. 19. *
Habitat in America. Dom. Hybner.

7. L. nigra, elytris puncto baſeos lineaque media *exclama-*
flauis. Ent. ſyſt. 2. 343. 20. * *tionis.*
Habitat in Suecia. Dom. de Paykull.
Varietas L. 6 guttatae teſte Dom. de Paykull.

8. L. cinerea, elytris teſtaceis: ſutura nigra. Ent. *ſuturalis.*
ſyſt. 2. 343. 21. *
Habitat Halae Saxonum. Dom. Hybner.

9. L. punctata, nigra, ſcutello albo. Ent. ſyſt. 2. *ſcutellata.*
343. 22. *
Panz. Fn. Germ. 69. tab. 15.
Payk. Fn. Sv. 3. 106. 6. *VI. 33.*
Habitat in Italia. Dr. Allioni.
Thorax et elytra valde punctata.

a. L. laeuis, nigra, elytris ſubacuminatis, ſubtus *Chamomil-*
cinereo-holoſericea. *lae.*
Habitat in Auſtriae Chamomilla. Dom. Scheidler.
Magnitudo et ſtatura L. ſcutellatae, at puncta im- *VI. 35*
preſſa omnino deſunt. Elytra apice in angulum
excurrunt. Corpus ſubtus cinereo-holoſericeum.

1. L. ferruginea, elytris teſtaceis. Ent. ſyſt. 2. *lurida.*
343. 23. *

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Vl. 34 Panz. Fn. Germ. 69. tab. 16.
Habitat in ſyluis Herciniae. Dom. Helwig.

*femorata.* 32. L. nigra, femoribus baſi ruſis. Ent. ſyſt. 2. 343. 24.
Payk. Fn. Sv. 3. 127. 29.
Habitat Halae Saxonum. Dom. Hybner.

*ruficornis.* 33. L. nigra, antennis pedibusque ruſis. Ent. ſyſt. 2. 344. 25. *
Payk. Fn. Sv. 3. 127. 30.
Habitat in Italia. Dr. Allioni.

*marginata* 34. L. nigra, elytrorum margine tibiisque poſticis ruſis. Ent. ſyſt. 2. 344. 26. *
Panz. Fn. Germ. 69. tab. 17.
Vl. 94 Schaeff. Icon. tab. 55. fig. 8.
Habitat in Norwegia. Muſ. D. Lund.

*lupata.* 35. L. atra, elytris macula lunata flaua.
Habitat in Carolina. Muſ. Dom. Boſc.
Media. Caput cum antennis atrum. Thorax pubeſcens, punctatus, ater, poſtice pilis aliquot aureis. Elytra punctata, atra, macula magna baſeos lunata, flaua, apice emarginata. Corpus atrum.

*nigra.* 36. L. elytris attenuatis, corpore nigro, nitido, abdomine rubro. Ent. ſyſt. 2. 344. 27. *
*Leptura nigra.* Linn. ſyſt. Nat. 2. 639. 14. Fn. Sv. 687.
Geoff. Inſ. 1. 227. 9.
Vl. 34. Schaeff. Icon. tab. 39. fig. 7.
Panz Fn. Germ. 69. tab. 18.
Payk. Inſ. Sv. 3. 120. 22.
Degeer Inſ. 5. 144. 24.
Habitat in Europae floribus.

*praeuſta.* 37. L. aureo-pubeſcens, capite elytrorumque apicibus nigris. Ent. ſyſt. 2. 344. 28. *

Panz

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 34. tab 17.
Habitat Halae Saxonum. Dom. Hybner.

38. L. fuſca, elytris nigris: punctis duobus baſeos ferrugineis. Ent. ſyſt. 2. 344. 29. * *4guttata.*
Habitat Halae Saxonum.

39. L. obſcure-aenea, pedibus flauis. Ent. ſyſt. 2. 344. 30. * *roſtrata.*
Habitat in Ammi maius Africae. Dom. Prof. Vahl.

40. L. lutea, thorace lineis duabus, elytris faſciis quatuor nigris. Ent. ſyſt. 2. 345. 31. * *luteicornis*
Habitat in Carolina. Muſ. Britt.

41. L. nigra, elytris teſtaceis: maculis duabus nigris. Ent. ſyſt. 2. 345. 32. * *4maculata*
Oliv. Inſ. 73. tab. 1. fig. 7.
Payk. Fn. Sv. 3. 102. 2.
Habitat in Germania. D. Smidt.

42. L. thorace ſpinoſo, pubeſcente, elytris faſtigiatis, liuidis: taenia obſcura longitudinali ſinuata. *lamed.*
*Stenocorus lamed.* Ent. ſyſt. 2. 293. 2.
*Cerambyx lamed.* Linn. ſyſt. Nat. 2. 630. 46.
Fn. Sv. 649.
~~Geoff. Inſ. 1. 226. 6.~~
Payk. Fn. Sv. 3. 101. 1.
Habitat in Europae floribus.

43. L. nigra, elytris maculis duabus remotis, ferrugineis. Ent. ſyſt. 2. 345. 33. * *4puſtulata.*
Habitat in Suecia. Dom. de Paykull.

44. L. nigra, elytris liuidis: maculis quatuor nigris. Ent. ſyſt. 2. 345. 34. * *8maculata*
~~*Leptura 8maculata.* Linn.~~ ſyſt. Nat. 2. 638. 10.
~~Fn. Sv. 684.~~
~~Schaeff. Elem. tab. 118.~~ fig. 2.
~~Icon. tab. 1. fig.~~ 7.
Habitat in Europae floribus.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*interrogationis.* 45. L. nigra, elytris flauis: linea longitudinali arcuata maculisque quatuor marginalibus nigris. Ent. syst. 2. 345. 35. *

Linn. Fn Sv. 685.

II. 42 Payk. Fn. Sv. 3. 121. 24.

Habitat in Suecia. Dom. de Souze.

*marginella* 46. L. nigricans, elytris futura maculis duabus marginalibus apiceque flauis. Ent. syst. 2. 346. 36. *

Habitat in Italia. Dr. Allioni.

Varietas praecedentis teste Dom. de Paykull.

*6 maculata* 47. L. nigra, elytris testaceis: fasciis tribus dentatis, nigris; anteriore subinterrupta. Ent. syst. 2. 346. 37. *

*Leptura 6 maculata.* Linn. syst. Nat. 2. 638. 11. Fn. Sv. 686.

~~Schaeff. Icon. tab. 6.~~ fig. 9.

Voet. Col. 2. tab. 26. fig. 12.
tab. 21. fig. 108.

VI. 35 Panz. Fn. Germ. 69. tab. 21.

Herbst. Arch. 5. tab. 26. fig. 26.

Payk. Fn. Sv. 3. 123. 25.

Degeer Inf. 5. 133. 9.

Habitat in Europae hortis.

*7 punctata.* 48. L. nigra, thorace testaceo: puncto nigro, coleoptris testaceis: punctis septem. Ent. syst. 2. 346. 38. *

VIII. 33.

Habitat in Hungaria. Dom. Hybner.

*12 maculata* 49. L. nigra, elytris flauis: maculis sex nigris. Ent. syst. 2. 346. 39. *

Habitat in Sibiria. Mus. D. Banks.

*attenuata.* 50. L. elytris attenuatis, fasciis quatuor nigris, pedibus testaceis. Ent. syst. 2. 346. 40.

*Leptura attenuata.* Linn. syst. Nat. 2. 639. 13. Fn. Sv. 688. Schaeff.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Schaeff. Icon. tab. 39. fig. 6.
Payk. Fn. Sv. 3. 116. 18.
Habitat in Europae floribus.

1. L. nigra, elytris faſciis quatuor flauis: anteriore annulari. *annularis.*
Habitat in Sibiria. Muſ. D. Lund.
Statura omnino L. attenuatae. Caput et thorax nigra, pube-aurea parum nitidula. Elytra laeuia, attenuata, nigra, annulo baſeos faſciisque tribus flauis. Corpus nigrum, pube aurulenta, nitidulum.

2. L. nigra, elytris punctis albis: exterioribus duobus minoribus. *12guttata.*
Habitat in Sibiria. Muſ. D. Lund.
Statura praecedentis. Caput et thorax nigra, immaculata. Elytra laeuia, truncata, atra, punctis ſex albis: quatuor ad ſuturam maioribus, duobus exterioribus minoribus.

3. L. nigra, coleoptris attenuatis, flauis: faſciis quatuor nigris; anteriore punctata, ſecunda interrupta, tibiis poſticis bidentatis. Ent. ſyſt. 2. 347. 41. *calcarata.*
~~Geoff. Inſ. 1. 226. 1.~~
Payk. Fn. Sv. 3. 114. 16.
Habitat in Germania. D. Smidt.

4. L. nigra, coleoptris teſtaceis: faſciis quatuor nigris, anteriore punctata, antennis pedibusque flauis. Ent. ſyſt. 2. 347. 42. * *ſubſpinoſa.*
*Leptura armata.* Herbſt. Arch. tab. 26. fig. 24.
Payk. Fn. Sv. 3. 115. 17.
Habitat Kiliae.

5. L. nigra, elytris maculis duabus punctisque duobus flauis. Ent. ſyſt. 2. 347. 43. * *ſinuata.*
Habitat Kiliae.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*4faſciata.* 56. L. nigra, elytris teſtaceis: faſciis quatuor de tatis, nigris. Ent. ſyſt. 2. 348. 44.
*Leptura 4faſciata.* Linn. ſyſt. Nat. 2. 639. 1 Fn. Sv. 687.
Oliv. Inſ. 73. tab. 2. fig. 17.
Schaeff. Icon. tab. 59. fig. 6.
Degeer Inſ. 5. 132. 8. tab. 4. fig. 11.
Payk. Fn. Sv. 3. 114. 15.
Habitat in Europae floribus.

*aurulenta.* 57. L. atra, thoracis margine antico poſticoque a reis, elytris teſtaceis: faſciis quatuor ſimpli bus nigris. Ent. ſyſt. 2. 348. 45. *
VIII. 93. Habitat Kiliae.

*zebrata.* 58. L. nigra, thorace aureo, tomentoſo, elytris faſ giatis, ſubſpinoſis: faſciis quatuor flauis, b ſeos lunata.
Habitat in Carolina. Muſ. Dom. Boſc.
Praecedentibus minor. Antennae nigrae. Cap et thorax tomento aureo tecta. Elytra attenu ta, apice ſpinoſa, nigra, faſciis quatuor flau baſeos lunata reliquis linearibus. Corpus n grum, abdominis ſegmentis margine albo-c liatis. Pedes rufi.

*dubia.* 59. L. ſubuilloſa, nigra, elytris teſtaceis, nigro-pu ctatis, pedibus nigris. Ent. ſyſt. 2. 348. 46.
Habitat in Siberia. Muſ. D. Banks.

*connexa.* 60. L. fuſca, elytris faſciis duabus cinereis, ſutu connexis. Ent. ſyſt. ſuppl. 153. 47. *
Habitat in Tanger. Dom. Schousboe.

*6guttata.* 61. L. nigra, elytris maculis tribus flauis. Ent. ſy 2. 349. 47. *
Herbſt. Arch. tab. 26. fig. 23.
VI. 55 Panz. Fn. Germ. 69. tab. 22.
Payk. Fn. Sv. 3. 119. 21.
Habitat in Germania. Dom. de Hattorff. 6

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

2. L. atra, elytris faſciis tribus flauis: anteriore interrupta. Ent. ſyſt. 2. 349. 48. * — *3faſciata.*
Schaeff. Icon. tab. 39. fig. 10.
Habitat in Suecia. Muſ. D. Lund.
Varietatem L. 6maculatae credit D. de Paykull.

3. L. nigra, elytris baſi teſtaceis, apice nigris, flauo-reticulatis. Ent. ſyſt. ~~ſuppl.~~ 453. 48. * — *reticulata.*
Habitat in Italia. Dr. Allioni.

4. L. viridi-coerulea, elytris ſubfaſtigiatis. Ent. ſyſt. 2. 349. 49. * — *ſericea.*
*Leptura ſericea.* Linn. ſyſt. Nat. 2. 638. 8. Fn. Sv. 683.
Schaeff. Icon. tab. 84. fig. 1.
Habitat in Europae floribus.

5. L. thorace orbiculato, corpore toto atro, nitido. Ent. ſyſt. 2. 349. 50. * — *morio.*
Habitat in Suecia. Muſ. D. de Seheſtedt.

6. L. thorace globoſo abdomineque rubris, elytris nigris. Ent. ſyſt. 2. 349. 51. — *collaris.*
*Leptura collaris.* Linn. ſyſt. Nat. 2. 639. 16. Fn. Sv. 691.
Geoff. Inſ. 1. 228. 11.
Schaeff. Icon. tab. 58. fig. 9.
Degeer Inſ. 5. 143. 22.
Payk. Fn. Sv. 3. 126. 28.
Habitat in Europae floribus.

7. L. thorace globoſo, nigro, elytris violaceis, abdomine ruſo. Ent. ſyſt. 2. 349. 52. * — *virginea.*
*Leptura virginea.* Linn. ſyſt. Nat. 2. 639. 15. Fn. Sv. 690.
Oliv. Inſ. 73. tab. 2. fig. 24.
Schaeff. Icon. tab. 58. fig. 8.
Degeer Inſ. 5. 144. 23.
Payk. Fn. Sv. 3. 125. 27.
Habitat in Europae floribus.

68.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*holoſericea* 68. L. nigra, obſcura, ano rufo.
Habitat in Germania. D. Smidt.
Magnitudo L. virgineae, tota nigra minime nitid ano ſolo late rufo.

*tenuicollis.* 69. L. teſtacea, antennis elytrorumque apicibus n gris.
Habitat in Guinea. Dom. Meier. Muſ. Dom. d Seheſtedt.
Statura ab hoc genere parum differt. Antenna nigrae: articulo primo teſtaceo. Caput exſer tum, ouatum, teſtaceum. Thorax rotundatus laeuis, teſtaceus, antice attenuatus. Elytra lae uia, teſtacea, apice late nigra. Corpus teſta ceum, pectore plantisque fuſcis.

*cylindri-collis.* 70. L. teſtacea, elytris nigris, apice teſtaceis.
Habitat in Guinea. Dom. Meier. Muſ. D. Lund.
Statura omnino praecedentis, at elytra nigra, api ce teſtacea. Antennae nigrae: articulo prim teſtaceo.

*attelaboi-des.* 71. L. teſtacea, elytris nigris immaculatis.
Habitat in Guinea. Dom. Meier. Muſ. D. Lund.
Statura et ſumma affinitas praecedentis et forte mera eius varietas, at elytra tota nigra. An tennae nigrae: articulo primo teſtaceo.
Vltimae tres ſpecies forte proprii generis mihi haud rite examinatur.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

45. MEGALOPVS. *Palpi* filiformes: articulo vltimo acuto.
*Maxilla* compreſſa, cornea, intus ferrata, apice fiſſa.
*Antennae* extrorſum craſſiores, ſubſerratae.

. M. teſtaceus, capitis vertice thoracisque dorſo nigris. *ruficornis.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Genus omnino diſtinctum. *Os* maxillis palpisque. *Palpi* quatuor aequales, filiformes: *anteriores* vix longiores, quadriarticulati; articulis inaequalibus: ſecundo longiori, vltimo acuto, adhaerentes maxillae dorſo; *poſteriores* triarticulati; articulis ſubaequalibus: vltimo obtuſiori adnati ligulae medio. *Mandibula* cornea, arcuata, edentula, acuta. *Maxilla* compreſſa, cornea, intus ferrata, apice fiſſa. *Ligula* porrecta, longitudine maxillae, cornea, bifida: laciniis approximatis, cylindricis, obtuſis. *Labium* transuerſum, corneum. *Antennae* extrorſum craſſiores: articulo primo longiori, ſecundo orbiculato, reliquis ſubſerratis.

Cor-

---

*Megalopi* corpus medium, oblongum, glabrum, immarginatum, tardum; *capite* ouato, exſerto; *oculis* magnis, globoſis, prominentibus, lateralibus; *antennis* diſtantibus, thorace breuioribus, in cantho oculorum antico inſertis; *thorace* breui, cylindrico, elytris anguſtiore; *ſcutello* breui triangulo; *elytris* longitudine abdominis, rigidis, incumbentibus; *pedibus* longiuſculis, curſoriis; *femoribus* poſticis, incraſſatis, *tibiisque* incuruis; *tarſis* omnibus quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Corpus medium, ſtatura et magnitudine fere X rufae. Caput cum antennis teſtaceum, vertice fuſco. Oculi magni, globoſi, prominuli. Thorax rotundatus, teſtaceus, dorſo fuſco. Elytra laeuia, teſtacea, ſubattenuata, immaculata. Pectus fuſcum, abdomine teſtaceo. Pedes teſtacei femoribus poſticis incraſſatis.

*nigricornis* 2. M. teſtaceus, antennis elytrorum margine tibiisque poſticis nigris.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et ſumma affinitas praecedentis. Differt tamen capite cum antennis, nigro ore, flaueſcente. Thorax teſtaceus, dorſo nigro. Elytra punctata, teſtacea, margine nigro et in hoc margine macula magna, baſeos ouata, teſtacea. Corpus teſtaceum, femoribus incraſſatis, tibiis nigris.

## 146. NECYDALIS. *Palpi* filiformes. *Maxilla* vnidentata. *Ligula* membranacea, late emarginata. *Antennae* filiformes.

*thalaſſina.* 1. N. thorace canaliculato, corpore viridi, pedibus nigris. Ent. ſyſt. 2. 350. 1. *

Panz.

---

*Necydalis* corpus oblongum, anguſtum, tomentoſum, immaculatum, agile; *capite* ouato, inſerto; *oculis* globoſis, prominulis, marginalibus; *antennis* thorace longioribus, approximatis, inter oculos inſertis; *thorace* rotundato; *dorſo* inaequali; *ſcutello* breui, rotundato; *elytris* rigidis, attenuatis, incumbentibus longitudine abdominis; *pedibus* longioribus, tenuibus; *femoribus* poſticis ſaepius clauatis; *tarſis* anticis quinque, poſticis quadriarticulatis; *colore* vario.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 5. tab. 15. I. 46
Habitat Kiliae.

. N. thorace teretiuſculo, corpore viridi, pedibus muticis, teſtaceis. Ent. ſyſt. 2. 350. 2. *viridiſſima*
*Cantharis viridiſſima.* Linn. ſyſt. Nat. 2. 650. 23. Fn. Sv. 717.
*Oedemera viridiſſima.* Oliv. Inſ. 50. t. 2. f. 15.
Geoff. Inſ. 1. 177. 14.
~~Degeer Inſ. 5. 15. 4. tab. 1. fig. 13.~~
Payk. Fn. Sv. 3. 134. 1.
Habitat in Europae hortis frequens.

. N. thorace teretiuſculo, corpore coeruleo, ſubopaco. Ent. ſyſt. 2. 3. * *coeruleſcens*
~~*Cantharis coerulea.* Linn. ſyſt. Nat. 2. 650. 22. Fn. Sv. 716.~~
~~*Oedemera coerulea.* Oliv. Inſ. 50.~~ tab. 2. ~~fig. 14.~~
~~*Telephorus villoſus.* Degeer Inſ. 4. 76. 8.~~
Payk. Fn. Sv. 3. 135. *varied. nigricolli*
Habitat in Europae hortis.

. N. thorace inaequali, corpore vireſcente, obſcuro, antennis pedibusque nigris. Ent. ſyſt. 2. 351. 4. * *vireſcens.*
*Cantharis vireſcens.* Linn. ſyſt. Nat. 2. 650. 24.
Payk. Fn. Sv. 3. 136. 6.
Habitat in Germania. Dom. Smidt.

. N. thorace ouato, cyanea, antennis pedibusque nigris. Ent. ſyſt. 2. 351. 5. * *cyanea.*
Habitat in Germania. Dom. Smidt.

. N. thorace ouato: maculis duabus baſeos nigris, corpore fuſco. *notoxoides*
Habitat in Carolina. Muſ. Dom. Boſc.
Statura et magnitudo praecedentium. Caput cum antennis nigrum. Thorax ouatus, rufus,



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

maculis duabus baſeos nigris. Elytra laeuia fuſca.

*ſanguinicollis.* 7. N. fuſca, thorace teretiuſculo, rufo: punctis duobus ~~com~~preſſis, elytris lineis tribus eleuatis. Ent. ſyſt. 2. 351. 6. *

II. 43. *Oedemera ſanguinicollis.* Oliv. Inſ. 70. tab. 1. fig. 12.

Habitat Halae Saxonum. Dom. Hybner.

*thoracica.* 8. N. thorace teretiuſculo, rufo, elytris laeuiſſimis fuſcis.

Habitat in Carolina. Muſ. Dom. Boſc.

Statura et ſumma affinitas praecedentis. Corpus fuſcum, at thorax rufus et elytra pubeſcentia laeuiſſima.

*barbara.* 9. thorace inaequali, vireſcens, elytris apice pedibusque flaueſcentibus. Ent. ſyſt. 2. 351. 7. *

Habitat in Barbaria. Dom. Desfontaines.

*ſeladonia.* 10. N. thorace inaequali, corpore viridi, nitidulo, elytris apice incraſſatis, coeruleis. Ent. ſyſt. 2. 351. 8. *

Habitat Kiliae.

*ruficollis.* 11. N. thorace teretiuſculo abdomineque rufis, capite elytrisque viridi-aeneis. Ent. ſyſt. 2. 352. 9. *

*Oedemera ruficollis.* Oliv. Inſ. 50. tab. 1. fig. 11.

Habitat in Italia. Dr. Allioni.

*melanocephala.* 12. N. nigra, thorace abdomineque fuluis, elytris teſtaceis. Ent. ſyſt. ~~Suppl.~~ Append. IV. 153. 9. *

III. 57. ~~Panz. Fn. Germ. 36. tab. 9.~~

Habitat in Italia. Dr. Allioni.

*triſtis.* 13. N. nigra, thorace teretiuſculo femorumque baſi rufis. Ent. ſyſt. 2. 352. 10. *

*Oedemera triſtis.* Oliv. Inſ. 50. tab. 2. fig. [illegible]

H[illegible]

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Terra Diemenii. Muſ. D. Banks.

4. N. elytris ſubulatis, atra, femoribus omnibus clauatis. Ent. ſyſt. 2. 352. 11. * *atra.*
Linn. ſyſt. Nat. 2. 646. 5.
Habitat in Europa auſtraliori.

5. N. elytris ſubulatis, nigris, baſi flauis. Ent. ſyſt. 2. 352. 12. * *humeralis.*
*Cantharis humeralis.* Oliv. Inſ. 50. t. 2. f. 20.
Geoff. Inſ. 1. 342. 2.
Forſt. nou. Inſ. ſp. Cent. 1. 48. 48.
Habitat in Anglia.

6. N. nigra, elytris baſi ſuturaque late teſtaceis. Ent. ſyſt. 2. 352. 13. * *vſtulata.*
*Oedemera vſtulata.* Oliv. Inſ. 50. tab. 2. fig. 19.
Habitat in Germaniae plantis.

7. N. nigra, thorace elytrisque teſtaceis: his apice nigris. Ent. ſyſt. 2. 353. 14. * *melanura.*
*Oedemera melanura.* Oliv. Inſ. 50. tab. 1. fig. 8. *M. 57.*
Habitat in Hiſpaniae plantis. Dom. Prof. Vahl.

8. N. capite thoraceque ferrugineis, elytris teſtaceis apice nigris Ent. ſyſt. 2. 353. 15. * *notata.*
~~Payk. Fn. Sv. 3. 132. 1.~~
Habitat in Germania. Dom. Hybner.

9. N. teſtacea, antennis nigris, elytrorum margine coeruleo, nitido. *marginella*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo omnino N. notatae. Caput teſtaceum, clypeo nigro. Antennae nigrae. Thorax rotundatus, teſtaceus, immaculatus. Elytra lineis duabus eleuatis, teſtacea, margine baſeos et laterali coeruleis, nitidis. Corpus teſtaceum.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fuluicollis.* 20. N. atra, thorace anoque fuluis, elytris subcoe-
ruleſcentibus. Ent. ſyſt. 2. 353. 16. *
Habitat in Germania. Muſ. D. Lund.

*longipes.* 21. N. niger, elytris aurulentis, dorſo cinereis, pe-
dibus quatuor, poſticis elongatis, compreſſ-
eleuatis. Ent. ſyſt. ſuppl. 155. 16. ♀
Habitat in Inſula St. Domingo. Muſ. Dom. Boſc

*rufa.* 22. N. elytris ſubulatis, nigra, femoribus omnibu
clauatis, elytris antennisque ruſis. Ent. ſyſt. 2
353. 17.
Linn. ſyſt. Nat. 2. 642. 6.
Geoff. Inſ. 1. 220. 22.
Schaeff. Icon. tab. 94. fig. 8.
Habitat in Europa auſtraliori.

*praeuſta.* 23. N. atra, elytris teſtaceis, apice nigris, femorib
omnibus clauatis. Ent. ſyſt. 2. 354. 18. ♀
Habitat in Barbaria. Muſ. Dom. Desfontaines.

*margari-*
*tacea.* 24. N. rufa, elytris ſubulatis, abbreuiatis, margar
taceis, baſi rufis, apice atris.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Statura omnino N. coeruleae. Caput ruſum. A
tennarum articulis apice atris. Thorax glob
ſus, ruſus, immaculatus. Elytra laeuiſſim
ſubulata, abdomine duplo fere breuiora, ba
ruſa, in medio margaritacea, pitidiſſima, api
atra. Abdomen nigrum, ano rufo. Pedes ru
femoribus clauatis.

*coerulea.* 25. N. elytris ſubulatis, coerulea, femoribus poſti
cis clauatis arcuatis. Ent. ſyſt. 2. 354. 19.
*Necydalis coerulea.* Linn. ſyſt. Nat. 2. 642. 4.
*Necydalis ceramboides.* Forſt. nou. Inſ. ſpe
Cent. 1. 47. 47.
*Oedemera coerulea.* Oliv. Inſ. 50. tab. 2. f. 1
Geo

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 342. 3.
Sulz. Hift. Inf. 48. tab. 6. fig. 2.
Schaeff. Icon. tab. 94. fig. 7.
Habitat in Oriente et Europa auftraliori.
*Variat* pedibus fimplicibus.

16. N. elytris fubulatis teftaceis, corpore nigro, femoribus poſticis clauatis baſi teſtaceis. Ent. ſyſt. 2. 354. 20. * *Podagrariae.*
*Oedemera Podagrariae.* Oliv. Inf. 50. tab. 1. fig. 10.
Linn. ſyſt. Nat. 2. 642. 9.
Geoff. Inf. 1. 343. 4.
Payk. Fn. Sv 3. 134. 4.
Habitat in Europae hortis.

17. N. coeruleſcens, thoracis margine abdomine pedibusque teſtaceis. Ent. ſyſt. 2. 354. 21. * *bicolor.*
Habitat ad Cap. Bon. Spei. Dom. de Paykull.

18. N. nigra, elytris attenuatis, vireſcentibus, femoribus poſticis incraſſatis, arcuatis. Ent. ſyſt. 2. 355. 22. * *clauipes.*
*Oedemera aenea.* Oliv. Inf. 50. tab. 2. fig. 18.
Payk. Fn. Sv. 3. 137. 7.
Habitat Kiliae.

29. N. nigra, thorace elytrisque teſtaceis, pedibus rufis. Ent. ſyſt. 2. 355. 23. * *teſtacea.*
Habitat in Germania.

30. N. nigra, elytris teſtaceis: margine omni nigro. Ent. ſyſt. ſuppl. 155. 23. * *marginata*
*Necydalis femorata.* Panz. Fn. Germ. 36. tab. 12.
Habitat in Gallia. Dom. Boſc.
De Synonymo D. Panzer haereo, cum femora in meo ſpecimine ſimplicia. Forte varietas ſexus.

31. N. elytris ſubulatis, glaucis, corpore nigro, abdominis inciſuris albis. Ent. ſyſt. 2. 355. 24. * *glauceſcens*
Linn.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. syst. Nat. 2. 642. 7.
Schaeff. Icon. tab. 94. fig. 7.
Habitat in Europa australi.

*simplex.* 32. N. elytris subulatis, testaceis, pedibus simplicibus. Ent. syst. 2. 355. 25. *
*Necydalis simplex.* Linn. syst. Nat. 2. 643. 10.
*Necydalis flauescens.* Payk. 3. 135. 5.
~~*Oedemera simplex.* Oliv.~~ Inf. 50. tab. 1. fig. 9.
Degeer Inf. 5. 155. 4.
Habitat in Europae boreali.

*Quercus.* 33. N. elytris subulatis, abdomine pedibusque flauescentibus. Ent. syst. 2. 355. 26. *
Habitat in Daniae Quercu. Mus. Dom. Lund.

147. MOLORCHVS. *Palpi* inaequales, filiformes.
*Maxilla* membranacea, bifida.
*Ligula* bifida.
*Antennae* setaceae.

*abbreuiatus.* 1. M. elytris dimidiatis, ferrugineis, immaculatis, antennis breuibus. Ent. syst. 2. 356. 1. *
*Necydalis maior.* Linn. syst. Nat. 2. 641. 1.
Schaeff. Monogr. 1753. fig. 12.
Elem. tab. 13. fig. 2.
*IV. 65* Icon. tab. 10. fig. 10. 11. Degeer

---

*Molorchi* corpus oblongum, angustatum, glabrum, immarginatum, agile; *capite* ouato, decliui, inserto; *oculis* reniformibus, vix prominulis, marginalibus; *antennis* thorace longioribus, approximatis, inter oculos insertis; *thorace* rotundato, angusto, dorso canaliculato; *elytris* rigidis, fornicatis, abbreuiatis; *alis* plicatis, abdomini incumbentibus; *abdomine* elongato: margine vtrinque reflexo; *pedibus* longiusculis; *femoribus* clauatis; *tarsis* quadriarticulatis; *colore* testaceo.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 5. 148. 1. tab. 5. fig. 1
Panz. Fn. Germ. 41. tab. 20.
Payk. Fn. Sv. 3. 129. 1.
Habitat in Europae floribus.

2. M. elytris abbreuiatis, abdomine atro: faſciis fuluis. Ent. ſyſt. 2. 357. 2. * *variegatus*
Habitat in noua Hollandia. Muſ. D. Banks.

3. M. elytris dimidiatis, teſtaceis, apice lineola alba, antennis longis. Ent. ſyſt. 2. 357. 3. *dimidiatus*
*Necydalis minor.* Linn. ſyſt. Nat. 2. 641. 2. Fn. Sv. 837.
Schaeff. Monogr. 1753. fig. 6. 7. Icon. tab. 95. fig. 5.
Sulz. Inſ. tab. 7. fig. 51.
Panz. Fn. Germ. 41. tab. 21.
Payk. Fn. Sv. 3. 130. 2.
Degeer. Inſ. 5. 151. 2.
Habitat in Europae hortis.

IV. 65

4. M. elytris dimidiatis, teſtaceis, immaculatis, antennis longis. Ent. ſyſt. 2. 357. 4. *vmbellatarum.*
*Necydalis vmbellatarum.* Linn. ſyſt. Nat. 2. 641. 3.
Schaeff. Icon. tab. 95. fig. 4.
Sulz. Hiſt. Inſ. tab. 6. fig. 1.
Payk. Fn. Sv. 3. 131. 3.
Habitat in Germaniae vmbellatis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

148. SPONDYLIS. *Palpi* filiformes.
*Maxilla* conica, craſſa, integra.
*Labium* bifidum.
*Antennae* moniliformes.

*bupreſtoides.* 1. Spondylis. Ent. ſyſt. 2. 358. 1.
*Attelabus bupreſtoides.* Linn. ſyſt. Nat. 2. 621. 3.
Friſh. Inſ. 3. 23. tab. 19.
Degeer Inſ. 5 34 21. tab. 5. fig. 21.
IV. 66 Paykull Fn. Sv. 3. 155. 1.
Habitat in Europae borealis ſyluis.

149. SINODENDRON. *Palpi* inaequales.
*Maxilla* vnidentata.
*Labium* filiforme, apice palpigerum.
*Antennae* lamellatae.

*cylindricum.* 1. S. thorace antice truncato quinquedentato, capitis cornu erecto. Ent. ſyſt. 2. 358. 94.
*Scarabaeus cylindricus.* Linn. ſyſt. Nat. 2. 544. 11. Fn. Sv. 380.
Oliv. Inſ. 1. 3. 47. 54. tab. 9. fig. 80.
I. 46 47. Degeer Inſ. 4. tab. 10. fig. 2.
Voet. Scarab. tab. 20. fig. 131.

Laichart.

---

*Sinodendri* corpus cylindricum, glabrum, immarginatum, antice poſticeque obtuſum, tardum; *capite* paruo, ouato, plano; *oculis* paruis, rotundatis, marginalibus; *antennis* breuibus, reflexis, ſub oculos inſertis; *thorace* gibbo, antice truncato, muricato, aut inaequali; *ſcutello* paruo, triangulari; *elytris* rigidis, fornicatis, abdomen includentibus, apice ſaepius retuſis, dentatis; *pedibus* breuibus, validis; *tibiis* anticis, dentatis; *tarſis* quinquearticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Laichart. Inſ. 4. 4.
Scopol. Intr. Hiſt. nat. 5. 75.
Panz. Fn. Germ. 1. tab. 1.
Payk. Fn. Sv. 3. 140. 1.
Habitat in Europae borealis truncis Cerafi.

2. S. thorace antice mucronato, mandibulis vnaris, porrectis, medio vnidentatis. *cornutum.*
Habitat in Terra Diemenii. Dom. Billardiere.
Statura et magnitudo omnino praecedentis, obſcure-ferrugineum. Caput minutum, vtrinque ante oculos dente eleuato, minuto, mandibulisque valde elongatis, fere rectis, in medio dente laterali, valido armatis. Antennae lamellis ſex. Thorax punctatus, canaliculatus, antico mucronatus. Elytra crenato-ſtriata. Tibiae anticae ſerratae.
Foemina ſimillima, at mandibulis haud porrectis inermibus.

3. S. atrum, nitidus, thorace cylindrico, elytris crenato-ſtriatis, pedibus ferrugineis. *digitatum*
*Scarites cylindricus.* Ent. ſyſt. ſuppl. 44. 8. *
Habitat in India orientali. D. Daldorff.

4. S. thorace muricato, nigrum, elytris apice retuſis vnidentatis. *vnidentatum.*
Habitat in India orientali. D. Daldorff. Muſ. D. de Seheſtedt.
Statura et magnitudo S. gibbi. Antennae lamellatae, flauae. Thorax gibbus, muricatus. Elytra punctata, apice retuſa, dente ad ſuturam eleuato, acuto.

5. S. thorace muricato, gibbo, elytris retuſis, ſtriatis. Ent. ſyſt. ſuppl. 156. 1. *gibbum.*
Habitat in India orientali. D. Daldorff.

6. S. thorace muricato, gibbo, elytris ante apicem biſpinoſis. Ent. ſyſt. 2. 359. 2. * *muricatum.*

Der-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Dermeſtes muricatus.* Linn. ſyſt. Nat. 2. 562. 6.
Oliv. Inſ. 77. tab. 1. fig. 1.
Roſſ. Fn. Etr. 1. 38. 95.
Habitat in Europae auſtralioris et Africae lignis.

*capucinus.* 7. S. elytris integris nigris, thorace antice ſcabro. Ent. ſyſt. 2. 359. 3. *
Habitat in Coromandel. Dr. Koenig.

*dominicum.* 8. S. laeue, nigrum, obſcurum, elytris ſtriatis, pedibus piceis. Ent. ſyſt. 2. 359. 4. *
Habitat in America meridionali. D. Pflug.

*puſillum.* 9. S. ferrugineum, elytris punctato-ſtriatis integris. Ent. ſyſt. ſuppl. 156. 5. *
Habitat in India orientali. D. Daldorff.
Corpus totum paruum, cylindricum, ferrugineum, immaculatum.

## 150. APATE.

*Palpi* aequales, filiformes.
*Maxilla* vnidentata.
*Labium* truncatum.
*Antennae* perfoliatae.

*muricata* 1. A. elytris reticulatis, poſtice retuſis dentatis, thorace muricato gibbo. Ent. ſyſt. 2. 360. 1.
*Ligniperda terebrans.* Pall. Specil. Zool. 9. tab. 1. fig. 3.
Voet. Col. 2. tab. 44. fig. 1.
Oliv. Inſ. 77. tab. 1. fig. 4.
Habitat in Americae meridionalis Saccharo.

2.

---

*Apates* corpus cylindricum, ſubtomentoſum, immarginatum, tardum; *capite* paruo rotundato, inflexo; *oculis* paruis, globoſis, marginalibus; *antennis* breuibus, diſtantibus, ſub oculos inſertis; *thorace* rotundato, gibbo, antice ſaepius muricato; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis; *tarſis* quadriarticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. A. elytris ante apicem bidentatis, thorace gibbo, antice muricato: dente laterali hamato denticulato. *franciſca.*

Habitat in Americae Inſulis. Dr. Pflug.

Statura praecedentis paullo minor. Antennae nigrae: claua compreſſa, ferruginea. Caput nigrum, pube ferruginea, clypeoque pube aurea ciliato. Thorax gibbus, ater, antice muricatus, vtrinque dente porrecto, hamato, dentato. Elytra punctata: lineis duabus eleuatis, ante apicem excurrentibus. Abdomen obſcurum.

3. A. elytris ante apicem tridentatis, thorace gibbo antice muricato, capite bidentato. *monacha.*

Habitat in Americae Inſulis.

Statura et fere magnitudo A.muricatae. Antennae nigrae: claua magna, compreſſa, fuſca. Caput nigrum, fronte dentibus duobus eleuatis et ſub his pili denſi, ferruginei. Thorax gibbus antice dentibus plurimis, eleuatis, ſcaber. Elytra ſubreticulata: lineis tribus eleuatis ante apicem in dentem excurrentibus. Corpus ferrugineo-villoſum.

4. A. elytris ante apicem ſubdentatis, thorace gibbo, muricato, capite rufo, hirto. *Carmelita.*

Habitat in Americae Inſulis. Muſ. D.de Seheſtedt.

Statura praecedentis, at paullo minor et alia. Antennae nigrae: claua compreſſa, ferruginea. Caput totum pilis denſis, ferrugineis tectum. Thorax gibbus, antice muricatus, at denticuli multo minores, vix eleuati, nullo modo recurui. Elytra punctata, lineis tribus eleuatis, ante apicem in dentes excurrentibus. Abdomen obſcure ferrugineum.

Specimina e Guinea et Tanger miſſa nullo modo differunt.

5.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Jeſuita.* 5. A. elytris integris varioloſis, thorace gibbo, muricato: dente vtrinque hamato. Ent. ſyſt. 2. 361. 5. *

Habitat in noua Hollandia. Muſ. D. Banks.

*rugoſa.* 6. A. thorace gibbo, muricato, elytris integris rugoſis.

Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

Statura et magnitudo A. 2maculatae. Corpus totum atrum, immaculatum. Thorax valde gibbus, muricatus. Elytra reticulata, apice rotundata, integra.

*dominicana.* 7. A. elytris retuſis dentatis, thorace muricato: dente laterali hamato, pedibus ruſis.

Habitat in India orientali. Muſ. D. de Seheſtedt

Duplo minor quam A. Ieſuita. Caput nigrum, fronte pube ferruginea. Antennae ferrugineae. Thorax niger, antice muricatus: dente laterali hamato. Elytra ſubpunctata, poſtice retuſa, dentata. Corpus ſubtus cinereo-villoſum. Pedes ferruginei.

*femoralis.* 8. A. elytris laeuibus, retuſis, cylindrica, atra, femoribus anticis incraſſatis. Ent. ſyſt. 2. 361. 6. *

Habitat in Americae Inſulis. Muſ. D. Lund.

*gonagra.* 9. A. elytris retuſis, punctatis, cylindricus, niger, femoribus anticis compreſſis, incraſſatis. Ent. ſyſt. ſuppl. 157. 7. *

Habitat in Americae meridionalis Inſulis. Dom. v. Rohr.

*cornuta.* 10. A. thorace antice vtrinque vltra caput porrecto, ſerrato, hirto. Ent. ſyſt. 2. 360. 2. *

*Boſtrichus cornutus.* Oliv. Inſ. 77. tab. 1. fig. 5.

Habitat in Madagaſcar. Dom. Olivier.

*hamata.* 11. A. elytris ante apicem vniſpinoſis, thoracis margine antico bihamato, dentato. Ent. ſyſt. 2. 360. 3. *

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Saxoniae ligno. Dom. Prof. Leske.

12. A. nigra, elytris abdomineque rufis, thorace emarginato, retuſo. Ent. ſyſt. 2. 362. 7. *capucina.*

*Dermeſtes capucinus.* Linn. ſyſt. Nat. 2. 562. 5. Fn. Sv. 416.

*Boſtrichus.* Geoff. Inſ. 1. 302. 1. tab. 5. fig. 1. Schaeff. Elem. tab. 28. Icon. tab. 189. fig. 1. Sulz. Hiſt. Inſ. tab. 2. fig. 5. Roſſ. Fn. Etr. 36. 87. Oliv. Inſ. 77. tab. 1. fig. 1. Panz. Fn. Germ. 34. tab. 18. Payk. Fn. Sv. 3. 141.

Habitat in Europae truncis emortuis.

Varietas abdomine nigro e Tanger in Muſeo D. Lund vix differt.

13. A. thorace gibbo ſcabro: vtrinque macula poſtica alba, nigro-punctata, elytris poſtice vniſpinoſis. Ent. ſyſt. 2. 362. 8. * *2maculata*

*Boſtrichus 2maculatus.* Oliv. Inſ. 77. tab. 2. fig. 14.

Habitat in Gallia prouincia. Muſ. Dom. Olivier.

14. A. thorace gibbo muricato, elytris apice retuſis, ſinuatis. Ent. ſyſt. 2. 362. 9. * *ſinuata.*

Habitat in Gallia. Muſ. Dom. Boſc.

15. A. elytris retuſis, dentatis, picea, thorace antice gibbo dentato. Ent. ſyſt. 2. 362. 10. * *tridens.*

Habitat in Inſula St. Thomas Americae. Muſ. D. de Seheſtedt.

16. A. elytris ſtriatis, retuſis, bidentatis: dente interiore elongato, thorace antice muricato. Ent. ſyſt. ſuppl. 157. 10. * *bidens.*

Habitat in Tanger. D. Schousboe. Muſ. D. Lund.

17. A. elytris retuſis, thorace muricato: punctis duobus dorſalibus rufis, elytris baſi rufis. *puſtulata.*

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717 Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Tanger. D. Schousboe. Muſ. D. de Seheſtedt.

Corpus paruum, nigrum. Caput pube denſa, cinerea. Thorax niger, antice muricatus, poſtice punctis duobus dorſalibus, rubris. Elytra ſubpunctata, nigra, baſi ferruginea, apice retuſa. Corpus nigrum, femoribus rufis.

*torquata.* 18. A. thoracis margine antico prominulo villoſo elytris poſtice retuſis tridentatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Magnitudo A. tridentis. Caput planum, antennis flauis: articulis tribus vltimis elongatis. Thorax rotundatus, ater, nitidus: margine prominulo, hirto. Elytra ſublaeuia, nigra, apice retuſa, tridentata.

*marginata.* 19. A. thorace gibbo, muricato, elytris brunneis apice retuſis ſubreflexis.

Habitat in India orientali. D. Daldorff. Muſ. D. de Seheſtedt.

Statura praecedentium, at diſtinctus, elytris brunneis, retuſis, margine ſuturaque eleuatis, apice parum reflexis at absque dente.

*limbata.* 20. A. atra, elytris integris pallidis: margine omni nigro. Ent. ſyſt. 2. 363. 13. *

*Dermeſtes domeſticus.* Linn. ſyſt. Nat. 2. 643. 13. Fn. Sv. 424.

*Boſtrichus limbatus.* Herbſt. Arch. tab. 20. f. 11.

Panz. Fn. Germ. 43. tab. 19.

Payk. Fn. Sv. 3. 144. 5.

Habitat Kiliae ſub cortice Fagi. D. Daldorff.

*diſpar.* 21. A. atra, thorace gibbo, antice ſcabro, tibiis teſtaceis. Ent. ſyſt. 2. 363. 11.

Payk. Fn. Sv. 3. 143. 4.

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Germaniae truncis. D. Daldorff.

12. A. nigra, thorace teſtaceo: punctis duobus lineolaque media nigris, elytris pallidis: linea poſtica nigra. Ent. ſyſt. 2. 363. 12. * *ſignata.*
Habitat in Germania. D. Smidt.

13. A. atra, laeuis, nitidus, elytris laeuiſſimis, pedibus piceis. Ent. ſyſt. 2. 363. 14. * *voluulus.*
Habitat in Gallia.

24. A. thorace gibbo, muricato, elytris laeuibus integris brunneis. *brunnipennis.*
Habitat in India. D. Daldorff. Muſ. D. de Seheſtedt.
Parua, atra, elytris integris, brunneis. Pedes brunnei.

25. A. nigra, elytris integris piceis, thorace antice ſcabro. Ent. ſyſt. 2. 363. 15. * *minuta.*
Habitat in noua Zelandia. Muſ. D. Banks.

26. A. thorace rugoſo, nigra, elytris piceis. Ent. ſyſt. ſuppl. 157. 16. * *Fagi.*
Habitat in Saxoniae Fago.
Antennae mihi perfoliatae videntur.

27. A. thorace ſcabro, elytris pedibusque griſeis. Ent. ſyſt. ſuppl. 157. 17. * *Tiliae.*
*Apate Tiliae* Panz. Fn. Germ. 8. tab. 14.
Habitat in Germaniae Tilia. Dr. Panzer.

28. A. thorace rugoſo, nigra, antennis pedibusque ferrugineis. *brunnipes.*
Habitat in Americae Inſulis. Dr. Pflug. Muſ. D. Lund.
Paruus. Antennae ferrugineae. Caput laeue, atrum. Thorax rugoſus, niger. Elytra laeuia, integra, nigra. Pedes ferruginei.

29. A. nigra, thorace elytrisque laeuiſſimis, nigris. *glabrata.*
Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Minuta, at diſtincta, thorace omnino laeui elytrisque integris.

## 151. BOSTRICHVS.

*Maxilla* cornea.

*Labium* cylindricum integrum, apice palpigerum.

*Antennae* claua ſolida compreſſa.

*flauicornis* 1. B. rufus, elytris nigris, apice ſpinoſo dentatis antennis flauis. Ent. ſyſt. 2. 364. 1. *

Habitat in America boreali.

*cylindrus.* 2. B. cylindricus, ater, elytris ſtriatis apice villoſis dentatis, pedibus compreſſis teſtaceis. Ent. ſyſt. 2. 364. 2. *

*Il. 43.* *Platypus cylindrus.* Herbſt. Col. 5. tab. 49. f. 3. Panz. Fn. Germ. 15. tab. 1.

Habitat in Germaniae Quercu. D. Helwig.

*parallelus.* 3. B. cylindricus, ferrugineus, elytris laeuibus pallidis: margine ferrugineo.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Corpus cylindricum magnitudine B cylindri. Antennae breues, flauae: claua magna, compreſſa.

Caput

---

*Boſtrichi* corpus paruum, cylindricum, tomentoſum, immarginatum, vtrinque obtuſum, tardum; *capite* paruo, rotundato recondito; *oculis* rotundatis, marginalibus; *antennis* breuibus, diſtantibus. ſub oculos inſertis; *thorace* rotundato, cylindrico; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis longitudine abdominis; *pedibus* breuibus, validis; *tarſis* quadriarticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Caput et thorax laeuia, ferruginea, immaculata. Elytra laeuia, pallida: margine omni ferrugineo. Corpus ferrugineum.

4. B. cylindricus, ater, nitidus, thorace elytris longiore. *thoracicus*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Corpus totum cylindricum, laeue, atrum, nitidum. Labrum ſtriatum, porrectum, acutum. Antennae claua ſolida, compreſſa. Thorax elongatus, elytris longior. Elytra laeuiſſima, abdomine breuiora. Abdomen acutum prominet. Pedes breues, tibiis quatuor anticis ſerratis.

5. B. cylindricus, ater, nitidus, thorace elytris longiore, pedibus ferrugineis. *proboſcideus.*

Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.

Statura omnino praecedentis, at duplo fere minor. Antennae ferrugineae, claua magna, compreſſa. Thorax elongatus, punctatus. Elytra laeuiſſima, atra, abdomine breuiora. Pedes ferruginei.

6. B. cylindricus, ater, nitidus, thorace elytris longiore, elytris puncto rufo. *2puſtulatus*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Affinis praecedentibus, at minor et elytra macula magna, rufa.

Species praecedentes omnes ab hoc genere, habitu antennarumque claua magna, compreſſa differunt. Forte proprii generis mihi haud rite examinatae.

7. B. piloſus, teſtaceus, elytris ſtriatis, retuſis praemorſo dentatis. Ent. ſyſt. 2. 365. 3. *typographus.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Dermeſtes typographus.* Linn. ſyſt. Nat. 2. 56
7. Fn. Sv. 418.
Payk. Fn. Sv. 3. 145. 1.
Degeer Inſ. 4. tab. 6. fig. 1. 2.
II. 43. Panz. Fn. Germ. 15. tab. 2.
Sulz. Inſ. tab. 2. fig. 4.
Act. Holm. 14. 163.
Schaeff. Icon. tab. 259. fig. 3.
Habitat intra arborum cortices, labyrinthos daed
leos puluere repletos formans, pinetorum peſti
Variat colore et magnitudine.

*varians.* 8. B. glaber, teſtaceus, elytris retuſis, ſerratis, vn
dentatis.
Habitat in America meridionali. D. Smidt. Mu
D. de Seheſtedt.
Statura et magnitudo praecedentis. Thorax gl
ber, antice parum rugoſus, teſtaceus. Elyt
breuia, valde retuſa, ſerrata, dente vnico, cl
uato, longiori. Corpus teſtaceum.

*vnidenta-tus.* 9. B. ater, thorace antice ſcabro, elytris retuſis, ſe
ratis, vnidentatis.
Habitat in America meridionali. D. Smidt. Mu
D. Lund.
Corpus totum glabrum, atrum, nitidum, ſtatu
fere Apate, at huius generis. Thorax antic
ſcaber. Elytra laeuia, atra, nitida, apice retu
ſa, ſerrata, denteque vnico elongato. Pede
ferruginei.

*Laricis.* 10. C. niger, elytris retuſis, praemorſo dentatis, p
dibus piceis. Ent. ſyſt. 2. 365. 4.
Payk. Fn. Sv. 3. 147. 3.
II. 44. Panz. Fn. Germ. 15. tab. 3.
Herbſt. Coleopt. 5. tab. 48. fig. 13.
Habitat ſub cortice Pini Laricis Germaniae. Dom
Helwig.

11

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

11. B. ater, elytris praemorſo dentatis rufis. Ent. ſyſt. 2. 365. 5. *chalcographus.*
*Dermeſtes chalcographus.* Linn. ſyſt. Nat. 2. 562. 8. Fn. Sv. 417.
Panz. Fn. Germ. 39. tab. 20. IV. 67
Payk. Fn. Sv. 3. 147. 4.
Habitat intra arborum cortices.

12. B. nigricans, elytris glaucis, obtuſiuſculis. Ent. ſyſt. 2. 365. 6. * *polygraphus.*
*Dermeſtes polygraphus.* Linn. ſyſt. Nat. 2. 562. 10. Fn. Sv. 420.
Panz. Fn. Germ. 15. tab. 5. II. 44.
Payk. Fn. Sv. 3. 155. 15.
Herbſt. Arch. 4. tab. 20. fig. 9.
Coleopt. 5. tab. 48. flg. 6.
Habitat ſub arborum corticibus.

13. B. nigricans, thorace rufo, elytris apice retuſo-dentatis. Ent. ſyſt. 2. 365. 7. * *monographus.*
Payk. Fn. Sv. 3. 149. 6.
Habitat in Germaniae arboribus. Dom. Koch.

14. B. villoſus, teſtaceus elytris integris, tibiis anticis dentatis. *Dactyliperda.*
*Boſtrichus Dactyliperda.* Panz. Fn. Germ.
Habitat in Dactylorum nucleis. Dom. de Meyerle.
Paruus, affinis certe B. micrographo Differt tamen corpore magis villoſo, teſtaceo tibiisque anticis teſtaceis.

15. B. ferrugineus, elytris integris teſtaceis. Ent. ſyſt. 2. 366. 8. * *micrographus.*
*Dermeſtes micrographus.* Linn. ſyſt. Nat. 2. 562. 11.
Payk. Fn. Sv. 3. 155. 13.
Panz. Fn. Germ. 66. tab. 13. VI. 36
Habitat in Europae ligno.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*flauipes.* 16. B. ater, nitidus, elytris apice retusis, serratis antennis pedibusque flauis.

Habitat in America meridionali. Mus. D. de Seheſtedt.

Vl. 96

Paruus. Caput atrum, antennis flauis. Thorax antice subrugosus. Elytra laeuia, atra, nitida apice valde retusa, serrata. Pedes flaui.

*serratus.* 17. B. niger, thorace depresso, elytris testaceis, apice retusis, serratis.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura parua praecedentis. Caput et thorax punctata, nigra. Elytra laeuia, testacea, apice retusa, serrata. Corpus et pedes testacea.

*rufipollis.* 18. B. ater, nitidus, thorace rufo, elytris retusis, bidentatis.

Habitat in America meridionali. Mus. D. de Seheſtedt.

Corpus paruum, cylindricum, atrum. Thorax laeuis, rufus, nitidulus. Elytra laeuia, atra, nitida, apice retusa, margine bidentato.

*compressicornis.* 19. B. glaber, ater, nitidus, elytris marginato-retusis, antennis elongatis, compressis.

Habitat in America meridionali. D. Smidt. Mus. Dom. Lund.

Paruus, ater, glaber, nitidus. Elytra laeuia, apice retusa, margine suturaque eleuatis. Antennae singulares, elongatae, compressae, flauae. Corpus atrum, pedibus flauis.

Alium in eodem museo a D. Smidt missum vidi antennis breuibus. Forte sexus varietas.

*ferrugineus.* 20. B. glaber, ferrugineus, nitidus, elytris subrotundis, integris.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Cor-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Corpus paruum, cylindricum, ferrugineum, nitidum. Elytra ſublaeuia, parum retuſa, inermia.
Varietas nigra, vix diſtincta.

21. B thorace antice criſtato, ferrugineus, elytris ſubretuſis, nigris *criſtatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura parua pracedentis. Thorax laeuis, ferrugineus, nitidus, antice criſta parua, eleuata, dentata. Elytra baſi laeuia, ferruginea, apice retuſa, ſubſtriata, nigra.

22. B. fuſcus, elytris apice retuſis, ante apicem vnidentatis. Ent. ſyſt. 2. 368. 24. *bidens.*
Payk. Fn. Sv. 3. 148. 5.
Panz. Fn. Germ. 39. tab. 21.
Habitat in Kiliae ligno.

23. B. ater, elytris ſtriatis, piceis, capite excauato. *frontalis.*
Habitat in Carolina. Muſ. Dom. Boſc.
Paruus. Caput valde excauatum, vel potius canaliculatum, dente paruo, vti videtur, vtrinque eleuato. Thorax rotundatus, punctatus, ater. Elytra crenato-ſtriata, picea. Pedes picei.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

## 152. HYLESINVS.

*Antennae* claua, ſolida, acuminata, roſtro obſoleto inſedentes.

*Scolytus.* 1. H. glaber, nigricans, elytris truncatis, abdomine retuſo, fronte villoſa, cineraſcente.

*Boſtrichus Scolytus.* Ent. ſyſt. 2. 366. 9. ♂

*Ekkoptogaſter Scolytus.* Herbſt. Coleopt. 5. tab. 49. fig. 1. 2.

*Scolytus.* Geoff. Inſ. 1. 350. 1. tab. 5. fig. 5.

Payk. Fn. Sv. 3. 150. 7.

Sulz. Hiſt. Inſ. tab. 2. fig. 13.

Schaeff. Icon. tab. 112. fig.

~~Panz. Fn. Germ. 15. tab. 6.~~

Habitat in Angliae vlmis. Dom. Lee.

Paullo minorem totum nigrum ex America meridionali miſit Dom. Smidt, vix tamen diſtinctum.

*crenatus.* 2. H. glaber, ater, elytris crenato-ſtriatis.

*Boſtrichus crenatus.* Ent. ſyſt. 2. 366. 10. ♂

Payk. Fn. Sv. 3. 150. 8.

Panz. Fn. Germ. 15. tab. 7.

Herbſt. Col. 5. tab. 48. fig. 10.

Habitat in Germania. Dom. Schneider.

*Fraxini.* 3. H. griſeus, elytris nigro-reticulatis.

Panz. Fn. Germ. 66. 15.

Ha-

---

*Hyleſini* corpus paruum, cylindricum, pubeſcens, immarginatum, tardum; *capite* ouato, roſtro breuiſſimo, fere obſoleto, inſerto; *antennis* breuibus: articulo primo longiſſimo, vltimo craſſiori, apice acuminato, inſerti ad roſtri baſin; *oculis* magnis, haud prominulis, reniformibus, lateralibus; *thorace* cylindrico, antice parum anguſtato; *elytris* rigidis, fornicatis, arcte includentibus, longitudine abdominis; *pedibus* breuibus, compreſſis; *tibiis* extus craſſioribus, ſaepius dentatis; *tarſis* quadriarticulatis; *colore* obſcuro.

Hyleſini genus haud rite determinatum. Character naturalis generis adhuc deſideratur.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Auſtriae Fraxino in truncis gregatim tubercula efficiens. Dom. Scheidler.

Diſtinctus omnino ab H. vario, duplo fere maior. Caput griſeum, vertice nigro. Thorax griſeus, annulo dorſali nigro. Elytra ſubſtriata, griſea, nigro-reticulata. Pedes nigri, tarſis griſeis.

4. H. flaueſcens, thorace elytrisque fuſco-variis. *varius.*

Habitat in Germaniae ligno. Dom. Smidt.

Medius. Caput cineraſcens, antennarum claua acuta. Thorax flaueſcens, annulo dorſali fuſco. Elytra laeuia, flaua, fuſco-variegata. Corpus flaueſcens.

5. H. villoſus, nigricans, tibiis quatuor poſticis ſerratis. *ligniperda.*

*Boſtrichus ligniperda.* Ent. ſyſt. 2. 367. 14. *

Payk. Fn. Sv. 3. 149. 7.

Habitat in Germaniae ligno. Dom. Hybner.

6. H. villoſus, niger, elytris breuibus, piceis. *abietinus.*

*Boſtrichus abietinus.* Ent. ſyſt. 2. 367. 15. *

Habitat in Germaniae Abiete. Dom. Smidt. III. 59

7. H. villoſus, piceus, pedibus teſtaceis. *villoſus.*

*Boſtrichus villoſus.* Ent. ſyſt. 2. 367. 16. *

Payk. Fn. Sv. 3. 154. 13.

Panz. Fn. Germ. 15. tab. 8. II. 45.

Herbſt. Coleopt. 5. 121.

Habitat in Germaniae arboribus. Dom. Helwig.

8. H. fuſcus, thorace lineis, elytris maculis albis. *maculatus*

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

Magnus. Caput cinereum. Thorax fuſcus, linea dorſali lateralique cinereis, et in hac macula magna, fuſca. Elytra laeuia, fuſca, maculis cinereis baſeos: futurali longiori, apicis ſublunata. Corpus cinereum.

9.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*piniperda.* 9. H. ſubuilloſus, niger, elytris crenato-ſtriatis, plantis rufis.

*Boſtrichus piniperda.* Ent. ſyſt. 2. 367. 17.

*Dermeſtes piniperda.* Linn. ſyſt. Nat. 2. 562. 9. Fn. Sv. 421.

Degeer Inſ. 4. tab. 6. fig. 8. 9.

Panz. Fn. Germ. 15. tab. 9.

Herbſt. Col. 5. tab. 48. fig. 4.

Payk Fn. Sv. 3. 152. 11.

Habitat in ramiculis Pini inferioribus, quos perforat, exſiccat, deſtruit.

*aeneipennis* 10. H. glaber, ater, elytris aeneis, nitidis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Maior. Antennae claua compreſſa, nigrae. Caput et thorax glabra, laeuia, atra, nitida. Corpus atrum, pedibus valde compreſſis.

Variat ſexu antennis longioribus piloſis, frontequeque ferrugineo-villoſa.

*faſciatus,* 11. H. ſubuilloſus, teſtaceus, elytris baſi ſuturaque nigris.

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

Paullo maior H. teſtaceo. Caput et thorax pubeſcentia, teſtacea, immaculata. Elytra laeuia, integra, teſtacea, baſi ſuturaque fuſcis. Pedes breues, compreſſi, teſtacei.

Variat ſexu capite retuſo.

*gibbus,* 12. H. glaber, gibbus, obſcure-rufus, nitidus, elytris teſtaceis.

Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.

Statura omnino praecedentis. Caput et thorax glabra, laeuia, obſcure-rufa, nitida. Elytra vix ſtria-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſtriata, teſtacea. Corpus nigrum, pedibus teſtaceis.

3. H. glaber, obſcure-rufus, elytris baſi ſuturaque nigris. *futuralis.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentium. Caput et thorax laeuia, ruſa, nitida. Elytra vix ſtriata, rufa, baſi ſuturaque nigris. Corpus nigrum, pedibus rufis.

14. H. glaber, teſtaceus, elytris laeuibus. *teſtaceus.*

*Boſtrichus teſtaceus.* Ent. ſyſt. 2. 367. 18.

Panz. Fn. Germ. 66. tab. 16.

Habitat in Pini truncis. Dom. Zſchuck.

15. H. niger, elytris ſtriatis. *niger.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura omnino H. ſuturalis. Caput et thorax laeuia, glabra, atra. Elytra ſtriata, nigra. Corpus nigrum, pedibus compreſſis, breuibus.

Variat ſexu: capite retuſo, ferrugineo, villoſo.

16. H. glaber, ater, elytris crenato-ſtriatis. *breuipes.*

Habitat in Amboina. Dom. Billardiere.

Statura omnino H. chloropodis, at duplo maior. Corpus cylindricum, atrum. Antennae breues, claua cineraſcente. Thorax vix punctatus. Elytra crenato-ſtriata. Pedes breues, femoribus compreſſis.

17. H. niger, elytris ſtriatis, antennis pedibusque rufis. *chloropus.*

*Curculio chloropus.* Ent. ſyſt. 2. 471. 323.

Linn. ſyſt. Nat. 2. 617. 82. Fn. Sv. 635.

Payk. Monogr. 116. 110.

Panz. Fn. Germ. 10. tab. 14.

Bonsd.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Bonsd. Curc. ſuec. 2. 18. fig. 19.
Habitat in Europae ligno quercino.

*pubeſcens.* 18. H. villoſus, niger, antennis pedibusque flauescentibus, fronte villoſa.
*Boſtrichus pubeſcens.* Ent. ſyſt. 2. 368. 19. *
Panz. Fn. Germ. 15. tab. 10.
Herbſt. Coleopt. 5. 121.
Habitat in Germaniae ligno. Dom. Prof. Helwig.

*ater.* 19. H. glaber, ater, nitidus, elytris laeuibus, pedibus piceis.
*Boſtrichus ater.* Ent. ſyſt. 2. 368. 20. *
*Boſtrichus anguſtatus.* Herbſt. Col. 5. 111. 9. tab. 48. fig. 9.
Payk. Fn. Sv. 3. 153. 12.
Oliv. Inſ. 2. 18. 9. 10. tab. 2. fig. 16.
Habitat in Sueciae ligno. D. de Paykull.

*voluulus.* 20. H. glaber, teſtaceus, ore anoque nigris.
*Boſtrichus voluulus.* Ent. ſyſt. 4. Append. 454. 18. *
Habitat in Americae ligno. D. v. Rohr.

*melanocephalus.* 21. H villoſus, griſeus, capite nigro, pedibus flaueſcentibus.
*Boſtrichus melanocephalus.* Ent. ſyſt. 2. 368. 21. *
Habitat in Daniae ligno. Muſ. D. Lund.

*flauus.* 22. H. glaber, flauus, nitidus.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Corpus paruum cylindricum, totum flauum, immaculatum, nitidum.
*Variat* forte ſexu: capite retuſo.

*oleiperda.* 23. H. villoſus, fuſcus, elytris ſtriatis griſeis, pedibus teſtaceis.

Bo-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Boſtrichus oleiperda.* Ent. ſyſt. 2. 366. 11. *
Habitat in Galliâ meridionali, Oleam deſtruens.
Muſ. Dom. Boſc.

4. H. villoſus, griſeus, antennarum claua elongata. *Oleae.*
*Boſtrichus Oleae.* Ent. ſyſt. 2. 366. 12. *
Habitat in Galliae meridionalis Olea. Muſ. D. Boſc.

5. H. ater, nitidus, elytris rufis, abdomine retuſo. *pygmaeus.*
*Boſtrichus pygmaeus.* Ent. ſyſt. 2. 367. 13. *
Habitat Halae Saxonum. Dom. Hybner.

6. H. ſubuilloſus, fuſcus, elytris vitta abbreuiata cinerea. *vittatus.*
*Boſtrichus vittatus.* Ent. ſyſt. 2. 368. 22. *
Habitat in Holſatiae ligno. D. Daldorff.

7. H. glaber, ater, immaculatus. *minutus.*
*Boſtrichus minutus.* Ent. ſyſt. 2. 368. 23. *
Panz. Fn. Germ. 15. tab. 11.
Habitat in Kiloniae Boleto verſicolore.

8. H. cinereus, elytris laeuibus obſcurioribus. *minimus.*
Habitat ſub Abietis cortice Saxoniae. D. Hybner.
Minutus. Thorax laeuis, cinereus. Elytra integra, obſcuriora.
*Variat* thorace nigricante; ſorte aetate.

29. H. thorace eleuato, gibbo, niger, obſcurus, pedibus flauis. *obſcurus.*
Habitat in Americae meridionalis ligno. D. Smidt.
Muſ. D. de Seheſtedt.
Minutus. Corpus totum obſcurum, pedibus ſolis flauis. Thorax gibbus.

30. H. cylindricus, ater, pedibus rufis, thorace canaliculato. *canaliculatus.*
*Curculio canaliculatus.* Ent. ſyſt. 2. 471. 324. *
Habitat in Americae meridionalis Inſulis. D. Smidt.

153. BRV-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

153. BRVCHVS. *Palpi* aequales, filiformes.
*Maxilla* membranacea, bifida.
*Ligula* acuminata.
*Antennae* filiformes.

*ſpinoſus.* 1. B. griſeus, thorace elytrisque ſpinoſis. Ent. ſyſt. 2. 369. 1. *
Habitat in Iamaica. D. Drury.

*nucleorum* 2. B. cinereus, elytris ſtriatis, femoribus poſticis ouatis, dentatis. Ent. ſyſt. 2. 369. 2. *
Herbſt. Arch. tab. 20. fig. 16.
Habitat in Indiae nucleis.

*Bactris.* 3. B. cinereus, elytris ſublaeuibus, femoribus poſticis ouatis, tibiis incuruis. Ent. ſyſt. 2. 369. 3. *
Linn. ſyſt. Nat. 2. 605. 4. Amoen. acad. 6. 392. 6.
Habitat in Bactris nucleis.

*vmbellatarum.* 4. B. ſquamoſus, ſupra griſeus, ſubtus cinereus. Ent. ſyſt. 2. 370. 4. *
Habitat in Barbariae vmbellatis. D. Prof. Vahl.

*Piſi.* 5. B. elytris nigris, albo-maculatis, podice albo: punctis duobus nigris. Ent. ſyſt. 2. 370. 5. *
*Bruchus Piſi.* Linn. ſyſt. Nat. 2. 604. 1. Muſ. Lud. Vlr. 35.
Geoff. Inſ. 1. 267. 1. tab. 4. fig. 9.
Vl. 37. Ledermyll. microſc. tab. 99. fig. 2.
Degeer Inſ. 4. tab. 16. fig. 3. 4.
Scop.

*Bruchi* corpus paruum, ouatum, gibbum, glabrum, inmarginatum, tardum; *capite* inflexo, obtuſo, ouato, inſerto; *oculis* rotundatis, lateralibus, vix prominulis: *antennis* breuibus, ſerratis, ante oculos inſertis; *thorace* ouato, gibbo; *ſcutello* paruo, triangulo; *elytris* rigidis, fornicatis, abdomine breuioribus, ano prominulo; *pedibus* breuibus, validis; *tarſis* quadriarticulatis; *colore* vario, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Scop. Ent. carn. 63.
Panz. Fn. Germ. 66. tab. 11.
Habitat in America boreali et Europa auſtrali, piſa deſtruens. Deuoratur a Gracula quiscula, Coruis, Picis aliisque.

. B. corpore fuſco, griſeo-maculato. Ent. ſyſt. 2. 370. 6. * *Cacao.*
Habitat in Theobromatis ſeminibus.

. B. fuſcus, thorace elytrisque cinereo-punctatis. Ent. ſyſt. ſuppl. 158. 6. * *punctatus.*
Habitat - - -

. B. griſeo-nitidulus, elytris vix abdomine breuioribus. Ent. ſyſt. 2. 370. 7. * *Robiniae.*
Habitat in Robiniae pſeudoacaciae ſeminibus Americae borealis.

. B. obſcure ferrugineus, elytris fuſco-maculatis, femoribus poſticis elongatis, clauatis. Ent. ſyſt. ſuppl. 158. 7. * *femoratus.*
Habitat in Indiae orientalis ſeminibus. D. Daldorff.

0. B. thorace gibboſo cupreo, elytris nigris tuberculato-ſpinoſis. Ent. ſyſt. 2. 370. 8. * *gibboſus.*
Habitat in America boreali.

1. B. elytris griſeis, nigro-maculatis, pedibus rufis, ſcutello albo. Ent. ſyſt. 2. 371. 9. * *Theobromatis.*
*Bruchus Theobromae.* Linn. ſyſt. Nat. 2. 605. 2.
Habitat in Theobromatis ſeminibus.

2. B. niger, elytris cinereis: maculis tribus nigris, margine connexis. Ent. ſyſt. 2. 371. 10. * *marginellus.*
Habitat in Germania. D. de Hattorff.

3. B. niger, elytris ſtriatis, ferrugineis: maculis tribus nigris, margine connexis. Ent. ſyſt. ſuppl. 158. 10. * *3maculatus.*

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in India orientali. Dom. Hybner.

*anilis.* 14. B. ſubferrugineus, elytris apice atris, ano atr linea apicis alba. Ent. ſyſt. 2. 371. 11. *

Habitat in India orientali. Muſ. D. Banks.

*ſparſus.* 15. B. cinereus, elytris albo-punctatis, femorib poſticis punctatis.

Habitat in Tanger. Muſ. D. de Sebeſtedt.

Medius. Antennae cinereae, baſi rufae. Thor cinereus, macula magna baſeos alba. Elyt ſtriata, cinerea: punctis albis ſparſis. An prominet maculis duabus fuſcis. Pedes cine femoribus poſticis acute dentatis.

*2 puſtulatus.* 16. B. niger, elytris ſtriatis: puncto rufo.

Habitat in America meridionali. D. Smidt. Muſ D. Lund.

Statura et magnitudo B. 2 punctati. Antenna craſſae, ſerratae. Caput et thorax atra, imma culata. Elytra ſtriata, nigra, puncto magn rufo. Corpus cinereo-tomentoſum, podice pro minulo, immaculato. Pedes nigri, femoribu poſticis rufis.

*2 punctatus.* 17. B. cinereus, elytris fuſcis: puncto baſeos ocel lari atro. Ent. ſyſt. 2. 371. 12. *

Sulz. Hiſt. Inſ. tab. 4. fig. 2.

Habitat in Heluetiae plantis.

*4 maculatus.* 18. B. thorace nigro: macula baſeos pallida, elytri rufeſcentibus: macula apiceque nigris. Ent ſyſt. 2. 371. 13. *

Habitat in Inſula St. Croix Americae. Dr. Pflug

*arthriticus.* 19. B. fuſcus, femoribus poſticis incraſſatis, com preſſis.

Habitat in Carolina. Muſ. D. Boſc.

Statura et magnitudo B. Piſi. Antennae ſerratae fuſcae. Elytra ſubſtriata. Femora poſtica ova ta, valde incraſſata, compreſſa.

20.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

20. B. cinereo-villoſus, elytris ſtriatis, femoribus poſticis incraſſatis, compreſſis. Ent. ſyſt. ſuppl. 159. 13. * *gonagra.*

Habitat in India orientali. D. Daldorff.

21. B. teſtaceo fuſcoque varius, elytris punctato-ſtriatis, femoribus poſticis compreſſo-clauatis, vnidentatis. *podagricus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Minor quam B. gonagra, et diſtinctus. Antennae extrorſum ſubcraſſiores, teſtaceae. Caput et thorax obſcura, fuſco-ſublineata. Elytra punctato-ſtriata, teſtacea, lituris aliquot fuſcis albidisque obſoletis. Pedes teſtacei, femoribus poſticis valde compreſſis, incraſſatis, acute vnidentatis.

22. B. ferrugineus, thoracis macula baſeos ſcutelloque albis. Ent. ſyſt. 2. 372. 14. * *ſcutellaris.*

Habitat - - - Muſ. D. Lund.

23. B. ferrugineus, cinereo-villoſus, antennarum articulis tribus vltimis incraſſatis. Ent. ſyſt. ſuppl. 159. 14. * *craſſicornis.*

Habitat in Indiae orientalis feminibus. D. Daldorff.

24. B. elytris nigris: atomis albis, femoribus poſticis vnidentatis. Ent. ſyſt. 2. 372. 15. * *granarius.*

*Bruchus granarius.* Linn. ſyſt. Nat. 2. 605. 5. Fn. Sv. 628. *vi. 37.*

Payk. Fn. Sv. 3. 157. 1.

Habitat in plantarum feminibus.

25. B. niger, ſcutello elytrorumque faſcia poſtica albis. *Erui.*

Habitat in Hiſpaniae Eruo, Muſ. D. de Seheſtedt.

Sta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo B. granarii. Caput et thorax nigra, puncto ad baſin thoracis albo. Scutellum album. Elytra vix ſtriata, nigra, atomis faſciaque maculari albis. Pedes ferruginei: poſtici nigri, femoribus muticis.

*Ciſti.* 26. B. ater, immaculatus, femoribus muticis. Ent. ſyſt. 2. 372. 16. *

Panz. Fn. Germ. 66. tab. 12.

Payk. Fn. Sv. 3. 159. 3.

Habitat in floribus Ciſti Helianthemi. Muſ. Dom. Banks.

*ſuturalis.* 27. B. niger, ſutura elytrorum flaueſcente. Ent. ſyſt. 2. 372. 17. *

Habitat Pariſiis. Muſ. D. Boſc.

*abdominalis.* 28. B. niger, abdomine cinereo-maculato, pedibus anticis teſtaceis. Ent. ſyſt. 2. 372. 18. *

Habitat in India orientali. Muſ. D. Banks.

*ſeminarius* 29. B. ater, antennarum baſi pedibusque anticis teſtaceis, femoribus muticis. Ent. ſyſt. 2. 373. 19. *

*Bruchus ſeminarius.* Linn. ſyſt. Nat. 2. 605. 6.

Geoff. Inſ. 1. 268. 3.

Habitat in Angliae floribus.

*villoſus.* 30. B. villoſus, cinereus, immaculatus. Ent. ſyſt. 2. 373. 20. *

Habitat in Germania. D. Smidt.

*nigricornis.* 31. B. villoſus, cinereus, antennis pedibusque nigris.

Habitat in Isle de France. D. Daldorff. Muſ. D. de Seheſtedt.

Statura parua B. villoſi, at alius. Antennae extrorſum parum craſſiores, nigrae. Caput, thorax elytraque ſtriata, cinerea, immaculata. Corpus cinereum, pedibus nigris.

32.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:25 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

32. B. niger, obſcurus, antennis pedibusque rufis, femoribus poſticis incraſſatis nigris. Ent. ſyſt. 2. 373. 21. * *rufipes.*

*Bruchus rufipes.* Oliv. Inſ. tab. fig.

Habitat Pariſiis. Muſ. D. Boſc.

33. B. obſcurus, elytris ſtriatis, vitta media ferrugi- nea. Ent. ſyſt. ſuppl. 159. 21. * *vittatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

34. B. elytris punctatis, teſtaceis, nigro-maculatis, podice griſeo: punctis binis nigris. Ent. ſyſt. 2. 373. 22. * *maculatus*

Habitat in America. D. v. Rohr.

35. B. teſtaceus, cinereo-villoſus, pedibus ſimpli- cibus. Ent. ſyſt. ſuppl. 159. 22. * *capſincola.*

Habitat in Cayennae feminibus.

36. B. teſtaceus, immaculatus, elytris ſubſtriatis, femoribus poſticis acute dentatis. Ent. ſyſt. 2. 374. 23. * *Mimoſae.*

Habitat in Mimoſae feminibus Americae meridionalis. Muſ. D. Lund.

37. B. cinereus, elytris ſtriatis, pallidis, fuſco-maculatis, pedibus teſtaceis. *minutus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Minutus. Caput et thorax villoſa, cinerea, immaculata, maculis quinque fuſcis 2. 2. 1. Corpus cinereum, pedibus teſtaceis.

38. B. antennis pectinatis, flauis, corpore atro, pedibus teſtaceis. Ent. ſyſt. 2. 374. 24. * *flauicornis*

Habitat Halae Saxonum. Dom. Hybner.

39. B. antennis pectinatis, corpore longioribus, corpore griſeo, fuſco, maculato. Ent. ſyſt. 2. 374. 25. * *ſerraticornis.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717 Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Oriente. Prof. Forſkahl.

*Coryphae.* 40. B. antennis pectinatis, fuſcis, thorace elytrisq fuſcis.

Habitat in Carolina. Muſ. Dom. Boſc.

Statura minimo B. ferraticornis. Antennae inp mis maris valde pectinatae, nigrae. Caput grum. Thorax punctatus, obſcure-rufus, i maculatus. Elytra punctato-ſtriata, laetius r fa. Corpus nigrum. Abdomen cinereum, poſ ce prominet punctis duobus paruis, fuſcis.

*2guttatus.* 41. B. thorace nigro, lineolis duabus baſeos, ſcute loque albis, elytris baſi rufis, antennis flabe latis.

Habitat in Carolina. Muſ. Dom. Boſc.

Paruus. Antennae corpore paullo breuiores, fa bellatae, nigrae, baſi rufae. Thorax niger, l neolis duabus approximatis, eleuatis baſeo albis. Scutellum album. Elytra ſtriata, baſi ru fa, apice nigra. Pedes rufi, tibiis nigris.

*2ſtriatus.* 42. B. thorace nigro, lineolis duabus baſeos approxi matis ſcutelloque albis, elytris rufis: ſtrig duabus albidis.

Habitat in America meridionali. D. Smidt. Mu D. Lund.

Affinis videtur B. 2 guttato. Antennae corpor paullo breuiores, nigrae. Caput nigrum. Tho rax niger, lineolis duabus baſeos eleuatis, ap proximatis, albis. Scutellum album. Elytra ſtriata, ferruginea, ſtrigis duabus approximatis albis. Pedes rufi, femoribus nigris.

Variat ſexu: antennis ſimplicibus.

*pectinicor-nis.* 43. B. antennis pectinatis, corpore ferrugineo, lon gioribus. Ent. ſyſt. 2. 374. 26.

*Bruchus pectinicornis.* Linn. ſyſt. Nat. 2. 605. 5

Degeer

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inf. 5. 281. 2.
Schaeff. Elem. tab. 86. fig. 1.
Habitat in Chinae feminibus.

4. B. antennis pectinatis, niger, elytris ſtriatis, ferrugineis, femoribus poſticis vnidentatis. *barbicornis.*
Habitat - - - Muſ. D. de Seheſtedt.
Statura et magnitudo praecedentis. Antennae pectinatae, nigrae, baſi ferrugineae. Caput et thorax nigra, macula ad baſin thoracis triangulari alba. Elytra ſtriata, ferruginea, immaculata. Corpus nigrum. Pedes ferruginei, pedibus poſticis vnidentatis.

5. B. nigricans, antennis pectinatis pedibusque ferrugineis. *flauipes.*
Habitat - - -
Praecedentibus paullo maior. Antennae magnae, ferrugineae, valde pectinatae: articulo vltimo ouato. Caput et thorax obſcura, nigra, immaculata. Elytra ſtriata, obſcura. Pedes inermes, ferruginei.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

154. ANTHRIBVS. *Palpi* aequales, filifo mes.
*Maxilla* bifida.
*Antennae* articulis fu aequalibus, infide tes roſtro breui, pl no.

*heros.* 1. A. griſeus, elytris punctato-ſtriatis, fuſco-m culatis.
Habitat in Guinea. Dom. Heſſe. Muſ. D. Lund.
Magnus in hoc genere. Antennae maris longit dine corporis, nigrae, articulo 8. 9. apice cin reo. Caput griſeum, obſcurum. Thorax g ſeus, lineolis duabus fuſcis. Elytra punctat ſtriata, maculis duabus magnis, marginalibu fuſcis. Pedes nigri, cinereo-varii.
Femina antennis breuioribus, ſubfiliformibus: a ticulis vltimis parum craſſioribus, compreſſis.

*gigas.* 2. A. griſeus, atomis albis, antennis longiſſimis.
Habitat in Guinea. D. Meier. Muſ. D. de Sehe ſtedt.
Antennae corpore duplo longiores, obſcurae: a ticulis vltimis incraſſatis: vltimo tamen ſetaceo. Roſtrum longiuſculum, canaliculatum Thorax et elytra griſea, obſcura, atomis ſpar ſis, albis. Pedes inermes, cinerei.

Va

---

*Anthribi* corpus ouatum, gibbum, ſubglabrum, immarginatum tardum; *capite* ouato, antice porrecto, plano, obtuſo, i ſerto; *oculis* globoſis, prominulis, marginali us; *antenn* longiuſculis, in fouea laterali ſub oculis inſertis; *thorace* r tundato: marginibus deflexis; *ſcutello* paruo, rotundato *elytris* rigidis, fornicatis, apice deflexis, longitudine abdom nis; *pedibus* breuibus, ſvalidis; *tarſis* quadriarticulatis; *co lore* vario, ſaepius obſcuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Variat paullo minor, elytris albo-maculatis.
Femina antennis breuibus.

A. cinereo nigroque varius, antennis atris apice albis. *atricornis.*

Habitat in Guinea. Dom. Heſſe. Muſ. D. Lund.

Statura praecedentium. Antennae in noſtro ſpecimine breues, atrae, nodoſae, nitidae. Caput et thorax ſcabra, fuſca, cinereo-varia. Elytra punctato-ſtriata, nigra, cinereo-varia. Corpus et pedes variegata.

A. griſeus, cinereo fuſcoque varius, antennarum claua, articulo vltimo acuto. *acuticornis*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Maior A. perfolicornis et alius. Antennae nigrae, ante apicem albidae: articulis tribus vltimis incraſſatis; vltimo acuto, breuiori. Caput et thorax griſea, fuſco-maculata, thoracis ſtriga poſtica clauata. Elytra punctato-ſtriata, griſea, cinereo fuſcoque variegata. Pedes variegati.

griſeus, antennis corpore longioribus. Ent. ſyſt. ſuppl. 160. 1. * *longicornis*

Habitat in India orientali. Muſ. D. de Seheſtedt.

A. ater, elytris ſulcatis, ferrugineo-maculatis. *puſtulatus.*

Habitat in America meridionali. D. Smidt. Muſ. Dom. Lund.

Magnus. Antennae corpore breuiores, nigrae. Caput atrum, bifaſciatum: ſulcis villoſo-albis. Thorax ater, ferrugineo-maculatus. Elytra dor-

---

Antennae in hoc genere et ſpecie et ſexu nimis differunt. Mox ſetaceae, articulo vltimo ſubulato, mox articulis tribus ouatis, craſſioribus, vltimo ſetaceo, mox claua ouata, compreſſa, obtuſa, mox moniliformes, extrorſum craſſiores: articulo vltimo ouato, acuto.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

dorſo plana, ſulcata, atra, maculis plurimis, ſparſis ferrugineis. Corpus atrum, albido-ſtrigoſum. Pedes atri.

*pilicornis.* 7. A. cinereus, thorace elytrisque fuſco-variis, antennis piloſis.

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

Corpus magnum, cinereum, vix maculatum. Antennae breues, vtrinque piloſae: articulo primo ſecundoque nudis. Caput cinereum. Thorax et elytra cinereo fuſcoque varia. Pedes cinerei.

*pubicornis.* 8. A. fuſcus: linea dorſali lata, cinerea, antennis piloſis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentis. Antennae vtrinque piloſae. Caput fuſcum, fronte cinerea. Thorax fuſcus, linea dorſali lata, cinerea, apice parum coarctata. Elytra vix ſtriata, fuſca: linea lata, dorſali, cinerea, baſi medio apiceque coarctata, punctis aliquot nigris. Pedes fuſci.

*barbicornis.* 9. A. cinereus, elytris fuſco-maculatis, antennis piloſis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Antennae piloſae, corpore paullo breuiores, nigrae. Caput et thorax cinerea, fuſco-maculata. Elytra punctato-ſtriata, cinerea, fuſco-maculata, tuberculoque paruo baſeos eleuata. Pedes cinerei, femoribus poſticis, faſcia atra.

*compreſſicornis.* 10. A. obſcurus, albo-varius, elytris punctis elevatis, atris, antennarum clava ouata, compreſſa.

Habitat in Guinea. Muſ. Dom. Lund.

Sta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura omnino et magnitudo praecedentium. Antennae ſingulares, craſſae, corpore breuibres, atrae, articulis baſi cinereis, apice claua ouata, compreſſa. Caput et thorax fuſco cinereoque varia. Elytra concolora, ſubſtriata, punctis aliquot eleuatis, atris, ſparſis. Corpus cinereum.

11. A. cinereo fuſcoque varius, elytris planis, laeuibus, antennis breuibus: claua ouata, compreſſa. *clauicornis*

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

Statura praecedentis. Antennae atrae, claua ouata, compreſſa. Caput villoſum, cinereum. Thorax et elytra cinereo fuſcoque variegata. Corpus cinereum.

12. A. thorace fuſco: punctis quatuor albis, elytris variegatis, antennis claua ouata, compreſſa. *craſſicornis.*

Habitat in Sumatra. D. Daldorff.

Praecedentibus minor. Antennae fuſcae, ante clauam albae, claua ipſa compreſſa, atra. Caput fuſcum, cinereo-varium. Thorax fuſcus, punctis quatuor albis. Elytra ſtriato-punctata, fuſca, albo-varia.

13. A. fuſco cinereoque varius, elytris punctato-ſtriatis, antennarum claua compreſſa. *perfolicornis.*

Habitat in Sumatra. D. Daldorff. Muſ. D. Lund.

Statura et affinitas A. clauicornis. Antennae breues, claua ouata, compreſſa. Caput et thorax fuſca, cinereo-varia, thorace poſtice ſtriga parua, eleuata. Elytra punctato-ſtriata, fuſca, cinereo-varia. Corpus atrum, pedibus cinereis.

14. A. niger, antennis elongatis, tenuibus: claua ouata, compreſſa. *tenuicornis.*

Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura omnino praecedentium. Antennae longitudine corporis, nigrae, articulis apice incraſſatis, vltimis tribus clauam ouatam, compreſſam, obtuſam conſtituentibus. Roſtrum breue, planum. Thorax et elytra nigra, punctis aliquot eleuatis ſcabra. Corpus nigrum.

*Albinus.* 15. A. niger, fronte anoque albis, thorace tuberculato. Ent. ſyſt. 2. 375. 1.
*Curculio Albinus.* Linn. ſyſt. Nat. 2. 616. 79.
Fn. Sv. 632.
Degeer Inſ. 5. 255. 44. tab. 8. fig. 1.
Payk. Monogr. 118. 112.
Fn. Sv. 3. 160. 2.
I 47. Bonsd. Curcul. 16. 1.
Habitat in Europa boreali.

*latiroſtris.* 16. A. roſtro latiſſimo, plano, elytris apice albis: punctis duobus nigris. Ent. ſyſt. 2. 376. 2.
*Platyrrhinus coſtiroſtris.* Ent. Helv. 1. 114. tab. 14. fig. 1. 2.
Geoff. Inſ. 1. 307. 8. tab. 5. fig. 8.
Payk. Monogr. 117. 111.
Fn. Sv. 3. 159. 1.
Sulz. Hiſt. Inſ. tab. 2. fig. 17.
Schaeff. Icon. tab. 89. fig 6.
Roſſ. Fn. Etr. 1. 132. tab. 5. fig. 5.
II 47 Panz. Fn. Germ. 15. tab. 12.
Bonsd. Curc. ſuec. 2. tab. 1. fig. 2.
Habitat in Europa boreali.

*albiroſtris.* 17. A. roſtro latiſſimo, plano, elytrisque poſtice albis nigro-variis. Ent. ſyſt. 2. 376. 3.
*Curculio albiroſtris.* Payk. Monogr. 120. 113.
Payk. Fn. Sv. 3. 162. 3.
II 47. Panz. Fn. Germ. 15. tab. 13.
Schall. Act. Hall. 1. 287.

Herbſt.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Arch. 80. 6. tab. 24. fig. 26.
Habitat in Saxoniae Betula. Dom. Hybner.

18. A. roſtro latiſſimo, plano, elytrorumque apicibus anoque albis. Ent. ſyſt. ſuppl. 160. 4. *Anthribus breuiroſtris.* Panz. Fn. Germ. 57. tab. 9. *niueirostris.*

Habitat Halae Saxonum. Dom. Hybner.

19. A. cinereus, thorace elytrisque ſpinoſis, antennarum claua compreſſa. *ſpinoſus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Magnitudo omnino A. albiroſtris. Antennae corpore breuiores, rufae: claua ouata, compreſſa, nigra. Caput cinereum. Thorax cinereus, ſpinis duabus anticis tribusque in medio eleuatis, acutis. Elytra vix ſtriata, ſpinis tribus ad ſuturam eleuatis. Corpus cinereum.

20. A. fuſcus, thorace inaequali, elytris ſpinoſis. *aculeatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentis. Antennae breues, fuſcae: articulis tribus vltimis craſſioribus, diſtinctis; vltimo orbiculato. Caput planum, fuſcum. Thorax inaequalis, fuſcus. Elytra fuſca, ſpinis tribus ad ſuturam maioribus, eleuatis, acutis aliquot minoribus ſparſis. Pedes teſtacei, fuſco-annulati.

21. A. obſcurus, capite, thoracis antico, elytrorumque macula obſcure-griſeis. *lunatus.*

Habitat in Carolina. Muſ. Dom. Boſc.

Caput planum, rotundatum, obſcure-griſeum. Thorax obſcurus, antice lateribusque griſeis. Elytra ſtriata, obſcura: punctis holoſericeis, atris, maculaque magna, in medio cinerea, quae

ſae-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

saepius obsoleta. Pedes albo nigroque variegati.

Mas minor, obscurior.

*griseus.* 22. A. griseus, elytris postice cinereis. Ent. syst. 2. 377. 4. *

Habitat in noua Hollandia. Mus. D. Banks

*Roboris.* 23. A, rostro plano, latissimo, nigro, aeneus, rostro thorace pedibusque rufis. Ent. syst. suppl. 160. 4. *

*Anthribus ruficollis.* Ent. Helv. 1. 122. 1. tab. 15. fig. 4. 5.

*Curculio ruficollis.* Linn. Fn. Sv. 595.

*Attelabus ruficollis.* Herbst. Arch. 5. tab. 25. fig. 1.

Habitat in Kiliae Robore.

*planirostris* 24. A. rostro plano, latissimo, aeneus, rostro pedibusque testaceis. Ent. syst. 2. 377. 5. *

*Curculio planirostris.* Payk. Monogr. 124. 117.

Panz Fn. Germ. 15. tab. 14.

Degeer Inst. 5. 252. 42.

Habitat in Holsatiae Aesculo Hippocastano. D. Daldorff.

*nigripennis.* 25. A. rostro plano, rufus, elytris atris.

Habitat in Carolina. Mus. D. Bosc.

Statura omnino A. Roboris. Antennae rufae, articulis tribus vltimis clauatis, fuscis. Thorax rotundatus, pubescens, rufus immaculatus. Elytra punctato-striata, atra.

*collaris.* 26. A. rostro plano, niger, thorace rufo.

Habitat in Carolina. Mus. D. Bosc.

A. Roboris paullo maior. Caput atrum, rarius fronte obscure-rufa. Thorax subcylindricus, pubescens, rufus, immaculatus. Elytra punctato-striata, atra. Corpus nigrum. Pedes nunc rufi, nunc nigri.

27.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

27. A. niger, elytris elevato-striatis, rufis: punctis nigris, sparsis. Ent. syst. 2. 377. 6. ✲ *scabrosus.*
*Anthribus scabrosus.* Payk. Fn. Sv. 3. 164. 4.
*Curculio scabrosus.* Payk. Monogr. 121. 114.
*Bruchus scabrosus.* Herbst. Arch. tab. 20. fig. 15.
Geoff. Inf. 1. 306. tab. 5. fig. 3.
Panz. Fn. Germ. 15. tab. 15. II. 46
Habitat in Europae Jacea.

28. griseus, elytris punctis albis nigrisque alternis, striatis. *Coffeae.*
Habitat in Indiae Coffea. D. Daldorff. Mus. D. de Seheſtedt.
Statura et magnitudo A. scabrosi. Caput et thorax obscure-grisea. Elytra grisea, striis e punctis albis nigrisque. Corpus griseum, pedibus subtestaceis.

29. A. elytris nigris: punctis albis nigrisque alternis, striatis. Ent. syst. 2. 378. 8. ✲ *varius.*
*Anthribus varius.* Payk. Fn. Sv. 3. 165. 5.
*Curculio varius.* Payk. Monogr. 122. 115.
Geoff. Inf. 1. 307. 2. II. 48
Panz. Fn. Germ. 15. tab. 16.
Habitat in Sueciae Pinetis. Dom. de Paykull.

30. A. obscure-ferugineus, griseo-pubescens, elytris apice griseis. Ent. syst. suppl. 161. 8. ✲ *pubescens.*
Habitat Halae Saxonum. Dom. Hybner.

31. A. cinereo fuscoque varius: punctis elevatis, pilosis. Ent. syst. 2. 378. 9. ✲ *sepicola.*
Habitat in Germaniae sepibus. Dom. Schneider.

32. A. villosus, cinereus, antennis pedibusque testaceis: femoribus posticis clava nigra. *sericeus.*
Habitat in Austria. Mus. D. Lund.

Minu-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Minutus. Caput, thorax, elytra cinerea, immaculata. Antennae et pedes teſtacea, femoribus poſticis elongatis, clauatis: claua nigra.

155. BRACHYCERVS. *Palpi* quatuor, breuiſſimi, *antici* biarticulati. *Antennae* breues, extrorſum craſſiores, roſtro inſidentes.

*apterus.* 1. B. thorace ſpinoſo: cruce impreſſa, elytris ferrugineo-punctatis. Ent. ſyſt. 2. 379. 1. *
Oliv. Inſ. 3. 82. tab. 1. fig. 3.
Linn. ſyſt. Nat. 2. 619. 95.
Degeer Inſ. 5. 275. tab. 16. fig. 1.
Habitat in India.

*ocellatus.* 2. B. thorace ſpinoſo, antice excauato, elytris cinereus: punctis atris, ſubocellatis. Ent. ſyſt. 2. 379. 2. *
Oliv. Inſ. 3. 82. tab. fig.
Habitat in Madagaſcar. Muſ. Britann.

*ſcalaris.* 3. B. thorace ſpinoſo, inaequali, ater, elytris ſtriis rufis, denticulatis. Ent. ſyſt. 2. 380. 3. *
Habitat ad Cap. Bon. Spei. Muſ. Dr. Schulze.

4.

---

*Brachyceri* corpus magnum, ouatum, gibbum, glabrum, immarginatum, tardum; *capite* paruo, inſerto, antice producto in roſtrum craſſum, planum, obtuſum, inaequale; *oculis* parvis, oblongis, marginalibus; *antennis* breuibus ſub medio marginis roſtri inſertis; *thorace* transuerſo, inaequali, ſaepius ſpinoſo; *ſcutello* nullo; *elytris* connatis, rigidis, fornicatis, deflexis, longitudine abdominis; *pedibus* breuibus, validis: *tarſis* quadriarticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. B. thorace ſpinoſo, inaequali, ater, elytris rubris, punctis nigris, confertis. Ent. ſyſt. 2. 380. 4. * *obeſus.*
*Brachycerus obeſus.* Oliv. Inſ 82. tab. fig.
*Curculio tuberculatus.* Degeer Inſ. 7. 658. 63. tab. 49. fig. 8.
*Curculio oedematoſus.* Sulz. Hiſt. Inſ. tab. 4. fig 10.
*Curculio ater.* Wulf. Inſ. cap. tab. 1. fig. 11.
Herbſt. Arch. tab. 45. fig. 8.
Habitat ad Cap. Bon. Spei.

5. B. niger, thorace ſpinoſo. canaliculato, punctato, elytris punctis eleuatis, ſtriatis. Ent. ſyſt. ſuppl. 161. 4. * *cordiger.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

6. B. thorace ſpinoſo, quinqueſulcato, elytris laeuibus. Ent. ſyſt. 2. 381. 5. * *globoſus.*
*Brachycerus globoſus.* Oliv. Inſ. 3. 82. t. f.
*Curculio globoſus.* Drury Inſ. 1. tab. 32. fig. 4.
Habitat in Indiis. Dom. Drury.

7. B. thorace ſpinoſo, ſulcatoque, niger, thoracis lateribus elytrorumque maculis ferrugineis. *maculatus.*
Habitat - - - Muſ. D. de Seheſtedt.
Statura et magnitudo omnino B. globoſi. Caput nigrum, immaculatum. Thorax acute-ſpinoſus, valde canaliculatus, niger, margine ferrugineo. Elytra connata, laeuia, atra: maculis plurimis ferrugineis. Pedes ſcabri, nigri.

8. B. fuſcus, capite thoraceque cylindricis, anguſtioribus, elytris poſtice ſpinoſis. Ent. ſyſt. 2. 381. 6. * *roſtratus.*
Oliv. Inſ. 82. tab. 1. fig. 4.
Habitat in Africa aequinoctiali. Muſ. D. Banks.

9. B. obſcurus, thorace varioloſo, elytris punctatis ſpinisque triplici ſerie. *ſpiniger.*

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Guinea. Dom. Heſſe. Muſ. D. de Seheſtedt.

Magnitudo B. obeſi. Caput cinereum. Thorax punctis plurimis alte-impreſſus. Elytra punctata, obſcura, lineis tribus ſpinarum eleuatarum acutarum, armata. Pedes cinerei.

*verrucoſus.* 10. B. cinereus, thorace ſubinermi, elytris ſcabri ſpinoſisque tuberculoque poſtico eleuato.

Habitat in Guinea. Dom. Meier. Muſ. D. Lund.

Magnus. Caput inaequale, cinereum. Thorax ſub inermis, cinereus, punctis plurimis eleuatis atris, glabris. Elytra cinerea, faſcia media lata, inaequali, fuſca, punctis eleuatis, atris ſpinisque breuibus, acutis, armata. Poſtice tuberculum obtuſum, eleuatum ad ſuturam. Pedes cinerei.

Variat absque elytrorum faſcia fuſca.

*barbarus.* 11. B. thorace ſpinoſo, ſulcatoque, elytris ſcabri anguloque duplici, criſpato. Ent. ſyſt. 2. 381. 8. *

*Curculio barbarus.* Linn. ſyſt. Nat. 2. 617. 87

Habitat in Africa.

*undatus.* 12. B. thorace ſpinoſo canaliculatoque, elytris lineis eleuatis, criſpatis, corpore nigro. Ent. ſyſt. ſuppl. 161. 8. *

Habitat in Gallia, Barbaria.

*inaequalis.* 13. B. thorace ſcabro, elytris lineis eleuatis, criſpatis, capite bidentato. Ent. ſyſt. 2. 382. 9. *

Oliv. Inſ. 82. tab. 2 fig. 12.

Habitat ad Cap. Bon. ſpei. Muſ. Dom. Olivier.

*criſpatus.* 14. B. niger, thorace ſpinoſo canaliculatoque, elytris cinereis: angulo exteriori criſpato, dentato. Ent. ſyſt. 2. 382. 10. *

Habitat in Barbaria. Muſ. Dom. Desfontaines.

15.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. B. cornutus, thorace elytrisque ſubſpinoſis, elytrorum ſpinis faſciculato-piloſis. Ent. ſyſt. 2. 382. 11. * *cornutus.*

*Brachycerus cornutus.* Oliv. Inſ. 82. tab. 2. fig. 14.

*Curculio cornutus.* Linn. ſyſt. Nat. 2. 618. 91. Muſ. Lud. Vlr. 57.

Habitat in Indiis.

6. B. cinereus, thorace ſpinoſo, ſulcato, elytris angulo duplice, ſpinoſis. Ent. ſyſt. 2. 382. 12. * *algirus.*

Oliv. Inſ. 82. tab. 2. fig. 16.

Habitat in Africa. Dom. Prof. Vahl.

7. B. griſeo-fuſcus, elytris poſtice retuſis, dentatis. Ent. ſyſt. 2. 382. 13. * *retuſus.*

Oliv. Inſ. 82. tab. 1. fig. 6.

Habitat ad Cap. Bon. Spei. Muſ. Dom. Banks.

8. B. fuſcus, thorace elytrisque globoſis. Ent. ſyſt. 2. 383. 14. * *ſpectrum.*

Oliv. Inſ. 82. tab. 1. fig. 5.

Habitat ad Cap. Bon. Spei.

9. B. thorace ſpinoſo, ſulcato, niger, obſcurus, elytris lineis tribus eleuatis, crenatis. Ent. ſyſt. 2. 383. 15. * *muricatus.*

Oliv. Inſ. 82. tab. 2. fig. 19.

Habitat in Hungaria. Dom. Hybner.

Thorax lineis duabus dorſalibus eleuatis, dentatis.

10. B. thorace criſta duplici, elytris dorſo planis, reticulatis. Ent. ſyſt. ſuppl. 161. 15. * *criſtatus.*

Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

11. B. thorace punctis eleuatis ſcabro, elytris varioloſis. *griſeus.*

Habitat ad Cap. Bon. Spei. Muſ. Dom. de Seheſtedt.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura omnino B. Morionis, at minor. Corpus piloſum, griſeum, obſcurum, immaculatum. Thorax punctis eleuatis ſcaber. Elytra punctis maioribus varioloſa.

*morio.* 22. B. ater, thorace elytrisque varioloſis.
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.
Corpus paruum, atrum, immaculatum. Caput tuberculis duobus baſeos. Thorax et elytra atra, punctis impreſſis, varioloſis, reticulata. Pedes inermi, fuſci.

*vau.* 23. B. thorace ſpinoſo inaequali, elytris tuberculis eleuatis, obtuſis, numeroſis. Ent. ſyſt. 2. 383. 16.
Habitat ad Cap. Bon. Spei. D. de Paykull.

## 156. ATTELABVS. *Palpi* filiformes.
*Maxilla* bifida.
*Antennae* moniliformes, rostro inſidentes.

*Coryli.* 1. A. niger, elytris rufis, reticulatis. Ent. ſyſt. 2. 384. 1.
*Attelabus Coryli.* Linn. ſyſt. Nat. 2. 619. 1. Fn. Sv. 638.
Degeer Inſ. 5. 257. 46. tab. 8. fig. 3.
Payk. Monogr. 128. 121.
- Fn. Sv. 3. 168. 1.
Ent. Helv. 1. 118. tab. 15. fig. 1. 2.

IX. Schaef

---

*Attelabi* corpus paruum, ouatum, glabrum, immarginatum tardum; *capite* exſerto, ouato, poſtice attenuato, antporrecto; *oculis* globoſis, marginalibus, ante roſtrum inſertis; *antennis* breuibus, approximatis, roſtri baſi inſertis; *thorace* rotundato, ouato; *ſcutello* rotundato; *elytris* rotundis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis; *tarſis* quadriarticulatis; *colore* vario, plerumque obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Schaeff. Icon. tab. 75. fig. 8.
Sulz. Inſ. tab. 4. fig. 1.
β. *Attelabus Auellanae.* Linn. ſyſt. Nat. 2. 619. 2.
Geoff. Inſ. 1. 237. 11.
Schaeff. Icon. tab. 56 fig. 5. 6.
Habitat in Coryli Auellanae foliis, quae in cylindrum voluit et vtrinque claudit.

A. rufus, pectore femoribusque atris, capite poſtice attenuato, cylindrico, longitudine corporis. *cygneus.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Statura ſingularis. Magnitudo A. Coryli. Roſtrum breue, planum, apice incraſſatum, rufum. Antennae rufae: articulo primo vltimisque tribus craſſioribus, fuſcis. Caput glabrum, rufum, poſtice attenuatum, cylindricum, longitudine fere totius corporis. Thorax antice attenuatus, rufus. Elytra ſtriata, rufa. Corpus rufum, pectore femoribusque nigris.

*Varietas* forte ſexus, capite triplo breuiore, vix differt.

A. rufus, capite poſtice attenuato, cylindrico, nigro. *longicollis.*

Habitat in Auſtralaſia. D. Riche. Muſ. D. Boſc.

Statura praecedentium. Antennae articulis ſex anterioribus rufis, interne obtuſe prominulis, vltimis tribus paullo maioribus, medio roſtri inſertae. Caput rufum, nitidum, poſtice productum, cylindricum, longitudine corporis, nigrum. Thorax antice anguſtatus, ferrugineus, lateribus nigris. Elytra ſtriata: ſtriis punctatis, rufa, baſi apiceque parum flaueſcentia. Corpus ferrugineum, pedibus parum elongatis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*2guttatus.* 4. A. ater, nitidus, elytris puncto medio flauo.
Habitat in Sumatra. D. Daldorff.
Caput poſtice attenuatum et thorax laeuia, atra, nitida. Elytra laeuia, atra, nitida, puncto magno medio flauo. Pedes ferruginei, femoribus clauatis, claua atra.

*2puſtulatus.* 5. A. ater, elytris macula baſeos rufa. Ent. ſyſt. 2. 383. 3.
Habitat in America boreali. D. Yeats.

*rufus.* 6. A. rufus, immaculatus, elytris ſtriatis.
Habitat in Iaua. Muſ. D. Boſc.
Statura et magnitudo A. Penſyluanici, totus rufus. Caput poſtice anguſtatum. Elytra ſtriata: ſtriis punctatis.

*Penſyluanicus.* 7. A. niger, elytris rufis: faſcia media apicisque nigra. Ent. ſyſt. 2. 385 4. *
Linn. ſyſt. Nat. 2. 620. 5.
Habitat in America boreali.

*gemmatus* 8. A. ferrugineus: tuberculis nigris ſparſis. Ent. ſyſt. 2. 385. 5. *
*Attelabus gemmatus.* Thunb. nou. Inſ. ſp. 3. 68. fig. 80.
Habitat ad Cap. Bon. Spei.

*notatus.* 9. A. obſcure ferrugineus, elytris flauo-notatis, pedibus flauis. Ent. ſyſt. 2. 385. 6. *
Habitat in India. D. de Paykull.

*Tranquebaricus.* 10. A. rufus, elytris faſciis duabus nigris: anteriore interrupta. Ent. ſyſt. ſuppl. 161. 6. *
Habitat Tranquebariae. Muſ. D. de Seheſtedt.

*longimanus.* 11. A. piceus, elytris apice retuſis, pedibus anticis elongatis: femoribus vnidentatis. Ent. ſyſt. 2. 385. 7. *
Habitat in Senegallia. D. de Paykull.

12.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

2. A. fuſcus, elytris variolosis, baſi vniſpinoſis, pedibus anticis elongatis: femoribus vnidentatis. Ent. ſyſt. ſuppl 162. 7. * *dentipes.*

Habitat in Americae Inſulis. Muſ. D. Lund.

3. A. ferrugineus, elytris laeuiſſimis, atris. Ent. ſyſt. 2. 386. 8. * *nigripennis.*

Habitat ad Cap. Bon. Spei. Dom. de Paykull.

4. A. ferrugineus, antennis tarſisque fuſcis, elytris punctato-ſtriatis. Ent. ſyſt. 2. 386. 9. * *fuſcicornis*

Habitat in Senegallia. Dom. de Paykull.

5. A. ferrugineus, elytris ſpinoſis, atris. *hyſtrix.*

Habitat in Sumatra. D. Daldorff.

Magnitudo A. Coryli. Caput ferrugineum, antennis palpisque nigris. Thorax ferrugineus, tuberculis duobus dorſalibus, obtuſis. Scutellum ferrugineum, ſpina parua, nigra. Elytra ſtriata, atra, ſpinis quatuor erectis, validis, acutis. Corpus ferrugineum, femoribus annulo fuſco.

6. A. ater, thorace rufo, elytris baſi angulatis, pedibus anticis elongatis: femoribus vnidentatis. *ruficollis.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo A. curculionoidis. Corpus atrum, nitidum. Roſtrum apice incraſſatum. Thorax laeuis, rufus, nitidus, immaculatus. Elytra ſubſtriata, atra, nitida, angulo baſeos prominente. Pedes atri, antici longiores, femoribus clauatis, dente valido, recuruo, acuto.

7. A. ater, nitidus, femoribus omnibus ſubſerratis. *atratus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentis, paullo minor, totus ater, nitidus, immaculatus. Elytra ſtriata. Femora omnia denticulis tribus paruis, ſerrata.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*cyanipennis.* 18. A. ater, elytris ſtriatis, cyaneis.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura praecedentis, at paullo minor. Thorax punctatus. Elytra ſtriata, cyanea, immaculata. Abdomen ſupra rufum. Pedes cyanei, femoribus ſubſerratis.

*nitidus.* 19. A. ater, nitidus, elytris ſtriatis, femoribus muticis.
Habitat in America meridionali. D. Smidt. Mus. Dom. de Seheſtedt.
Affinis A. atrato, at minor. Caput et thorax atra nitida, immaculata. Elytra ſtriata, angulo baſeos parum prominulo. Pedes antici elongati, femoribus incraſſatis, at muticis.
Variat rarius thorace parum rufeſcente.

*ſpinoſus.* 20. A. ferrugineus, thorace elytrisque ſpinoſis. Ent. ſyſt. 2. 386. 10. *
Habitat ad Cap. Bon. Spei. Dom. de Paykull.

*indicus.* 21. A. ferrugineus, capite elytrorum baſi faſciaque media cyaneis. Ent. ſyſt. 2. 386. 11. *
Thunb. nou. Inſ. ſp. 3. 68. fig. 81.
Habitat in India orientali.

*curculionoides.* 22. A. niger, thorace elytrisque ſtriato-punctatis rubris. Ent. ſyſt. 2. 386. 12.
Geoff. Inſ. 1. 273. 10.
Payk. Monogr. 130. 122.
Fn. Sv. 3. 171. 2.
Schaeff. Icon. tab. 56. fig. 7.
Sulz. Inſ. tab. 4. fig. 12.
Habitat in Europae Corylo, Salice.

*varioloſus.* 23. A. ater, thorace elytrisque ferrugineis, elytra varioloſis.
Habitat in Barbaria. Muſ. Dom. Boſc.





Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et ſumma affinitas A. curculionoidis. Roſtrum paullo breues, atrum. Thorax laeuis, rufus, immaculatus. Elytra haud ſtriata, at punctis plurimis, latioribus, rugis, varioloſa. Corpus atrum, pedibus anticis parum elongatis.

24. A. ferrugineus, coleoptris angulatis: diſco nigro. Ent. ſyſt. 2. 387. 13. ♀ *angulatus.*

Habitat in Cayenna. Dom. v. Rohr.

25. A. violaceus, hirtus, roſtro atro. Ent. ſyſt. 2. 387. 14. ♂ *pubeſcens.*

Habitat in Germania. Dom. de Hattorff.

26. A. hirtus, ater, elytris crenato-ſtriatis, coerulescentibus. *hirtus.*

Habitat in Carolina. Muſ. Dom. Boſc.

Statura et magnitudo omnino A. pubeſcentis. Roſtrum elongatum, atrum, antennis articulis tribus vltimis incraſſatis. Thorax hirtus, ater. elytra valde crenato-ſtriata, coeruleſcentia. Corpus nigrum, pedibus inermibus.

27. A. aureus, roſtro plantisque nigris. Ent. ſyſt. 2. 387. 15. *Bacchus.*

*Curculio Bacchus.* Linn. ſyſt. Nat. 2. 611. 38.
Payk. Monogr. 133. 124.
Fn. Sv. 3. 172. 4.
Geoff. Inſ. 1. 270. 4.
Degeer Inſ. 5. 250. 38.
Schaeff. Icon. tab. 39. fig. 18.
Sulz. Inſ. tab. 4. fig. 4.
Habitat in Vite, Corylo.
Variat ſexu: thorace antrorſum ſpinoſo.

28. A. corpore viridi-aurato, ſubtus concolore. Ent. ſyſt. 2. 387. 16. *Betuleti.*

*Curculio Betulae.* Linn. ſyſt. Nat. 2. 611. 39.
Fn. Sv. 605.
Geoff. Inſ. 1. 270. 2.

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Monogr. 131. 123.
Fn. Sv. 3. 170. 3. *a*.
Degeer Inſ. 5. 248. 5. tab. 7. fig. 25.
Habitat in Betula alba, in Vite, folia contorquens.
Variat ſexu: thorace antrorſum ſpinoſo.

*Populi.* 29. A. corpore viridi-ignito, ſubtus atro, coeruleſcente. Ent. ſyſt. 2. 387. 17.
*Curculio Populi.* Linn. ſyſt. Nat. 2. 611. 40.
Fn. Sv. 606.
Payk. Monogr. 132. 123. β.
Fn. Sv. 3. 171. 3. β.
Ent. Helvet. 1. 110. 2. tab. 13. fig. 3. 4.
Geoff. Inſ. 1. 270. 3.
Degeer Inſ. 5. 249. 38.
Habitat in Betula, Populo.
Varietatem praecedentis credit Dom. de Paykull, at vix merito.

*bicolor.* 30. A. niger, capite thorace elytrisque rubris, roſtro longitudine corporis, atro. Ent. ſyſt. 2. 388. 18. *
Habitat in America. Muſ. Dr. Hunter.

*hungaricus* 31. A. villoſus, ater, thorace elytrisque rufis: ſuturа atra. Ent. ſyſt. 2. 388 19. *
*Curculio hungaricus.* Herbſt. Arch. t. 24. f. 10.
Habitat in Hungaria.

*aequatus.* 32. A. cupreo-niger, elytris rufis. Ent. ſyſt. 2. 388. 20. *
*Curculio aequatus.* Linn. ſyſt. Nat. 2. 607. 9.
Payk. Monogr. 135. 126.
Fn. Sv. 3. 173. 6.
Panz. Fn. Germ. 20. tab. 8.
Geoff. Inſ. 1. 270. 4.
Habitat in Europae Salice.
Variat duplo minor, ſcutello ſuturaque nigris, an diſtinctus?

33.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

3 A. ater, elytrorum ſtriis ſimpliciſſimis, femoribus ruſis. *femoralis.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Maior A. aequato. Roſtrum elongatum, incuruum, atrum. Thorax antice attenuatus, laeuis, ater, immaculatus. Elytra ſtriis laeuiſſimis, atra. Pedes atri, femoribus ruſis.

4. A. violaceus, nitens, thorace elytrisque teſtaceis. Ent. ſyſt. 2. 389. 21. * *coeruleocephalus.*

Herbſt. Arch. tab. 24. fig. 11.

Act. Hallenſ. 1. 282.

Habitat in Saxoniae Crataego. Dom. Hybner.

5. A. obſcure-aeneus, ſubtus obſcurior. Ent. ſyſt. 2. 389. 22. *cupreus.*

*Curculio cupreus.* Linn. ſyſt. Nat. 2. 608. 21.

Fn. Sv. 593.

Payk. Monogr. 134. 125.

Fn. Sv. 3. 173. 5.

Habitat in Europae borealis hortis.

6. A. coeruleus, immaculatus, antennis tibiisque fuſcis. Ent. ſyſt. ſuppl. 162. 22. * *coeruleus.*

Habitat in Germania. D. Daldorff.

7. A. niger, elytris aeneis. Ent. ſyſt. 2. 389. 23. * *aeneus.*

*Curculio aeneus.* Payk. Monogr. 142. 134.

*Attelabus aeneus.* - Fn. Sv. 3. 179. 14.

Habitat in Hyſſopo officinali Europae borealis.

8. A. niger, obſcurus, elytris ſtriatis, cyaneis. Ent. ſyſt. ſuppl. 163. 23. * *triſtis.*

Habitat in Germania. Dom. Schneider.

9. A. obſcurus, elytris ſtriatis, cinereis: faſcia lata, media atra. Ent. ſyſt. ſuppl. 163. 23. * *ſellatus.*

Habitat in Boruſſia. Dom. Schneider.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fuſciroſtris* 40. A. griſeus, pedibus rufis, roſtro femoribusque baſi nigris. Ent. ſyſt. 2. 389. 24. *

Habitat in Germania. Dom. de Hattorff.

*cupriroſtris.* 41. A. viridi-aeneus, elytris ſtriatis, roſtro cupreo Ent. ſyſt. 2. 389. 25. *

*Curculio viridi ſericeus.* Geoff. Inſ. 1.

Habitat in Saxoniae Betula. Dom. Zſchuck.

*4puſtulatus* 42. A. ater, thorace ſcabro: puncto antico, elytris ſtriatis: maculis duabus rufis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paruus. Caput atrum. Thorax rotundatus, ſcaber, ater, puncto in medio marginis antici obſcure rufo. Elytra ſtriata, atra, maculis duabus obſcure rufis.

*rufiroſtris.* 43. A. niger, roſtro dimidiato pedibusque rufis. Ent. ſyſt. 2. 390. 26. *

Habitat in Anglia. Muſ. D. Banks.

*nigriroſtris.* 44. A. cinereus, roſtro arcuato, atro, pedibus teſtaceis. Ent. ſyſt. ſuppl. 163. 26. *

Habitat in floribus Onobrychis prope Tanger. D. Schousboe. Muſ. D. Lund.

*atriroſtris* 45. A. niger, obſcurus, roſtro atro nitido, pedibus teſtaceis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Minutus. Roſtrum elongatum, arcuatum, atrum nitidum. Thorax, elytra, corpus nigra, obſcura. Pedes rufi.

*Crotalariae.* 46. A. ater, capitis vertice impreſſo, thorace vtrinque retuſo.

Habitat in Crotalariae leguminibus Americae Muſ. D. Lund.

Praecedentibus maior, totus ater, parum nitidus. Capitis vertex parum impreſſus. Roſtrum elongatum,

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

gatum, cylindricum, incuruum. Thorax margine exteriore, antice retuſus, laeuis, ater. Elytra ſtriata. Pedes inermes, atri.

47. A. pubeſcens, cyaneus, elytris ſtriatis. Ent. ſyſt, 2. 390. 27. *Alliariae.*

*Curculio Alliariae.* Linn. ſyſt. Nat. 2. 506. 4. Fn. Sv. 580.
Friſh. Inſ. 9. 35. tab. 78.
Degeer Inſ. 5. 251. 39.
Payk. Fn. Sv. 3. 175. 8.
Habitat in Eryſimo Alliaria.

48. A. niger, obſcurus, elytris ſtriatis, obſcure coeruleis, roſtro baſi compreſſo, apice ſubulato. Ent. ſyſt. ſuppl. 164. 27. * *Pomonae.*

*Curculio ruficornis.* Herbſt. Col. tab. 112. f. 8. at antennae in noſtris vix ruſae.
Habitat in Kiliae Pomona.

49. A. ater, nitidus, elytris ſtriatis, roſtro elongato apice depreſſo. *planiroſtris.*

Habitat in Germania. D. Smidt.
Paullo maior A. Craccae. Antennae moniliformes, extrorſum craſſiores. Roſtrum elongatum, apice dilatatum, depreſſum. Elytra ſtriata. Corpus atrum.

50. A. ater, roſtro porrecto cylindrico, elytris ſtriatis globoſis. *Piſi.*

Habitat in Auſtria. Dom. de Meyerle.
Paullo maior A. Craccae, totus ater. Roſtrum porrectum, arcuatum, cylindricum. Elytra ſtriata, globoſa, atra. Pedes inermes.

51. A. niger, obſcurus, elytris ſtriatis, obſcure coeruleis, roſtro apice depreſſo. Ent. ſyſt. 2. 390. 28. * *Craccae.*

*Curculio Craccae.* Linn. ſyſt. Nat. 2. 506. 6. Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Monogr. 140. 131.
- Fn. Sv. 3. 177. 11.
Degeer Inf. 5. 253. 83. tab. 6. fig. 31. 32.
Habitat in Holfatiae hortis.

*Sorbi.* 52. A. ater, elytris ftriatis, obfcure coeruleſcentibus, roſtro arcuato, longitudine corporis. Ent. fyſt. 2. 390. 29. *
*Curculio laeuigatus.* Payk. Monogr. 141. 138.
- Fn. Sv. 3. 178. 12.
Habitat in Germaniae arboribus.

*cyaneus.* 53. A. ater, elytris cyaneis nitidulis. Ent. fyſt. 2. 391. 30. *
*Curculio cyaneus.* Linn. fyſt. Nat. 2. 506. 5. Fn. Sv. 581.
Herbſt. Arch. tab. 24. fig. 5.
Habitat in Europae Malua.

*globoſus.* 54. A. gibbus, fupra ater, fubtus cinereus. Ent. fyſt. 2. 391. 31. *
*Curculio Hypoleucos.* Payk. Monogr. 75. 72.
Quenz. Diff. Inf. 16.
Panz. Fn. Germ. 57. tab. 10.
Habitat in Europae arena.

*ouatus.* 55. A. ouatus, gibbus, cyaneus; nitidus, elytris abbreuiatis ftriatis.
Habitat in Carolina. Muf. D. Boſc.
Statura parua A. globoſi. Caput cyaneum, roſtro breui, atro, antennarum articulis tribus vltimis incraſſatis. Thorax planus, punctatus, cyaneus. Elytra abdomine breuiora, crenato-ftriata, cyanea, nitida. Corpus cyaneum.

*Maluae.* 56. A. griſeus, elytris pedibusque teſtaceis. Ent. fyſt. 2. 391. 32. *
*Curculio Pineti* Payk. Monogr. 61. 58.
Habitat in Angliae Maluis.

57.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

57. A. ater, femoribus luteis. Ent. ſyſt. 2. 391. 34. *flauipes.*
*Curculio flauipes.* Payk. Monogr. 143. 135.
Geoff. Inſ. 1. 272. 8. *II. 51.*
Habitat primo vere frequens in locis apricis, calidis.

58. A. ater, antennis pedibusque flauis. *flauicornis*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura parua, A. flauipedis. Roſtrum elongatum, cylindricum, incuruum. Thorax antice attenuatus, niger. Elytra ſtriata. Antennae et pedes omnes flaua.

59. A. ſanguineus, elytris crenato-ſtriatis. Ent. ſyſt. 2. 392. 35. *frumentarius.*
*Attelabus purpureus.* Ent. ſyſt. 2. 391. 34.
*Curculio purpureus.* Linn. ſyſt. Nat. 2. 607. 14. Fn. Sv. 585.
*Curculio frumentarius.* Linn. ſyſt, Nat. 2. 608. 15. Fn. Sv. 586.
*Rhinomacer frumentarius.* Ent. Helv. 1. 108. 1. tab. 13. fig. 1. 2.
Payk. Monogr. 139. 130. *II. 52.*
- Fn. Sv. 3. 177. 10.
Act. Holm. 1750. 186. 1.
Leuwenh. Arc. 183. Aug. 6. 83. fig. 3.
Degeer Inſ. 5. 251. 48.
Habitat in frumento diutius aſſeruato.

60. A. nigricans, elytris cinereis: faſciis duabus nigris, pedibus rufis. Ent. ſyſt. 2. 392. 36. * *vernalis.*
*Curculio vernalis.* Payk. Monogr. 138. 129.
- Fn. Sv. 3. 183. 18.
Habitat in Holſatiae Vrticae, primo vere.

61. A. pedibus ſaltatoriis, corpore atro. Ent. ſyſt. 2. 392. 37. Linn. ſyſt. Nat. 2. 620. 7. Fn. Sv. 640. *Betulae.* *II. 52*

*Cur-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Betulae.* Payk. Monogr. 127. 120.
Degeer Inſ. 5. 209. 47.
Habitat in Betulae foliis frequens, quae rodendo pulcre criſpa reddit.

157. RHINOMACER. *Palpi* extrorſum craſſiores: articulo vltimo truncato.
*Antennae* ſetaceae, roſtro inſidentes.

*varius.* 1. R. nigro alboque varius. Ent. ſyſt. ſuppl. 164. 1.
Habitat ad Cap. Bon. Spei. D. Weber.

*curculioides.* 2. R. villoſo-griſeus, antennis pedibusque nigris. Ent. ſyſt. 2. 393. 1. *
*Mycterus griſeus.* Ent. Helv. 1. 124. 1. tab. 16. fig. 1. 2. 3.
I. 48 Panz. Fn Germ. 12. tab. 8.
*Curculio Rhinomacer.* Payk. Monogr. 126. 119.
*Anthribus Rhinomacer.* Payk. Fn. Sv. 3. 166. 8.
Habitat in Italia. Dr. Allioni.

*attelaboides.* 3. R. villoſo-piceus, antennis pedibusque teſtaceis. Ent. ſyſt. 2. 393. 2. *
*Curculio Rhinomacer.* Payk. Monogr. 126. 119. β.
*Anthribus Rhinomacer.* Payk. Fn. Sv. 3. 166. 8. β.
Habitat in Sueciae plantis. D. de Paykull.

*Va-*

---

*Rhinomacri* corpus parvum, ouatum, villoſum, immarginatum, tardum; *capite* inſerto, ouato, antice roſtro breui, planiuſculo: *oculis* globoſis, marginalibus; *antennis* longitudine thoracis in medio roſtri inſertis; *thorace* ouato; *ſcutello* rotundato: *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, tenuibus; *tarſis* quadriarticulatis: *colore* obſcuro, minime nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Varietas* praecedentis teſte D. de Paykull.

4. R. villoſus, ater, elytris fuſcis. *lepturoides*
Habitat in Auſtria. D. de Meyerle.
Statura omnino praecedentium, at totus villoſus, ater, elytris ſolis fuſcis aut minus nigris. IX.

## 158. CALANDRA.

*Mandibula* lata, fornicata, truncata.
*Antennae* claua truncata: articulo vltimo ſpongioſo, retractili.

1. C. thorace elytrisque ſcabris, antennis apice albis. *gigas.*
*Curculio gigas.* Ent. ſyſt. 2. 394. 1. ♀
Oliv. Inſ. 83. tab. 12. fig. 146.
Habitat in Iaponia. Muſ. Britann.

2. C. obſcura, roſtro apice ſpina eleuata, compreſſa. *ferriroſtris*
Oliv. Inſ. 83. tab. 17. fig. 211.
Drury Inſ. 2. tab. 33. fig. 3?
Habitat in Iaua. Muſ. D. de Seheſtedt.
Maior C. palmarum. Roſtrum porrectum, rectum, dorſo apice ſerratum, dente eleuato, compreſſo. Antennae claua truncata, nigra. Thorax laeuis, obſcure rufus, lineis duabus magnis dorſalibus abbreuiatis, atris. Elytra ſubſtriata, obſcura, immaculata. Corpus atrum, femoribus tibiisque omnibus interne pilis aureis ciliatis.

3.

---

*Calandrae* corpus ouatum, planum, glabrum, immarginatum, tardum; *capite* paruo, gibbo, inſerto, antice producto in roſtrum ſubcompreſſum; *oculis* magnis, oblongis, lateralibus; *antennis* longitudine thoracis, in forma oblonga ante roſtri baſin inſertis; *thorace* plano, antice annulari caput arcte cingente, poſtice rotundato; *elytris* rigidis, planis, fornicatis, abdomine breuioribus; *pedibus* breuibus, compreſſis, validis; *tarſis* quadriarticulatis; *colore* vario, obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Palmarum.* 3. C. atra, elytris abbreuiatis ſtriatis.
*Curculio Palmarum.* Ent. ſyſt. 2. 395. 2.
Linn. ſyſt. Nat. 2. 506. 1. Muſ. Lud. Vlr. 42.
Oliv. Inſ. 2. 83. tab. 2. fig. 16.
Degeer Inſ. 5. tab. 15. fig. 26.
Rumph. Amb. 1. 79. tab. 17. fig. 9.
Petiv. Gazoph. tab. 35. fig. 5.
Gronov. Zooph. 578. tab. 16. fig. 4.
Mer. Surin. tab. 48. fig. 3.
Sulz. Hiſt. Inſ. tab. 3. fig. 20.
Habitat in Indiae Palmis.

*Phoenicis.* 4. C. atra, roſtro antice, elytrorum margine pedibusque anticis rufo-ciliatis.
Habitat in Guinea. D. Meier. Muſ. D. Lund.
Statura et ſumma affinitas C. Palmarum. Roſtrum ſubcompreſſum, antice criſta parua e pilis erectis ferrugineis. Thorax planus, laeuis, poſtice rotundatus, at minus dilatatus, ater, lineis duabus obſoletis, rubris. Elytra ſtriata, abdomine breuiora, margine exteriori rufo ciliata. Corpus atrum, pedibus quatuor anticis denſe rufo ciliatis.

*Boraſſi.* 5. C. atra, nitida, roſtro apice vtrinque reflexo, pedibus ciliatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo omnino C. longipedis. Tota totus ater, immaculatus, nitidus. Roſtrum ſubcompreſſum, apice vtrinque reflexum. Thorax planus, laeuis, ater, immaculatus, nitidus, poſtice rotundatus. Elytra abdomine breuiora, ſtriata. Pedes atri, ſubtus pilis nigris ciliati, tibiis apice ſpina acuta.

*cruentata.* 6. C. atra, thorace lineis, elytris punctis duobus ferrugineis.

*Cur-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio cruentatus.* Ent. ſyſt. 2. 395. 3. *
Oliv. Inſ. 33. tab. 12. fig. 147.
Habitat in Carolina. Muſ. D. Banks.

6. C. nigricans, elytris ferrugineis, roſtro apice vtrinque reflexo, pedibus anticis elongatis. *longipes.*
*Curculio longipes.* Ent. ſyſt. 2. 395. 4. *
Oliv. Inſ. 83. tab. 15. fig. 191.
Voet. Col. 2. tab. 86. fig. 10.
Habitat in China. Muſ. D. Lee.
*Variat* ſexu, roſtri dorſo ſerrato.

8. C. atra, thorace elytrisque fuſcis, roſtro apice vtrinque reflexo. *heros.*
Habitat in Sumatra. D. Daldorff.
Maximus in hoc genere et affinis C. Coloſſo. Caput laeue, atrum, roſtro compreſſo, apice vtrinque reflexo. Thorax planus, obſcurus. Elytra vix ſtriata, fuſca. Corpus atrum.

9. C. nigricans, elytris brunneis aut cinereis, pedibus elongatis. *Coloſſus.*
*Curculio Coloſſus.* Ent. ſyſt. 2. 396. 5. *
Oliv. Inſ. 83. tab. 3. fig. 32.
Voet. Col. 2. tab. 36. fig. 9.
Habitat in India orientali.

10. C. atra, thorace ſubouato, excauato, punctato, elytris rugoſo-ſulcatis, tibiis ſpinoſis. *Indus.*
*Curculio Indus.* Ent. ſyſt. 2. 396. 6.
Linn. ſyſt. Nat. 2. 506. 2. Muſ. Lud. Vlr. 43.
Degeer Inſ. 5. 45. 22.
Habitat in India.

11. C. ſupra cinerea, nigro-irrorata, elytris maculis duabus nigris. *4punctata.*
Habitat in Sumatra. D. Daldorff.
Statura omnino C. hemipterae, at paullo maior. Antennae atrae, claua ouata, compreſſa, apice alba. Roſtrum breue, atrum, dorſo baſeos cinereum.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nereum. Thorax planus, cinereus, atro varius. Elytra abbreuiata, ſtriata, cinerea, nigro parum irrorata, maculisque duabus diſtinctis nigris. Anus prominet cinereus, ſtriga nigra pone apicem elytrorum. Corpus cinereum, medio atrum. Pedes nigri.

*Guineenſis* 12. C. griſea, roſtro bidentato, elytris ſubſpinoſis.
*Curculio Guineenſis*. Ent. ſyſt. ſuppl. 164. 6.
Habitat in Guinea. Muſ. D. Lund.

*diſcors.* 13. C. punctata, cinerea, albo-lineata.
Habitat in Sumatra. D. Daldorff.
Triplo praecedenti minor. Antennae longitudine roſtri, cinereae: articulis aequalibus breuibus, cylindricis; vltimo maiori, ouato, fuſco, apice albo ad baſin roſtri inſertae. Roſtrum baſi craſſius, cinereum, apice arcuatum, atrum. Thorax varioloſus, cinereus, lineis tribus niueis. Scutellum niueum Elytra connata, varioloſa, cinerea: vitta ſubmarginali, ſtrigam mediam obliquam, interne flexuoſam exſerente. Corpus cinereum, pedibus anticis longioribus.

*granulata.* 14. C. obſcura, thorace noduloſo, elytris punctato-ſtriatis.
Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.
Statura et magnitudo omnino C. guineenſis. Antennae claua ouata, compreſſa, apice pallida. Thorax fuſcus, nodulis eleuatis, ſubacutis, inaequalis. Elytra obſcura, punctato-ſtriata. Pedes ſcabri, cinerei, antici parum elongati.

*pagana.* 15. C. griſea, thoracis dorſo fuſco: arcubus cinereis, roſtro biſulcato.
*Curculio paganus*. Ent. ſyſt. 2. 396. 7. *
Habitat in Indiis. Dom. Torſtenſon.

16.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

16. C. obſcure-purpuraſcens, elytris abbreuiatis, femoribus mediis ciliatis. *ferruginea*

*Curculio ferrugineus* Ent. ſyſt. 2. 396. 8.
Oliv. Inſ. 83. tab 2. fig. 16.
Sulz. Hiſt. Inſ. tab. 4. fig. 5.
Habitat in Braſilia.
In altero ſexu roſtri dorſum carinatum.
Inſtrumenta cibaria in hisce Curculionum generibus minutiſſima difficillime eruuntur, nec mihi adhuc ſatis faciunt.
Os porrectum, in roſtrum elongatum, incuruum, corneum, cylindricum. Palpos haud obſeruaui. Mandibulae latae, fornicatae, truncatae, dentatae. Maxilla breuiſſima, cornea, vti videtur, integra. Labium transuerſum, corneum, breue, integrum. Antennae articulo ſecundo longiſſimo, reliquis aequalibus, ſenſim craſſioribus, vltimo craſſiori, compreſſo, apice ſpongioſo, retractili, baſi roſtri inſertae.

17. C. atra, thoracis dorſo antennarumque claua purpureis, femoribus omnibus ciliatis. *Schach.*

Oliv. Inſ. 83. tab. 16. fig. 204.
Habitat in Sumatra. D. Daldorff.
Statura magna C. ferrugineae, cuius forte mera varietas. Roſtrum cylindricum, atrum, ante apicem pilis rufis ciliatum. Antennae atrae, claua purpurea, apice truncata, cineraſcente. Thorax laeuis, ater, obſcurus, dorſo purpuraſcente. Elytra abbreuiata, ſtriata, atra. Corpus atrum, abdomine macula media baſeos ſanguinea. Pedes atri, femoribus tibiisque intus pilis ferrugineis, ciliatis.

18. C. obſcure-purpuraſcens, elytris abbreuiatis, maculatis. *hemiptera.*

*Curculio hemipterus.* Ent. ſyſt. 2. 396. 9. *



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. ſyſt. Nat. 2. 506. 3. Muſ. Lud. Ulr. 44.
Oliv. Inſ 83. tab. 1. fig. 1.
Degeer Inſ. 5. tab. 15. fig. 25.
Habitat in Cayenna. Muſ D. Olivier.

*cribraria.* 19. C. nigra, thorace elytrisque purpureis, nigro-punctatis.
*Curculio cribarius.* Ent. ſyſt. ſuppl. 165. 9.*
Habitat in America boreali. Dom. Mauerhoff.

*variegata.* 20. C. rufo nigroque varia, roſtro apice nigro.
*Curculio variegatus.* Ent. ſyſt. 2. 397. 10.*
Oliv. Inſ. 83. tab. 13. fig. 158.
Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

*melanocardia.* 21. C. cinerea, coleoptris macula commune cordata fuſca.
*Curculio melanocardius.* Ent. ſyſt. 2. 397. 11.*
Linn. ſyſt. Nat. 2. 608. 18. Muſ. Lud. Vlr. 45.
Sulz. Hiſt. Inſ. tab. 4. fig. 11.
Habitat in India.

*limbata.* 22. C. nigricans, thoracis elytrorumque margine rufo.
*Curculio limbatus.* Ent. ſyſt. 2. 397. 12.*
Oliv. Inſ. 83. tab. 2. fig. 22.
Habitat in Senegallia.

*cruciata.* 23. C. nigra, thorace ſubſpinoſo: lineis elytris poſtice cruce albis.
*Curculio cruciatus.* Ent. ſyſt. 2. 397. 13.*
Oliv. Inſ. 83. tab. 11. fig. 131.
Habitat in noua Hollandia. Muſ. D. Banks.

*ſanguinolenta.* 24. C. atra, elytris faſcia, baſeos ſanguinea.
*Curculio ſanguinolentus.* Ent. ſyſt. 2. 398. 14.*
Oliv. Inſ. 83. tab. 10. fig. 118.
Habitat in Inſula Tabago Americae. Dom. Lee.

*faſciata.* 25. C. atra, elytris ſtriatis: faſcia media ſanguinea.

*Cu-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio faſciatus.* Ent. ſyſt. 2. 398. 15. *
Oliv. Inſ. 83. tab. 11. fig. 136.
Habitat - - - Muſ. Dom. Francillon.

6. C. laeuis, atra, nitida, elytris maculis duabus rufis. *4puſtulata.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo praecedentium. Roſtrum elongatum, arcuatum, atrum. Thorax laeuiſſimus, ater, nitidus, immaculatus. Elytra laeuiſſima, apice parum ſtriata, atra, maculis duabus rufis. Corpus atrum.

7. C. plana, atra, nitida, elytris ſtriatis, femoribus rufis. *planicollis.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo praecedentis Roſtrum elongatum, arcuatum, atrum. Thorax planus, margine vtrinque rotundato. Elytra ſtriata, apice decliuia. Pedes nigri, femoribus rubris.

8. C. glabra, nitida, roſtro arcuato. *coruina.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Tota atra, nitidiſſima, immaculata.

9. C. nigra, obſcura, thorace ſubpunctato, elytris ſtriato-punctatis. *necydaloides.*
Habitat in Carolina. Muſ. D. Boſc.
Statura omnino C. abbreuiatae, at diſtincta et alia. Caput atrum, nitidum, roſtro incuruo. Thorax niger, obſcurus, vix punctatus. Elytra abdomine breuiora, ſtriato-punctata.

10. C. cinerea, capite atro. *melanocephala.*
Habitat in Carolina. Muſ. Dom. Boſc.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paullo minor C. abbreuiata. Caput atrum. Thorax et elytra laeuia, cinerea, fere immaculata. Corpus et pedes atra.

*abbreuiata* 31. C. atra, thorace punctato, elytris ſubſtriatis.
*Curculio abbreuiatus.* Ent. ſyſt. 2. 403. 37.
*Calandra abbreuiata.* Ent. Helv. 1. 64. 2. tab. 2. fig. 3. 4.
Panz Fn. Germ. 42. tab. 3.
Habitat Halae Saxonum. Dom. Hybner.

*lituſata.* 32. C. nigra, obſcura, elytris ſubtuberculatis: litura baſeos rufa.
Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.
Media. Roſtrum arcuatum, atrum. Thorax planus, pubeſcens, niger, obſcurus. Elytra tuberculis plurimis, eleuatis, obſoletis, inaequalibus nigra, obſcura, litura ſinuata, baſeos rufa.

*ſignum.* 33. C. atra, nitida, elytris ſtriatis: lineola baſeos abdomine maculis albis.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Statura omnino C. nitentis. Roſtrum arcuatum, atrum. Caput et thorax antice attenuatus, atra, nitida, immaculata. Elytra ſtriata, lineola parua baſeos ad ſuturam alba. Corpus atrum, maculis aliquot albis ad baſin abdominis.

*porcula.* 34. C. atra, thorace elytrisque obſcuris, griſeo-punctatis.
*Curculio porculus.* Ent. ſyſt. ſuppl. 166. 36.
Habitat Halae Saxonum.

*nitens.* 35. C. atra, nitida, thorace plano, laeui, elytris ſtriatis, longitudine abdominis.
*Curculio nitens.* Ent. ſyſt. 2. 403. 38.
Habitat in Gallia meridionali.

D.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Diſtincta omnino a C. coruina, duplo minor.

6. C. nigra, thorace varioloſo, elytris ſtriatis, albo-irroratis. *viduata.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Magnitudo praecedentis. Roſtrum longum, arcuatum, laeue. Thorax niger, obſcurus, varioloſus lineaque dorſali eleuata. Elytra crenato-ſtriata, nigra, atomis obſoletis, albidis, irrorata. Pedes nigri, obſcuri.

7. C. atra, nitida, elytris ſtriatis: lineola poſtica alba. *carbonaria*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Roſtrum longum, arcuatum. Thorax laeuiſſimus, ater, nitidus. Elytra abdomen omnino tegentia, ſtriata, atra, ante apicem lineola parua, alba. Corpus atrum.

8. C. plana, atra, thorace punctato, elytris ſtriatis. *Sabæ.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Praecedentibus affinis, at minor. Roſtrum longum, incuruum. Thorax planus, punctatus, ater. Elytra ſtriata.

9. C. picea, thorace punctato, longitudine elytrorum. *granaria.*

*Curculio granarius.* Ent. ſyſt. 2. 414. 88.

*Calandra granaria.* Ent. Helv. 1. 62. 1. tab. 2. fig. 1. 2.

Linn. ſyſt. Nat. 2. 608. 16. Fn. Sv. 587.

Geoff. Inſ. 1. 255. 18.

Payk. Monogr. 54. 51.

Panz. Fn. Germ. 17. tab. 11.

Joblot. Obſ. Microſc. 1754. tab. 7. fig. 1.

Degeer

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inf. 5. 239. 25.
Habitat in Europae granariis, frumentorum grana exedens. Pellitur Iſatide, Hyoſcyamo, Sambuco, Thlaſpide.

*Frumenti.* 40. C. nigra, thorace punctato, ferrugineo, nigro maculato, elytris ſtriatis: baſi maculaque ferrugineis.
Habitat in Sumatra. Dom. Daldorff.
Statura omnino C. Oryzae, at paullo maior. Antennae nigrae, claua ouata, apice niuea. Thorax elongatus, planus, punctatus, ferrugineus macula media lunulaque magna baſeos nigris. Elytra abbreuiata, ſtriata, nigra, macula magna baſeos minorique media, ferrugineis. Corpus nigrum, abdominis baſi pedibusque ferrugineis.

*Oryzae.* 41. C. picea, thorace punctato, longitudine elytrorum, elytris punctis duobus rufis.
*Curculio Oryzae.* Ent. ſyſt. 2. 414. 89.
Linn. Amoen. acad. 6. 395. 19.
Oliv. Inſ. 83. tab. 8. fig. 81.
Habitat in Oryza ſaepius lecta inter grana ex Oriente adportata.

*2 tuberculata.* 42. C. ferruginea, thorace longitudine elytrorum punctisque duobus dorſalibus, eleuatis.
*Curculio 2 tuberculatus.* Ent. ſyſt. 2. 414. 90.
Oliv. Inſ. 83. tab. 9. fig. 165.
Habitat in noua Zelandia. Muſ. D. Banks.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

59. RHYNCHAENVS. *Palpi* filiformes.
*Maxilla* cylindrica, vnidentata.
*Antennae* clauatae: articulo primo longiſſimo, inſidentes roſtro elongato ante apicem.

* *femoribus ſimplicibus.*

. R. niger, elytris holoſericeis, ſtriatis. ***ſtriatus.***
*Curculio ſtriatus.* Ent. ſyſt. 2. 398. 16. *
*Curculio ſtriatulus* Oliv. Inſ. 83. 140.
Habitat in Inſula Terre neuve. Muſ. D. Banks.

:. B. ater, antennis apice griſeis, femoribus ſulcatis. ***rubetra.***
*Curculio rubetra.* Ent. ſyſt. 2. 398. 17. *
*Curculio gagates.* Oliv. Inſ. 83. fig. 104.
Habitat in Cayenna. Dom. v. Rohr.

. R. gibbus, ater, glaber, nitidus, roſtro ſubcompreſſo, tibiis anticis vnidentatis. ***gagates.***
*Curculio gagates.* Ent. ſyſt. 2. 398. 18. *
Habitat in Cayenna. Dom. de Paykull.

4.

---

***Rhynchaeni*** corpus oblongum, glabrum, immarginatum, tardum; ***capite*** paruo, rotundato, inſerto, antice producto in roſtrum elongatum, cylindricum, vtrinque ſulcatum ad receptionem ſecundi articuli antennarum; ***oculis*** magnis, oblongis, latentibus, ad baſin roſtri; ***antennis*** longitudine thoracis: articulo ſecundo longiſſimo, reliquis ſenſim craſſioribus, vltimis tribus clauatis, claua acutiuſcula, ante apicem roſtri inſertis; ***thorace*** breui, rotundato; ***ſcutello*** minuto, ſaepe nullo; ***elytris*** rigidis, fornicatis, inflexis, longitudine abdominis; ***pedibus*** breuibus, validis, ſaepe dentatis; ***tarſis*** quadriarticulatis; ***colore*** vario.

**Characterem generis in generibus Inſectorum dedi.**

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*aeneus.* 4. R. viridi-aeneus, thorace punctato, elytris ſtriatis
Habitat in America meridionali. D. Smidt. Muſ
Dom Lund.
Statura et magnitudo C. gagatis. Corpus totum
viridi-aeneum. Capitis vertex parum impreſ
ſus. Thorax punctatus. Elytra ſtriata.

*mendicus.* 5. R. ouatus, cineraſcens, elytris ſtriatis.
*Curculio mendicus.* Ent. ſyſt. 2. 399. 19.
Oliv. Inſ. 83. tab. 9. fig. 108.
Habitat in Inſula Madagaſcar. Muſ. Gall.

*Pineti.* 6. R. niger, elytris ſtriatis, albo-maculatis.
*Curculio Pineti.* Ent. ſyſt. 2. 399. 20.
*Curculio confuſus.* Payk. Monogr. 35. 33.
Fn. Sv. 3. 223. 41.
Herbſt. Col. 6. 293. 264. tab. 82. fig. 10.
Habitat in Europae Pino ſylueſtri frequens.

*Pini.* 7. R. elytris rufeſcentibus, faſciis nebuloſis.
*Curculio Pini.* Ent. ſyſt. 2. 399. 21.
Linn. ſyſt. Nat. 2. 608. 19. Fn. Sv. 589.
Payk. Monogr. 36. 34.
Fn. Sv. 3. 225. 43.
Oliv. Inſ. 83. tab. 4. fig. 42.
Panz. Fn. Germ. 42. tab. 1.
Degeer Inſ. 5. 222. 15.
Friſh. Inſ. 11. tab. 23. fig. 5.
Schaeff. Icon. tab. 25. fig. 7.
Habitat in Sueciae Pino.

*Onopordinis.* 8. R. niger, cinereo-villoſus, roſtro atro: vtrinque
ſulco abbreuiato baſeos.
*Curculio Onopordon.* Ent. ſyſt. 2. 399. 22.
Habitat in Africae Onopordine. D. Prof. Vahl.

*Vrſus.* 9. R. ferrugineo-fuſcus, albo-lineatus, roſtro nigro
*Curculio Vrſus.* Ent. ſyſt. 2. 399. 23.
Habitat in Italia. Dr. Allioni.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

10. R. niger, cinereo-irroratus, elytris puncto baſeos diſtincto. *laceae.*

*Curculio laceae.* Ent. ſyſt. 2. 400. 24. *
Geoff. Inſ. 1. 281. 8.
Panz. Fn. Germ. 18. tab. 2.
Habitat in Centaurea lacea. Prof. Hermann.

11. R. niger, immaculatus, roſtro cylindrico, elytris ſtriatis. *planus.*

*Curculio planus* Ent. ſyſt. 2. 400. 25. *
Habitat in Germania. D. Smidt.

12. R. flauo fuſcoque varius, abdomine cinereo, nigro-punctato. *punctulatus.*

*Curculio punctatus.* Ent. ſyſt. 2. 400. 26. *
Habitat in America. Muſ. D. Romani.

13. R. nigro-vireſcens, roſtro ferrugineo: orbita oculorum niuea. *ocularis.*

*Curculio ocularis.* Ent. ſyſt. 2. 400. 27. *
Habitat in Barbaria. Muſ. Dom. Desfontaines.

14. R. niger, vireſcenti-irroratus, roſtro nigro, ſubcarinato. *Cynarae.*

*Curculio Cynarae.* Ent. ſyſt. 2. 401. 28. *
*Curculio Cardui.* Roſſ. Fn. Etr. 1. 111. 284. tab. 5. fig. 11.
Panz. Naturf. 24. 18. 25. tab. 1. fig. 25.
Habitat in Cynarae floribus Africae. D. Vahl.

15. R. griſeus, elytris puncto albo. *Colon.*

*Curculio Colon.* Ent. ſyſt. 2. 401. 29.
Linn. Mant. 531.
Oliv. Inſ. 83. tab. fig. 76.
Geoff. Inſ. 1. 280. 6.
Herbſt. Arch. tab. 24. fig. 1.
Ent. Helv. 1. 64. 3. tab. 7. fig. 1. 2.
Schaeff. Icon. tab. 25. fig. 9. tab. 62. fig. 10.
Payk. Monogr. 33. 32.

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 223. 42.
Panz. Fn. Germ. 42. tab. 2.
Habitat in Germaniae plantis.

*2maculatus.* 16. R. fuſcus, elytris puncto cinereo, roſtro pedibusque atris.
*Curculio 2maculatus.* Ent. ſyſt. 2. 401. 30. *
Habitat Halae Saxonum. Dom. Hybner.

*punctum.* 17. R. ater, elytris ſtriatis: puncto medio albo, antennis pedibusque rubris.
*Curculio punctum.* Ent. ſyſt. 2. 401. 31. *
Habitat Halae Saxonum. Dom. Hybner.

*ſerratus.* 18. R. ferrugineus, thorace varioloſo, elytris ſtriatis: ſtriis ſubſerratis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura praecedentium. Roſtrum incuruum. Caput et thorax varioloſo-punctata, brunnea. Elytra paullo pallidiora, ſtriata, ſtriis dentibus breuibus, acutis ſubſerrata ſulcisque punctatis.

*biguttatus* 19. R. niger, elytris puncto eleuato, abdomine pedibusque poſticis flauis.
*Curculio 2guttatus.* Ent. ſyſt. 2. 402. 32. *
Habitat in America. Muſ. Dr. Hunter.

*bilineatus.* 20. R. fuſcus, elytris lineis duabus punctorum alborum.
*Curculio bilineatus.* Ent. ſyſt. 2. 402. 33. *
Habitat in Germania. D. de Hattorff.

*ſordidus.* 21. R. niger, obſcurus, immaculatus, elytris ſtriatis.
*Curculio ſordidus.* Ent. ſyſt. 2. 402. 34. *
Habitat in Americae meridionalis Inſulis. Dr. Pflug.

*ſuillus.* 22. R. obſcurus, thorace ſcabro, elytrorum ſtriis piloſis.

*Cur-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio ſuillus.* Ent. ſyſt. 2. 402. 35. *
Habitat in Americae meridionalis Inſulis. Dom. Smidt.

3. R. cinereus, elytris apice ſtriis albis nigro-punctatis. *teſſellatus.*
*Curculio teſſellatus.* Ent. ſyſt. 2. 403. 36. *
Habitat in Germania. D. de Hattorff.

4. R. thorace laeui, elytris muricatis, nigris: punctis duobus apiceque albis. *Equiſeti.*
*Curculio Equiſeti.* Ent. ſyſt. 2. 403. 37. *
*Curculio ſcaber.* Linn. ſyſt. Nat. 2. 609. 22.
Fn. Sv. 592.
Herbſt. Arch. tab. 24. fig. 2.
Panz. Fn. Germ. 4. tab. 4.
Degeer Inſ. 5. 224. 17.
Payk. Monogr. 39. 36.
Fn. Sv. 3. 226. 44.
Herbſt. Col. 6. 287. 258. tab. 82. fig. 4.
Habitat in Angliae et Germaniae Equiſeto aruenſi.

5. R. niger, obſcurus, thorace faſciculis duabus, elytris tribus eleuatis. *faſciculatus.*
Habitat in Amboina. D. Billardiere.
Statura R. Equiſeti, at maior. Caput nigrum, roſtro cylindrico, inflexo, baſi parum ferrugineo-villoſo. Thorax niger, obſcurus, dorſo plano, canaliculato, pari tuberculorum e pilis eleuatis, faſciculatis, nigris. Elytra ſubuarioloſa, nigra, tuberculis tribus e pilis faſciculatis, vno pone alterum.

6. R. niger, obſcurus, elytris granulatis, poſtice biſpinoſis. *granulatus.*
Habitat in Amboina. D. Billardiere.
Statura et magnitudo praecedentis. Roſtrum incuruum, fuſcum. Thorax punctis maioribus, ele-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

eleuatis, granulatis. Elytra granulata, poſtice ſpinis duabus eleuatis, validis, vna pone alteram. Pedes inermes.

*dimidiatus.* 27. R. niger, elytris ſubſtriatis, rufis.
*Curculio dimidiatus.* Ent. ſyſt. 2. 403. 40. *
Oliv. Inſ. 83. tab. 1. fig. 5.
Habitat - - -

*ſubſpinoſus.* 28. R. obſcurus, thorace varioloſo, elytris ſubſpinoſis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Medius. Roſtrum elongatum, fuſcum. Thorax punctis varioloſis confluentibus. Elytra ſpinis paruis, eleuatis ſtriata.

*palpebra.* 29. R. griſeus, fuſco-maculatus, thoracis margine antico orbitaque oculorum eleuatis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura praecedentium. Roſtrum elongatum, incuruum. Orbita oculorum eleuata. Thorax rotundatus, villoſus, teſtaceo parum maculatus: margine eleuato. Elytra ſtriata, villoſa griſea, teſtaceo parum maculata. Pedes griſei.

*Bufo.* 30. R. fuſcus, elytris ſubreticulatis: faſcia media alba.
*Curculio Bufo.* Ent. ſyſt. 2. 404. 41. *
Oliv. Inſ. 83. fig. 118.
Habitat in Sibiria. Muſ. D. Banks.

*famulus.* 31. R. ſupra ferrugineus, thoracis dorſo elytrorumque ſutura inaequaliter atris.
*Curculio famulus.* Ent. ſyſt. ſuppl. 165. 41.
Habitat Tranquebariae. Muſ. D. Lund.

*armillatus* 32. R. ater, elytris cinereo-vndatis, tibiis poſticis apice albis.

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Tanger. D. Sc. ousboe. Muſ. D. de Seheſtedt.

Maior R. atriroſtris. Roſtrum cylindricum, incuruum, atrum. Thorax punctatus, ater. Elytra ſtriata, atra, ſtrigis faſciisque plurimis, fere obſoletis, cinereis vndata. Pedes inermes, femoribus quatuor poſticis annulo baſeos apice albis.

33. R. ater, thorace plano, elytris ſtriatis, abdomine ferrugineo. *Muſculus.*

Habitat in Sumatra. D. Daldorff.

Magnitudo R. atriroſtris. Caput atrum, roſtro arcuato, elongato, piceo. Antennae nigrae, clauа cinerea. Thorax planus, laeuis, ater, immaculatus. Elytra ſtriata, nigra. Corpus nigrum, abdomine rufo. Pedes picei.

34. R. cinereus, roſtro arcuato atro. *atriroſtris*

*Curculio atriroſtris* Ent. ſyſt. 2. 404. 42. *

Payk. Monogr. 40. 37.

Fn Sv. 3 227. 45.

Habitat Lipſiae. Muſ. D. Prof. Leske.

35. R. griſeus, roſtro pedibusque ferrugineis. *brunniroſtris.*

*Curculio brunniroſtris* Ent. ſyſt. 2. 404. 43.*

Habitat in Daniae plantis. Muſ. D. Lund.

36. R. ſupra ater, thorace lineis tribus, elytris vittis duabus albis. *4vittatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Medius. Roſtrum valde elongatum, arcuatum, tenue, atrum. Thorax ater, lineis tribus lutis albidis. Elytra vix ſtriata, atra, vittis duabus poſtice coëuntibus, albidis. Corpus cinereum, pedibus atris.

37. R. aeneo-nitidus, elytris baſi angulo obtuſo, roſtro, apice antennisque fuſcis. *feſtiuus.*

*Cur-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio feſtiuus.* Ent. ſyſt. 2. 404. 44.
Habitat in Surinamo.

*Tragiae.* 38. R. aeneus, roſtro pedibusque ſubconcoloribus.
*Curculio Tragiae.* Ent. ſyſt. 2. 404. 45.
Oliv. Inſ. 83. fig. 112.
Habitat in Tragiae volubilis ſeminibus Braſiliae. Muſ. D. Banks.

*Chloris.* 39. R. thorace elytrisque obſcure viridibus, corpore nigro.
*Curculio Chloris.* Ent. ſyſt. ſuppl. 144. 46.
*Tr. 64.* Panz. Fn. Germ. 18. tab. 8.
Habitat in Germania. Dr. Panzer.

*curuiro-ſtris.* 40. R. ater, nitidus, roſtro arcuato.
*Curculio curuiroſtris.* Ent. ſyſt. 2. 405. 47.*
Oliv. Inſ. 83. fig. 115.
Habitat in noua Hollandia. Muſ. D. Banks.

*Aethiops.* 41. R. ater, antennis tibiisque piceis, elytris oblongis ſtriatis.
*Curculio Aethiops* Ent. ſyſt. 2. 405. 48.*
Payk. Monogr. 29. 46.
Fn. Sv. 3. 236. 54.
Herbſt. Col. 6. 270. 241. tab. 80. fig. 10.
Habitat in Suecia. D. de Paykull.

*Rubi.* 42. R. niger, thorace antice anguſtato, elytris oblongis ſtriatis.
*Curculio Rubi.* Herbſt. Col. 6. 271. 242.
Habitat in Germaniae Rubo. D. de Meyerle.
Statura omnino R. Aethiopis, at paullo minor. Roſtrum elongatum, tenue, cylindricum, atrum. Thorax punctatus, niger, antice anguſtatus. Scutellum paruum, cinereum. Elytra ouata, ſtriata, nigra. Pedes nigri.

*Scirpi.* 43. R. fuſcus, elytris ſubſtriatis ferrugineo-irroratis.

*Cur-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Scirpi.* Ent. ſyſt. 2. 405. 49. *
Habitat in Galliae Scirpo. Muſ. D. Boſc.

4. R. ater, nitidus, elytris ſtriatis: puncto baſeos vittaque media albis. *Vnca.*
Habitat in Iaua. Muſ. D. Billardiere.
Statura et magnitudo R. Scirpi. Antennae nigrae: articulo primo rufo. Caput nigrum, roſtro elongato, incuruo. Thorax laeuis, ater, vtrinque linea albida. Elytra ſtriata, abdomine breuiora, atra: puncto oblongo baſeos vittaque media albis. Pedes nigri.

5. R. ater, antennis ferrugineis, thorace bitubercu- *Pruni.*
lato.
*Curculio Pruni.* Ent. ſyſt. 2. 405. 50. *
Linn. ſyſt. Nat. 2. 507. 12.
Payk. Monogr. 137. 128.
Fn. Sv. 3. 246. 45.
Herbſt. Col. 6. 445. 443.
Habitat in Pruni Ceraſi foliis.

6. R. ater, immaculatus, elytris crenato-ſtriatis. *Armeniacae.*
*Curculio Armeniacae.* Ent. ſyſt. 2. 405. 51.
Habitat Kiliae in Pruni foliis.

7. R. fuſcus, thorace quadrituberculato, elytris ſtriatis tuberculatisque. *5 tuberculatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura R. Cameli. Roſtrum inflexum. Thorax margine antico parum reflexo tuberculisque quatuor in medio eleuatis: intermediis duobus maioribus, compreſſis. Elytra ſtriata tuberculisque aliquot eleuatis, ſparſis.

8. R. fuſcus, thorace elytrisque tuberculatis, roſtro apice rufo. *Camelus.*
*Curculio Camelus.* Ent. ſyſt. 2. 406. 52. *

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Germaniae hortis. D. Smidt.

*4tuberculatus.* 49. R. thorace quadrituberculato nigro, elytris ſtriatis, cinereo-variegatis.
*Curculio 4tuberculatus.* Ent. ſyſt. 2. 406. 53
Payk. Monogr. 71. 69.
Fn. Sv. 3. 258. 79.
Herbſt. Col. 6. 409. 390. tab. 92. fig. 3.
Habitat in Europa boreali. D. Daldorff.

*teter.* 50. R. depreſſus, villoſus, fuſcus, roſtro atro.
*Curculio teter.* Ent. ſyſt. 2. 406. 54. *
III. 61. *Curculio Linariae.* Panz. Fn. Germ. 26. tab. 18
Habitat ad radices Antirrhini Linariae Italiae et Germaniae, Gallas excitans. Dr. Allioni.

*Chamaeropis.* 51. R teſtaceus, capite thoracis macula pectoreque nigris. Ent. ſyſt. ſuppl. 164. 54 *
Habitat in floribus Chamaeropis humilis Barbariae. D. Schousboe. Muſ D. Lund.

*Campanulae.* 52. R. ouatus, niger, elytris ſtriatis obtuſis.
*Curculio Campanulae* Ent. ſyſt. 2. 406. 55.
Linn. ſyſt. Nat. 2. 506. 7.
Payk. Monogr. 28. 27.
Fn. Sv. 3. 212. 29.
Degeer Inſ. 5. 236. 7.
Herbſt. Col. 6. 161. 119. tab. 70. fig. 15.
Habitat in Campanulae rotundifoliae pericarpiis.

*nigriroſtris.* 53. R. viridis, roſtro atro.
*Curculio nigriroſtris.* Ent. ſyſt. 2. 407. 56. *
Payk. Monogr. 56. 58.
Fn. Sv. 3. 247. 47.
Herbſt. Col. 6. 281. 254.
Arch. tab. 24. fig. 3.
III. 60. Panz. Fn Germ. 36. tab. 14.
Habitat in Anglia. Muſ. D. Banks.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. R. ſubteſtaceus, thorace viridi-lineato, roſtro apice fuſco. *variabilis.*

*Curculio variabilis.* Ent. ſyſt. 2. 407. 57. *
*Curculio nigriroſtris.* Payk. Monogr. 57. 53. β.
Fn. Sv. 3 247. 47. β.
Herbſt. Col. 6. 283. 254. tab. 81. fig. 12.
Habitat in Europa boreali. Dr. Schulz.

5. R. oblongus, niger, argenteo-holoſericeus, roſtro dimidiato pedibusque piceis. *piciroſtris.*

*Curculio piciroſtris.* Ent. ſyſt. 2. 407. 58. *
Payk. Monogr. 63. 60.
Fn. Sv. 3. 253. 73.
Herbſt. Col. 6. 446. 446.
Habitat in Trifolii capitulis.

6. R. cinereus, thorace elytrisque cinereo-virentibus, roſtro nigro. *Viſcariae.*

*Curculio Viſcariae.* Linn. ſyſt. Nat. 2. 609. 27.
Fn. Sv. 598.
Payk. Fn. Sv. 3. 261. 82.
Habitat in floribus Lychnidis Viſcariae.
Paruus. Roſtrum elongatum, cylindricum, incuruum, apice atrum. Thorax vireſcens. Elytra vireſcentia, ſtriata. Corpus cinereum.

7. R. cinereus, thorace elytrisque cinereo-virentibus, ſcutello niueo. *bifrons.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo R. Viſcariae. Caput cum roſtro longo, arcuato, atrum. Thorax punctatus, elytraque ſtriata, vireſcentia. Scutellum album. Corpus cinereum, pedibus nigricantibus.

8. R. niger, antennarum baſi, coleoptrorum diſco tibiisque teſtaceis. *Salicariae.*



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Salicariae.* Ent. ſyſt. 2. 407. 58. *
Payk. Monogr. 74. 48. β.
Fn. Sv. 3. 363. 85. β.
Panz. Fn. Germ. 17. tab. 4.
Habitat in Lythro Salicaria Angliae.

*floralis.* 59. R. obſcure griſeus, elytrorum futura pallida.
*Curculio floralis.* Ent. ſyſt. 2. 408. 59. *
Payk. Monogr. 77. 74.
Fn. Sv. 3. 266. 88.
Habitat in Pomonae floribus. Dom. de Paykull

*Vmbellae.* 60. R. nigricans, elytris ſtriatis, ſcutello albo.
Habitat in Carolina. Muſ. Dom. Boſc.
Statura R. pericarpii, at maior et alius. Caput grum. Thorax laeuis, nigricans, immacu tus. Elytra parum pallidiora, ſtriata: ſtriis te apicem coëuntibus. Scutellum album, at tura elytris omnino concolor. Corpus ci reum, pedibus nigris.

*Vulpeculus* 61. R. ſupra niger, elytrorum futura corpore ſubtus brunneis.
Habitat in Carolina. Muſ. Dom. Boſc.
Statura omnino R. pſeudacori. Caput nigr apice retuſum. Roſtrum longum. arcuat nigrum. Thorax lineola abbreuiata, media ſeos alba. Elytra ſubſtriata, nigra, futura f ruginea lineolisque duabus ad futuram baſ albis. Corpus ferrugineo-villoſum. Pedes ſcuri.

*Pſeudacori* 62. R. ſupra niger, thoracis lateribus ferrugin elytris ſtriatis: futura baſi alba.
*Curculio Pſeudacori.* Ent. ſyſt. 2. 408. 61.
Panz. Fn. Germ. 17. tab. 5.
Habitat in Galliae Pſeudacoro. Muſ. Dom. Bo

*ſcanicus.* 63. R. teſtaceus, coleoptrorum diſco fuſco: futu alba.

C

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio ſcanicus.* Payk. Monogr. 59. 56.
Payk. Fn. Sv. 3. 251. 70. IX.
Habitat in Scania. Muſ. D. Lund.
Minor R. pſeudacori. Caput et thorax teſtacea, vertice nigro. Elytra ſtriata, teſtacea, ad ſuturam fuſca, ſutura ipſa, lunulisque aliquot indiſtinctis, albis. Pedes teſtacei.

64. R. ouatus, thorace tuberculato, coleoptris ſtriatis: ſutura baſi albida, pedibus ru s. *Caſtor.*
*Curculio Caſtor.* Ent. ſyſt. 2. 408. 62. *
Habitat in Germania. D. Smidt.

65. R. ſubgloboſus, nebuloſus, elytrorum ſutura baſi alba. *pericarpius.*
*Curculio pericarpius.* Ent. ſyſt. 2. 408. 63.
Linn. ſyſt. Nat. 2. 609. 31. Fn. Sv. 602.
Panz. Fn. Germ. 42. tab. 5.
Herbſt. Arch. tab. 24. fig. 4. IV. 69
Coleopt. 6. 401. 381. tab. 91. fig. 12.
Payk. Monogr. 67. 65.
Fn. Sv. 3 255. 75.
Habitat in Scrophulariis.

66. R. ater, thorace tuberculato, elytris ſtriatis: ſutura baſi alba, pedibus nigris. *quercicola.*
*Curculio quercicola.* Ent. ſyſt. 2. 408. 64. *
Payk. Monogr. 70. 68.
Fn. Sv. 3. 215. 33.
Herbſt. Col. 6. 403. 383. tab. 91. fig. 13.
Habitat in Sueciae Quercu. Dom. de Paykull.

67. R. thorace bituberculato canaliculatoque, elytris ſtriatis. *aſſimilis.*
*Curculio aſſimilis.* Ent. ſyſt. 2. 409. 65.
Panz. Fn. Germ. 42. tab. 6. IV. 69
Payk. Monogr. 69. 67.
Fn. Sv. 3. 257. 77.
Habitat in Suecia. Dom. de Paykull.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Sisymbrii.* 68. R. albo fuscoque varius, elytris puncto baseos eleuato, atro, rostro nigro.
*Curculio Sisymbrii.* Ent. syst. 2. 409. 66. *
Panz. Fn. Germ. 17. tab. 6.
Payk. Monogr. 55. 52.
Fn. Sv. 3. 247. 66.
Herbst. Coleopt. 6. 159. 116. tab. 70. fig. 12.
Habitat in Sisymbrio amphibio Kiliae.

*Capreae.* 69. R. coleoptris fasciis duabus abbreuiatis, albis.
*Curculio Capreae.* Ent. syst. 2. 409. 67. *
Habitat in Salice caprea Angliae. Mus. D. Banks.

*2punctatus* 70. R. villosus, cinereus, elytris macula media fusca.
*Curculio 2punctatus.* Ent. syst. 2. 409. 68.
Linn. syst. Nat. 2. 609. 28. Fn. Sv. 599.
Panz. Fn. Germ 42. tab. 7.
Payk. Monogr. 58. 55.
Fn. Sv. 3. 250. 69.
Herbst. Coleopt. 6. 234. 198. tab. 77. fig. 2.
Habitat in Salicis floribus. Dom. de Paykull.

*Carpini.* 71. R. villosus, virescens, rostro nigro, pedibus testaceis.
*Curculio Carpini.* Ent. syst. 2. 409. 69.
*Curculio Carpini.* Knoch. Beytr. 1.
Habitat in Germaniae Carpino. Mus. D. de Schestedt.

*Erysimi.* 72. R. niger, thorace bituberculato, virescente, elytris cyaneis.
*Curculio Erysimi.* Ent. syst. 2. 210. 70. *
Panz. Fn. Germ. 17. tab. 7.
Payk. Monogr. 76. 73.
Fn. Sv. 3. 265. 87.
Herbst. Coleopt. 6. 407. 388. tab. 92. fig. 1.
Habitat Kiliae. D. Daldorff.

*testaceus.* 73. R. testaceus, rostro elongato arcuato.

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Paruus, totus teſtaceus, immaculatus. Roſtrum elongatum, arcuatum, tenue. Elytra ſtriata.

4. R. nigricans, coleoptris maculis quatuor albidis. *4maculatus.*
*Curculio 4maculatus* Ent. ſyſt. 2. 410. 71.
Linn. ſyſt. Nat. 2. 609. 29. Fn. Sv. 600.
Geoff. Inſ. 1. 187. 22.
Habitat in Europae plantis.

5. R. ſupra fuſcus, elytris faſcia media fuſca. *vnifaſciatus.*
*Curculio vnifaſciatus* Ent. ſyſt. 2. 410. 72. *
Habitat Halae Saxonum. Dom. Hybner.

6. R. ater, elytris faſcia media abbreuiata puncto- *Lythri.*
que poſtico albis, pedibus flauis.
*Curculio Lythri.* Ent. ſyſt. 2. 410. 73. *
*Cionus Lythri.* Ent. Helv. 1. 68. 2. tab. 3. fig. 3. 4.
Panz. Fn. Germ. 17. tab. 8.
*Curculio pygmaeus.* Herbſt. Coleopt. 6. 142. 102. tab. 69. fig. 7.
Payk. Monogr. 74. 71.
Fn. Sv. 3. 264. 85.
Habitat in Lythri floribus Sueciae. Dom. de Paykull.

7. R. obſcurus, thorace tuberculato, elytris crena- *vmbraculatus.*
to-ſtriatis, oculorum orbita eleuata.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus paruum, breue, gibbum. Roſtrum longum, arcuatum, canaliculatum, orbita oculorum valde eleuata. Thorax obſcure-griſeus, linea media ſubcompreſſa tuberculoque vtrinque eleuato. Elytra breuia, crenato-ſtriata. Pedes nigri.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*2faſciatus.* 78. R. niger, elytris faſciis duabus cinereis: baſeos maiore vndata.
*Curculio 2faſciatus.* Ent. ſyſt. 2. 410. 74. *
Habitat Kiliae. Muſ. Dom. Lund.

*acridulus.* 79. R. niger, antennis pedibusque piceis, abdomine ouato.
*Curculio acridulus.* Ent. ſyſt. 2. 411. 75. *
Linn. ſyſt. Nat. 2. 507. 13. Fn. Sv. 584.
Degeer Inſ. 5. 235. 22.
*IV. 76* Panz. Fn. Germ. 42. tab. 10.
Herbſt Arch. tab. 24. fig. 12.
Coleopt. 6. 272. 244. tab. 81. fig. 1.
Payk. Monogr. 50. 47.
Fn. Sv. 3. 238 56.
Habitat in Europae floribus.

*ſubrufus.* 80. R. teſtaceus, thorace tuberculato, elytris ſtriatis: ſtrigis cineraſcentibus.
*Curculio ſubrufus.* Herbſt. Col. tab. 92. fig. 14.
Habitat in Zelandiae Quercu. Muſ. D. Lund.
Statura parua praecedentium. Corpus totum teſtaceum, nunc magis, nunc minus pallidum: ſtrigis aliquot elytrorum indiſtinctis, cinereis. Thorax vtrinque tuberculo eleuato.

*ſcabratus.* 81. R. fuſcus, thorace tuberculato, elytris ſcabris, pedibus piceis.
*Curculio ſcabratus.* Ent. ſyſt. 2. 411. 76.
Habitat in Germania. D. Smidt.

*alauda.* 82. R. ſupra ater, thorace bituberculato margineque antico eleuato.
*Curculio alauda.* Ent. ſyſt. 2. 411. 77. *
*V. 16* *Curculio affinis* Payk. Monogr. 68. 66.
Habitat in Germania. Dom. Smidt.

*dorſalis.* 83 R. elytris rubris, ſutura dimidiata, nigra.
*Curculio dorſalis.* Ent. ſyſt. 2. 411. 78. *
Linn.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. syst. Nat. 2. 608. 17. Fn. Sv. 588.
Panz. Fn. Germ 17. tab. 9.
Oliv. Inf. 83. fig. 169.
Payk. Monogr. 14. 73.
Habitat in Ranunculo ficaria.

4. R. cinereus, thoracis dorso fusco, elytris testaceis, cinereo-vndatis. *Quercus.*
*Curculio Quercus.* Ent. syst. 2. 412. 79. *
Linn. syst. Nat. 2. 614. 59.
Herbst. Col. 6. tab. 92. fig. 7.
Payk. Monogr. 72. 70.
Fn. Sv. 3. 262. 83.
Habitat in Kiliae Quercu.

5. R. fuscus, elytris puncto strigisque duabus vndatis, albidis. *Tigris.*
Habitat in America meridionali. Mus. D. de Schestedt.
Paruus. Rostrum minus elongatum, cinereum. Thorax obscurus, fuscus, linea dorsali cinerea. Elytra striata, fusca, macula baseos fasciisque duabus vndatis, cinereis. Apex parum albicat. Corpus cinereum.

6. R. ouatus, fuscus: linea longitudinali alba. *futuralis.*
*Curculio futuralis.* Ent. syst. 2. 412. 80. *
Habitat in Germaniae Salice.

7. R. ater, thorace punctis duobus baseos, elytris futura punctisque sparsis albis. *Crux.*
*Curculio Crux.* Ent. syst. 2. 412. 81. *
Habitat Hamburgi. Dr. Schulz.

8. R. ater, rostro apice plano, elytris striatis. *Lemnae.*
*Curculio Lemnae.* Ent. syst. 2. 412. 82. *
Panz. Fn. Germ. 17. tab. 10.
Herbst Col. 6. 453. 457.
Payk. Monogr. 78. 75.

Payk.

Digitized by Google. Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 3. 267. 89.
Habitat in Europae borealis Lemna. Dom. Prof. Helwig.

*exclamationis.* 89. R. ater, elytris puncto lineolaque apicis albis.
*Curculio exclamationis.* Ent. syst. 2. 412. 83.
Oliv. Inf. 83. fig. 167.
Habitat in noua Hollandia. Mus. D. Banks.

*venustus.* 90. R. fuscus, thorace elytrisque albo-lineatis, pedibus testaceis.
*Curculio venustus.* Ent. syst. 2. 412. 84.
*Curculio albovittatus.* Herbst. Archiv.
Habitat in Anglia. Mus. D. Banks.

*Plantaginis.* 91. R. elytris cinereis: macula media fusca.
*Curculio Plantaginis.* Ent. syst. 2. 415. 85.
Degeer Inf. 5. 237. 24. tab. 7. fig. 17. 18.
Payk. Monogr. 44. 41.
Fn. Sv. 3. 231. 48.
Herbst. Col. 6. tab. 70. fig. 2.
Habitat Halae. Dom. Hybner.

*Artemisiae* 92. glaber, ater, thorace punctato, elytrorum striis simplicibus.
*Curculio Artemisiae.* Payk. Fn. Sv. 3. 237. 55.
Herbst. Col. 6. tab. 76. fig. 5.
Habitat in Sueciae Artemisia.
Statura R. aethiopis, at duplo minor. Thorax punctis latius impressus. Elytra striis laeuibus.

*Rumicis.* 93. R. griseus, nigro-nebulosus, antennis fuscis.
*Curculio Rumicis.* Ent. syst. 2. 413. 86.
Linn. syst. Nat. 2. 613. 60. Fn. Sv. 390.
Degeer Inf. 5. 531. 20. tab. 7. fig. 10. 11.
Payk. Monogr. 43. 40.
Fn. Sv. 3. 229. 47.
Habitat in Europae borealis Rumice.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

94. R. griſeus, thorace nigro, cinereo, lineato, elytris punctis nigris, teſſellatis. *Pollux.*
*Curculio adſperſus.* Ent. ſyſt. 2. 413. 87. *
Habitat Kiliae.
Affinis certe R. Rumicis, cuius forte mera varietas.

** *femoribus dentatis.*

95. R. femoribus dentatis, niger, elytris fuſco-irroratis, tibiis anticis acute-dentatis. *calcaratus.*
*Curculio calcaratus.* Ent. ſyſt. 2. 420. 114. *
Oliv. Inſ. 83. tab. 15. fig. 186.
Habitat - - - Muſ. Dom. Geoffroy.

96. R. femoribus poſticis dentatis, niger, elytris vniſpinoſis. *bidens.*
*Curculio bidens.* Ent. ſyſt. 2. 420. 115. *
Oliv. Inſ. 83. fig. 113.
Habitat in noua Zelandia. Muſ. Dom. Banks.

97. R. femoribus dentatis, griſeus, thorace elytriſque tuberculatis, roſtro cornubus duobus arcuatis. *taurus.*
*Curculio taurus.* Ent. ſyſt. 2. 421. 116. *
Oliv. Inſ. 83. fig. 60.
Habitat Cayennae.

98. R. femoribus dentatis, thorace tuberculato, roſtro vtrinque acute-ſpinoſo. *cornutus.*
*Curculio cornutus.* Ent. ſyſt. 2. 421. 117.
Oliv. Inſ. 83. fig. 193.
Habitat Cayennae. Muſ. Dom. Geoffroy.

99. R. femoribus dentatis, fuſcis, thorace elytriſque tuberculis numeroſis, atris. *miliaris.*
*Curculio miliaris.* Ent. ſyſt. 2. 421. 118. *
Oliv. Inſ. 83. fig. 83.
Habitat Cayennae. Muſ. Dom. Boſc.

100.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*cyanicollis.* 100. R. femoribus dentatis, oblongus, nigricans, thorace cyaneo: lateribus ſcabris, elytris ſtriatis.
*Curculio cyanicollis.* Ent. ſyſt. 2. 421. 119.
Oliv. Inſ. 83. fig. 121.
Habitat - - - Muſ. Dr. Hunter.

*Jamaicenſis.* 101. R. femoribus dentatis, obſcurus, ſcaber, thorace vtrinque faſciculato, tuberculato, elytris ſtriatis.
*Curculio Jamaicenſis.* Ent. ſyſt. 2. 421. 120.*
Oliv. Inſ. 83 tab. 4. fig. 44.
Habitat in America meridionali. Muſ. D. Banks.

*vaginalis.* 102. R. femoribus dentatis, ater, elytris tuberculatis acuminatisque: maculis duabus tomentoſo-aureis.
*Curculio vaginalis*, longiroſtris, femoribus dentatis, elytris ſtriatis: tuberculis ſparſis, iuxta apicem mucronatis.
Linn. ſyſt. Nat. 2. 612. 51.
*Curculio Senegalenſis.* Ent. ſyſt. 2. 422. 121.*
Degeer Inſ. 5. tab. 15. fig. 23.
Habitat in America meridionali.
*Mas* cornubus duobus validis arcuatis, ad baſin roſtri.

*validus.* 103. R. femoribus anticis dentatis, oblongus, ſcaber, niger, tibiis anticis dentatis.
*Curculio validus.* Ent. ſyſt. 2. 422. 122.♂
Oliv. Inſ. 83. fig. 186.♀
Habitat in Cayenna.

*Veſpertilio* 104. R. femoribus dentatis, niger, roſtro plano, vtrinque dente obtuſo, elytris tuberculatis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Minor quam R. vaginalis. Roſtrum atrum, planum, in medio vtrinque dente valido, obtuſo, de-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

depreſſo, margineque a dente ad apicem reflexo. Thorax laeuis, niger, obſcurus. Elytra punctata, ſtriata, ſtriis tuberculis eleuatis, acutis, ſerrata. Pedes atri, tarſis vtrinque auro ciliatis.

105. R. femoribus dentatis, niger, thorace antice ſpinis ciliato, elytris ſtriatis. *coronatus.*

*Curculio coronatus.* Ent. ſyſt. 2. 422. 123. *
Oliv. Inſ. 83. fig. 70.
Habitat - - -

106. R. femoribus quatuor anticis dentatis, elytris polline flaueſcentibus, ſupra apicem gibboſis. *mucoreus.*

*Curculio mucoreus.* Ent. ſyſt. 2. 423. 124.
Linn. ſyſt. Nat. 2. 612. 49. Muſ. Lud. Vlr. 53.
Habitat in India orientali.

107. R. femoribus quatuor poſticis dentatis, elytris ſtriatis nigris: linea repanda griſea. *puſio.*

*Curculio puſio.* Ent. ſyſt. 2. 423. 125.
Linn. ſyſt. Nat. 2. 612. 50. Muſ. Lud. Vlr. 46.
Oliv. Inſ. 83. tab. 16. fig. 206.
Habitat in India orientali.

108. R. femoribus dentatis, niger, thorace lineis duabus, elytris ſtrigis quatuor albis, tibiis anticis ſpinoſis. *ſpinipes.*

*Curculio ſpinipes.* Ent. ſyſt. 2. 423. 126. *
Habitat in Americae meridionalis Inſulis. Muſ. Dr. Hunter.

109. R. femoribus dentatis, fuſco-griſeus, thorace elytrisque vitta laterali flaueſcente. *Rana.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Corpus medium, ſupra planum, fuſco-griſeum, punctis paruis, eleuatis, numeroſis, atris, ſcabrum, vitta laterali ab apice thoracis ad apicem ely-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

elytrorum ducta flaua. Roſtrum elongatum, arcuatum, atrum, apice vtrinque parum dilatatum. Corpus ſubtus ferrugineo-villoſum.

*diſcoides.* 110. R. femoribus dentatis, laeuis, ater, thorace elytrisque vitta laterali flaueſcente.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentis. Roſtrum breue, compreſſum, atrum, antennarum claua cinereſcente. Thorax planus, laeuis, ater, vitta laterali flaueſcente, albo-marginata. Elytra laeuia, atra: vitta lata laterali, quae ad apicem dilatatur, totum apicem occupat. Corpus atrum.

*lineatus.* 111. R. femoribus dentatis, ater, thorace elytrisque ſtriatis: vitta laterali flaueſcente.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura R. Ranae, at duplo minor. Roſtrum elongatum, atrum. Thorax ater, vittis duabus flauis. Elytra ſtriata, atra: vitta media flaua. Corpus atrum, vtrinque vitta communi, maculaque magna ad baſin abdominis, flauis. Pedes atri.

*bombinus.* 112. R. femoribus dentatis, ferrugineo-fuſcus, elytris ſtriatis: tuberculis eleuatis atris.

*Curculio bombinus.* Ent. ſyſt. 2. 423. 127. *

Oliv. Inſ. 83. fig. 12.

Habitat in Cayenna. D. v. Rohr.

*clauipes.* 113. R. femoribus clauatis dentatis, ſcaber, nigro pallidoque varius, elytris poſtice tuberculatis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Medius. Roſtrum elongatum, incuruum, atrum. Thorax punctis eleuatis, atris ſcaber, fuſco cinereoque va-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

varius. Elytra punctis ſcabra, variegata, poſtice tuberculo obtuſo. Pedes obſcuri, femoribus omnibus elongatis, clauatis, dentatis.

14. R. femoribus clauatis, dentatis, nigricans, elytris fuſcis: macula media holoſericea, atra. *teſtudo.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo praecedentis. Roſtrum breuius, vtrinque ſulco exaratum, nigricans, obſcurum. Thorax ſcaber, niger, lateribus obſcure brunneis. Elytra punctato-ſtriata, fuſca, macula media holoſericea atra, apice parum acuminata. Femora valde elongata, clauata, dentata, poſtice compreſſa.

15. R. femoribus dentatis, ſcaber, nigro ferrugineoque varius, elytris punctis ocellaribus ſtriatis. *chamaeleon.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentium. Roſtrum elongatum, incuruum, atrum. Thorax ſcaber, ater, ferrugineo-varius. Elytra varia, ſtriata, punctis impreſſis atris: pupilla parua ferruginea. Femora parum elongata, ſubclauata, dentata.

16. R. femoribus dentatis, fuſcus, thorace lateribus albis, elytris ſtriatis: apice macula niuea, apice teſtaceis. *Draco.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Praecedentibus paullo minor. Roſtrum breue cylindricum, obſcure brunneum. Thorax ſcaber, obſcurus, lateribus late albis. Elytra ſtriata, obſcura: macula magna lunata baſeos niuea, apice late teſtacea. Pedes obſcure ferruginei, femoribus clauatis, dentatis.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Scorpio.* 117. R. femoribus dentatis, ater, thorace plano basi cinereo, coleoptris tuberculatis acuminatisque, medio cinereis.
*Curculio Scorpio* Ent. syst. 2. 424. 128. ♂
Oliv. Inf. 83. fig. 14.
Habitat in Cayenna. D. v. Rohr.

*Chimaridis.* 118. R. femoribus posticis dentatis, niger, albo-maculatus, pedibus anticis longissimis.
*Curculio Chimaridis.* Ent. syst. 2. 424. 129.
Oliv. Inf. 83. 130.
Habitat in Americae meridionalis Chimaride carnosa. Dr. Isert.

*guttatus.* 119. R. femoribus subdentatis, niger, thorace maculis duabus baseos, elytris tuberculatis: punctis albidis.
*Curculio guttatus.* Ent. syst. 2. 424. 130. ♂
Oliv. Inf. 83. tab. 5. fig. 45.
Habitat in Cayenna.

*fascicularis.* 120. R. femoribus dentatis, niger, elytris fasciculato-pilosis, pedibus cinereo-variis.
*Curculio fasciicularis.* Ent. syst. 2. 424. 131. ♂
Oliv. Inf. 83. tab. 1. fig. 9.
Habitat in Cayenna.

*Hystrix.* 121. R. femoribus dentatis, niger, elytris crenato-striatis: punctis duobus albis.
*Curculio Hystrix* Ent. syst. 2. 425. 132. ♂
Oliv. Inf. 83. tab. 15. fig. 182.
Habitat in Cayenna. Mus. D. Geoffroy.

*marmoreus.* 122. R. femoribus dentatis, brunneus, albo-maculatus.
*Curculio marmoratus.* Ent. syst. 2. 425. 133.
Oliv. Inf. 83. tab. 13. fig. 153.
Habitat in Cayenna.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

123. R. femoribus dentatis, thorace carinato, elytris ſulcatis, tuberculato-ſpinoſis. *ſcaber.*
*Curculio ſcaber.* Ent. ſyſt. 2. 425. 134. *
Habitat in Cayenna. Dr. Mallet.

124. R. femoribus dentatis, thorace linea laterali, elytris albo-punctatis maculisque duabus brunneis. *3faſciatus.*
*Curculio 3faſciatus.* Ent. ſyſt. 2. 425. 135. *
Habitat in Cayenna. Muſ. Dr. Hunter.

125. R. femoribus dentatis, thorace ſcabro antice bicorni, elytris poſtice bituberculatis *cylindriroſtris.*
*Curculio cylindriroſtris.* Ent. ſyſt. 2. 426. 136.
Oliv. Inſ. 83. fig. 128.
Habitat in noua Hollandia. Muſ. D. Banks.
Thorax antice cornubus duobus, e faſciculis pilorum eleuatorum, quae aetate pereunt.

126. R. femoribus dentatis, elytris macula magna ferruginea. *ſtigma.*
*Curculio ſtigma* Ent. ſyſt. 2. 426. 137. *
*Curculio pudicus.* Ent. ſyſt. ſuppl. 165. 19.
*Curculio piger.* Ent. ſyſt. ſuppl. 169. 160.
Linn. ſyſt. Nat. 2. 612. 52. Muſ. Lud. Vlr. 48.
Habitat in Indiis.
*Variat* ſutura baſeos faſciaque media ferrugineis.

127. R, femoribus ſubdentatis, thorace ſcabro, elytris ſulcatis tuberculatis, ore barbato. *hebes.*
*Curculio hebes.* Ent. ſyſt. 2. 426. 138. *
Oliv. Inſ. 83. fig. 144.
Habitat in Bengala. Muſ. D. Banks.

128. R. femoribus dentatis, pallidus, thorace elytrisque ſtrigis nigris. *annulatus*
*Curculio annulatus.* Ent. ſyſt. 2. 426. 139. *
Linn. ſyſt. Nat. 2. 613. 54. Muſ. Lud. Vlr. 51.
Oliv.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Oliv. Inf. 83. fig. 62.
Habitat in India.

*angulus.* 129. R. femoribus dentatis, niger, elytris vitta abbreuiata, faſciaque poſtica cinereis, tibiis anticis compreſſis dentatis.
Habitat in Sumatra. D. Daldorff. Muſ. D. Lund.
Statura et magnitudo R. Abietis. Caput fuſcum. Thorax ſcaber, niger, immaculatus. Elytra punctato-ſtriata, nigra, vitta cinerea, a baſi ad faſciam ducta, faſciaque pone medium cinereis. Macula magna, cinerea, fere obſoleta, elytron terminat. Corpus cinereo-villoſum. Pedes nigri, antici elongati, tibiis quatuor anticis compreſſis, acute dentatis.

*Abietis.* 130. R. femoribus dentatis, niger, elytris faſciis linearibus, interruptis albis.
*Curculio Abietis.* Ent. ſyſt. 2. 428. 144. *
Linn. ſyſt. Nat. 2. 613. 57. Fn. Sv. 615.
Degeer Inſ. 5. 205. 1. tab. 6. fig. 11.
Panz. Fn. Germ. 42. tab. 14.
Schaeff. Icon. tab. 25. fig. 1.
Petiv. Gazoph. 14. tab. 8. fig. 9.
Habitat in Pini ſylueſtris cortice et reſina.

*Erinaceus.* 131. R. femoribus dentatis, niger, elytris albo-irroratis, tuberculisque duobus eleuatis piloſis, atris.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Magnitudo et ſtatura C. Abietis. Roſtrum inflexum, atrum. Thorax ſcaber, niger. Elytra ſtriata, nigra, atomis numeroſis albidis irrorata, tuberculis duobus eleuatis, e pilis faſciculatis, atris: altero baſeos, altero ante apicem. Corpus obſcurum.

*carinatus.* 132. R. femoribus dentatis, obſcurus, thorace carinato, elytris lineis tribus carinatis.

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Praecedente minor. Roſtrum arcuatum, atrum, baſi griſeum. Thorax ſubſcaber, obſcurus: linea media eleuata, connata. Elytra punctato-ſtriata, lineis tribus eleuatis, coſtatis. Pedes antici elongati, tibiis incuruis.

33. R. femoribus dentatis, brunneo cinereoque variegatus, thorace tuberculis quatuor, elytris duobus baſeos. *6 tuberculatus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Minor. Roſtrum inflexum, atrum. Thorax variegatus, tuberculis quatuor eleuatis, dorſalibus, obtuſis. Elytra concolora, ſtriata: ſtriis ſubtiliſſime-ſerratis, tuberculisque duobus baſeos eleuatis. Femora clauata, ferruginea, annulo atro.

34. R. femoribus dentatis, ferrugineus, elytris ſtriatis: puncto medio albo. *verruca.*

Habitat in America meridionali. D. Smidt. Muſ. D Lund.

Statura R. abietis. Roſtrum obſcurum, puncto baſeos obſoleto, albo. Thorax ſcaber, tuberculis duobus paruis eleuatis. Elytra ſtriata: ſtriis ſubſerratis, obſcure-brunnea: puncto paruo, medio, albo.

35. R. femoribus dentatis, thorace albo, elytris nigris, alboſtriatis, tibiis dentatis. *dentipes.*

*Curculio dentipes.* Ent. ſyſt. 2. 428. 145. *

Oliv. Inſ. 83. tab. 8. fig. 90.

Habitat in Senegalliae Goſſypio. Muſ. Gallicum.

36. R. femoribus dentatis, niger, thorace elytrisque albo-punctatis. *multiguttatus.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio multiguttatus*. Ent. ſyſt. 2. 428. 146.
Oliv. Inſ. 83. tab. 13. fig. 163.
Habitat in America meridionali.

*pupillator.* 137. R. femoribus dentatis, tuberculatis, fuſcus, elytris macula marginali cinerea, nigro-punctata.
*Curculio pupillator*. Ent. ſyſt. 2. 428. 147.
Oliv. Inſ. 83. tab. 15. fig. 183.
Habitat in Cayenna. Muſ. Dom. Geoffroy.

*Nero.* 138. R. femoribus dentatis, niger, fuluo-maculatus, pedibus anticis elongatis.
Habitat in Sumatra. D. Daldorff.
Roſtrum cylindricum, nigrum. Thorax punctis eleuatis ſcaber, obſcurus, lineis duabus fuluis. Elytra punctato-ſtriata, nigra: linea baſeos faſcia media maculaque apicis fuluis. Pedes rufi, antici valde elongati, femoribus poſticis ſubbidentatis.

*porcatus.* 139. R. femoribus dentatis, cinereus, thoracis elytrorumque dorſo nigris, pedibus anticis poſticisque elongatis.
Habitat in Sumatra. D. Daldorff.
Statura omnino praecedentis. Caput cinereum, roſtro cylindrico, nigro. Thorax obſcure-cinereus, dorſo nigro. Elytra ſulcata: ſulcis punctatis, cinereis, a baſi ad medium ad futuram nigris. Verſus apicem tuberculum paruum, eleuatum. Pedes antici poſticique elongati, femoribus clauatis.

*Lapathi.* 140. R. femoribus bidentatis, albo nigroque variis, thorace elytrisque muricatis.
*Curculio Lapathi*. Ent. ſyſt. 2. 429. 149.
Linn. ſyſt. Nat. 2. 608. 20. Fn. Sv. 591.
Oliv. Inſ. 83. tab. 6. fig. 96.
Panz. Fn. Germ. 42. tab. 15.

Dege

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 5. 223. 16. tab. 7. fig. 1. 2.
Payk. Monogr. 37. 35.
Habitat in Europae Salice.

41. R. femoribus dentatis, obſcurus, elytris lineolis eleuatis, apice atro-brunneis. *abdominalis.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo R Lapathi. Roſtrum arcuatum, atrum, apice ferrugineum. Thorax rugoſus, niger, obſcurus. Elytra nigra, lineolis breuibus, eleuatis, ante apicem atra, apice brunnea. Corpus nigrum, ano brunneo. Pedes cinereo-variegati.

42. R. femoribus dentatis, obſcurus, albo-ſubmaculatus, elytris lineis eleuatis, plurimis, compreſſis. *argula.*
Habitat in Carolina. Muſ. Dom. Boſc.
Statura et magnitudo R. Lapathi. Roſtrum incuruum, fuſcum. Thorax antice vtrinque impreſſus, obſcurus, lineis duabus, vndatis, albidis. Elytra obſcura, punctis aliquot baſeos albis maculaque pone medium communi brunnea, lineolisque plurimis breuibus, eleuatis, carinatis. Pedes antici elongati, femoribus bidentatis.

43. R. femoribus dentatis, albus: ſupra fuſcus, albo-maculatus, femoribus albo-annulatis. *irroratus.*
*Curculio irroratus.* Ent. ſyſt. 2. 429. 150.
Habitat in Cayenna. Dom. v. Rohr.

44. R. femoribus dentatis, griſeus, elytris ſtriis tribus eleuatis: exterioribus abbreuiatis. *porculus.*
*Curculio porcatus.* Panz. Fn. Germ. V. 17.
Habitat in Auſtria Dom. de Meyerle.
Magnitudo omnino R. Lapathi, at magis depreſſus. Caput nigrum, roſtro craſſiuſculo, vtrinque ſul-



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

cato. Thorax ſubuarioloſus, griſeus. Elytra griſea: lineis tribus eleuatis: exterioribus ante apicem coëuntibus.

*pilipes.* 145. R. femoribus dentatis, cinereus, pedibus anticis elongatis: tibiis tarſisque ciliato-piloſis.
Habitat in Iana. Muſ. Dom. Boſc.
Paullo maior R. ſtatua. Caput cinereum, roſtro cylindrico, incuruo, nigro, apice ſubbiſerrato. Thorax ſubpunctatus, cinereus. Elytra ſtriata, cinerea, immaculata. Pedes antici elongati, femoribus in medio vnidentatis, tibiis tarſisque pilis denſis, elongatis, ſubtus ciliatis, intermedii breues, poſtici elongati, femoribus clauatis.

*ſtatua.* 146. R. femoribus dentatis, ater, elytris ſulcatis macula communi alba, femoribus poſticis elongatis.
*Curculio ſtatua.* Ent. ſyſt. 2. 430. 152.
Habitat - - - Muſ. Dom. Boſc.

*tricinctus.* 147. R. femoribus dentatis, ater, thoracis lateribus elytrorumque faſciis tribus albis, pedibus anticis elongatis.
*Curculio tricinctus.* Ent. ſyſt. 2. 430. 151.
Oliv. Inſ. 83. tab. 6. fig. 61.
Habitat in Inſula Guadeloupe. Dom. Badier.

7 *tuberculatus.* 148. R. femoribus dentatis, thorace quadritubercula-to, elytris ſubſpinoſis: tuberculis tribus.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Medius. Caput cinereum. Thorax cinereus, in medio tuberculis quatuor eleuatis. Elytra ſtriata, cinerea: ſtriis ſpinis paruis, acutis, nigris, ſubſerrata. Praeterea tubercula tria eleuata obtuſa ad ſuturam. Corpus cinereum.

149

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

49. R. femoribus dentatis, thorace ferrugineo: ocellis duobus, elytris griſeis, apice brunneis. *coccus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Medius. Caput obſcurum, roſtro nigro. Thorax villoſus, ferrugineus: maculis duabus ocellaribus, griſeis. Elytra eleuato-ſtriata, griſea, apice brunnea. Corpus brunneum.

50. R. femoribus dentatis, obſcure-cinereus, elytris lineis eleuatis, compreſſo-carinatis. *Coluber.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Medius et diſtinctus, lineolis elytrorum plurimis, eleuatis, valde compreſſis et fere carinatis.

51. R. femoribus dentatis, fuſcus elytris lineis eleuatis, compreſſis, ſubſpinoſis. *Aſpis.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et affinitas R. Colubri, at diſtinctus. Roſtrum tenue, inflexum, nigrum. Thorax gibbus, ſcaber, fuſcus: puncto baſeos albo. Elytra fuſca, lineis plurimis eleuatis, compreſſo-carinatis, ſubſpinoſis. Corpus fuſcum, pectoris lateribus albidis.

152. R. femoribus dentatis, fuſcus, thorace tuberculo medio ſubbifido, elytris ſtriatis tuberculatisque. *Ceraſtes.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Corpus medium. Roſtrum tenue, arcuatum, inflexum. Antennae ſubferrugineae, claua ouata. Thorax ſcaber, fuſcus, tuberculo eleuato, ſubbifido in medio. Elytra ſtriata, ſulcis punctatis tuberculisque aliquot eleuatis, compreſſis. Femora clauata.

153.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*haemorrhoes.* 153. R. femoribus dentatis, fuſcus, elytris coſtatis, femoribus poſticis bidentatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura praecedentium. Roſtrum inflexum, atrum. Thorax ſcaber, fuſcus, linea dorſali eleuata. Elytra magis brunnea, ſulcata: ſulcis quatuor et in his ſtriae duae e punctis impreſſis. Femora poſtica bidentata.

*6guttatus.* 154. R. femoribus dentatis, niger, elytris maculis tribus albis.
*Curculio 6guttatus.* Ent. ſyſt. 2. 430. 153. *
Oliv. Inſ. 83. tab. 14. fig. 170.
Drury Inſ. 2. tab. 33. fig. 1. 2.
Habitat in America. Muſ. D. Hunter.

*luridus.* 155. R. femoribus dentatis, ouatus, obſcure-niger, elytris punctato-ſtriatis.
*Curculio luridus.* Ent. ſyſt. 2. 431. 154. *
Oliv. Inſ. 83. fig. 175.
Habitat in noua Hollandia. Muſ. D. Banks.

*ſtolidus.* 156. R. femoribus dentatis, fuſcus, tibiis poſticis incuruis, dentatis.
*Curculio ſtolidus.* Ent. ſyſt. 2. 431. 155. *
Oliv. Inſ. 83. fig. 145.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*mixtus.* 157. R. fuſcus, cinereo-varius, elytris ſtriatis: lituris duabus lateralibus, ferrugineis.
Habitat in Amboina. Dom. Billardiere.
Corpus cylindricum, colore cinereo fuſcoque mixto. Thorax vix punctatus. Elytra ſtriata, lituris duabus ante medium fere marginalibus, ferrugineis. Femora acute-dentata.

*frigidus.* 158. R. femoribus dentatis, fuſcus, elytris ſtriatis, ſubtuberculatis, ferrugineo-variis,

Cr.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio frigidus.* Ent. ſyſt. 2. 431. 156. *
*Curculio Chinenſis.* Oliv. Inſ. 83. tab. 8. fig. 97.
Habitat in Amboina. Dom. de Paykull.

159. R. femoribus dentatis, ſupra fuſcus, thorace elytrisque albo-ſtriatis. ***pictus.***
*Curculio pictus* Ent. ſyſt. ſuppl. 166. 157. *
Habitat in Tanger. Dom. Schousboe. Muſ. D. de Seheſtedt.

160. R. femoribus dentatis, obſcurus, thorace elytrisque faſciculato-piloſis. ***piloſus.***
Habitat in Iaua. Muſ. Dom. Boſc.
Statura omnino R. frigidi, at paullo minor. Caput nigrum. Thorax punctatus, obſcurus: faſciculis piloſis, antice duobus, in medio quatuor. Elytra ſtriata, faſciculis pilorum plurimis, ſparſis. Pedes obſcuri.

161. R. femoribus dentatis, cinereus, roſtro atro, elytris ſulcato-punctatis. ***alienatus.***
Habitat in Sumatra. D. Daldorff.
Minor R. frigido. Roſtrum arcuatum, inflexum, atrum. Thorax vix punctatus, obſcure-cinereus, immaculatus. Elytra ſulcata, ſulcis punctatis. Pedes cinerei, femoribus dentatis.

162. R. femoribus dentatis, niger, elytris ſtriatis, ferrugineis: lineis poſticis duabus albidis. ***Fabricii.***
*Curculio Fabricii.* Ent. ſyſt. ſuppl. 166. 157. *
Habitat Tranquebariae. D. Daldorff.

163. R. femoribus dentatis, niger, elytris excauato-punctatis: ſtriga abbreuiata, albida. ***gibbus.***
*Curculio gibbus.* Ent. ſyſt. 2. 431. 157. *
Oliv. Inſ. 83. tab. 8. fig. 88.
Habitat in Inſula Bourbon. Muſ. Gallicum.

164. R. femoribus dentatis, ater, nitidus, immaculatus. ***globus.***

Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura breuis praecedentium, totus laeuis, ater, nitidus, elytra vix ſtriata.

*Ebenus.* 165. R. femoribus dentatis, ater, nitidus, elytris tuberculatis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Affinis praecedenti, at paullo minor et alius, totus ater, nitidus, immaculatus. Thorax laeuiſſimus, nitidiſſimus. Elytra nitida, tuberculis plurimis obſoletis inaequalia.

*calidus.* 166. R. obſcure-aeneus, roſtro atro, thorace rugoſo, elytris punctato-ſtriatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Medius. Roſtrum incuruum, ſtriatum, atrum. Thorax rugoſus, obſcure-aeneus. Elytra punctato-ſtriata, aenea. Corpus aeneum.

*flauicornis* 167. R. femoribus dentatis, thorace elytrisque tuberculatis, niger, elytris cinereo-faſciatis, antennis flauis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Roſtrum incuruum, nigrum, antennis flaueſcentibus. Thorax tuberculatus, niger. Elytra ſtriata, tuberculata, nigra, faſciis aliquot obſoletis nigris. Pedes cinerei, femoribus anticis incraſſatis.

*ocellatus.* 168. R. femoribus dentatis, obſcure-cinereus, elytris macula ocellari atra.

*Curculio ocellatus.* Ent. ſyſt. 2. 432. 158. Oliv. Inſ 83. fig. 31.

Habitat in Cayenna.

16

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

169. R. femoribus dentatis, elytris ſtriatis, poſtice acuminatis. *meditabundus.*

*Curculio meditabundus.* Ent. ſyſt. 2. 432. 159.*
Oliv. Inſ. 83. fig. 132.
Habitat in noua Hollandia. Muſ. D. Banks.

170. R. femoribus dentatis, fuſcus, elytris reticulato-ſtriatis, poſtice tuberculatis, apice acuminatis. *Pacca.*

Habitat in India orientali. Muſ. D. Lund.
Roſtrum craſſiuſculum, incuruum. Thorax punctis eleuatis ſcaber. Elytra ſulcata: ſulcis ſtriis transuerſis, reticulatis, tuberculoque ante apicem eleuato, obtuſo, apiceque acumine paruo, fuſca, punctis faſciculatis, brunneis, ſparſis. Corpus fuſcum, pedibus pilis ferrugineis, maculatis.

171. R. femoribus dentatis, obſcurus, thoracis linea laterali ſcutello elytrorumque faſcia cinereis. *Caprea.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura omnino R. Pacca. Roſtrum longitudine thoracis. Thorax ſcaber, obſcurus: linea laterali cinerea. Elytra ſulcata: ſulcis rugoſis, obſcura. Pedes ferruginei, femoribus acute-dentatis.

172. R. femoribus dentatis, niger, thoracis lateribus rotundatis, elytris ſubſpinoſis. *ſtrepidus.*

*Curculio ſtrepidus.* Ent. ſyſt. 2. 432. 160.*
Oliv. Inſ. 83. fig. 151.
Habitat in noua Hollandia. Muſ. D. Banks.

173. R. femoribus dentatis, obſcurus, thorace ſcabro: linea dorſali alba, elytris reticulatis. *Mangiferae.*

*Curculio Mangiferae.* Ent. ſyſt. 2. 432. 161.*
Oliv. Inſ. 83. fig. 137.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Hybn. Naturf. 24. 46. 12. tab. 2. fig. 13.
Habitat in Mangiferae nucleis Indiae orientalis. Koenig.

*ſtultus.* 174. R. femoribus dentatis, griſeus, elytris macula communi lunata, cinerea, roſtro atro.
*Curculio ſtultus.* Ent. ſyſt. 2. 433. 162. *
Habitat in Coromandel. Prof. Vahl.

*Araneus.* 175. R. femoribus dentatis, obſcurus, cinereo-virius, thoracis lateribus rotundatis.
*Curculio Araneus.* Ent. ſyſt. 2. 433. 163. *
Habitat in Americae Inſulis. Dr. Pflug.

*Bubo.* 176. R. femoribus dentatis, thorace cinereo, villoſo, elytris punctato-ſtriatis, brunneis: litura poſtica cinerea.
Habitat - - - Muſ. D. Lund.
Corpus praecedentibus magis cylindricum. Roſtrum craſſiuſculum, brunneum. Thorax cinereo-villoſus, dorſo fuſceſcente: linea cinerea. Punctum fuſcum praeterea ad latera. Elytra punctis latioribus ſtriata, brunnea, litura transuerſa, villoſo-cinerea. Corpus cinereo-villoſum. Pedes brunnei.

*Strix.* 177. R. femoribus dentatis, niger, thorace ferrugineo, lineato, elytris ſtriatis, ferrugineo-punctatis.
*Curculio Strix.* Ent. ſyſt. 2. 433. 164. *
Oliv. Inſ. 83. fig. 16.
Habitat in Cayenna. Muſ. Dom. Olivier.

*ſulcatulus.* 178. R. femoribus dentatis, ater, elytris ſulcis tribus villoſo-albis.
Habitat in Amboina. Dom. Billardiere.
Statura cylindrica R. ſqualidi. Caput atrum. Thorax trilobus, punctis paruis, eleuatis, ſcaber: ſtrigis duabus in medio interruptis, impreſ-

ſis.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſis, villoſo-albis. Elytra punctato-ſtriata, et praeterea ſulcis tribus villoſo-albis exarata, qui tamen apicem haud attingunt. Corpus atrum, femoribus acute-dentatis.

79. R. femoribus dentatis, villoſo-griſeus, roſtro teſtaceo, elytris ſtriis tribus eleuatis, albidis. *ſqualidus.*

*Curculio ſqualidus.* Ent. ſyſt. 2. 433. 165. *
Oliv. Inſ. 83. fig. 184.
Habitat Surinami. Muſ. D. Banks.

80. R. femoribus dentatis, ater, thorace punctis duobus teſtaceis. *germanus.*

*Curculio germanus.* Ent. ſyſt. 2. 433. 166.
Linn. ſyſt. Nat. 2. 613. 58.
Panz. Fn. Germ. 42. tab. 16.
Sulz. Hiſt. Inſ. tab. 4. fig. 5.
Schaeff. Icon. tab. 62. fig. 9.
tab. 101. fig. 6.
Friſh. Inſ. 13. 28. tab. 26.
Payk. Monogr. 1. 1.
Habitat in Germania haud infrequens.

IV. 72

81. R. femoribus dentatis, fuſco albo ferrugineoque variegatus pectore biſpinoſo. *biſpinoſus.*

Habitat in Sumatra. Dom. Daldorff.
Forte cum ſequentibus proprii generis, Cionus Ent. Helv. Differt habitu, corpore magis depreſſo, planiuſculo, antennarum tertio quartoque articulis longioribus clauaque ouata, at inſtrumenta cibaria mihi haud rite nota. Cionus Lythri vero Ent. Helv. tab. 3. fig. 3. nec ſtatura nec antennis huc pertinet.
Maior R. Thapſi. Roſtrum elongatum, nigrum, baſi parum teſtaceum. Oculi magni, approximati. Thorax et elytra variegata, ſcutello albo. Corpus albo-villoſum, pectore antice ſpinis dua-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

duabus validis, acutis. Pedes elongati, atri, cinereo-irrorati.

Femina mutica.

*ſanctus.* 182. R. femoribus dentatis, ater, thorace vtrinque albo-cruciato, elytris cinereo-punctatis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Maior R. Scrophulariae. Roſtrum incurvum, baſi ſulcatum. Orbita oculorum albida. Thorax ater, linea dorſali et vtrinque cruce albis. Elytra atra, punctis numeroſis, ſparſis, albis. Pectus cinereo-villoſum, atro-maculatum. Abdomen atrum, lineis quatuor albis. Femora poſtica compreſſa.

*deplanus.* 183. R. femoribus dentatis, ater, thorace albo-lineato, elytris acuminatis: ſutura ſtrigisque albis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura praecedentis. Roſtrum arcuatum, atrum, baſi compreſſum, cinereum. Thorax ſubrugoſus, ater: lineis quinque integris, albis. Elytra ſtriata, apice parum acuminata, atra: ſutura ſtrigisque aliquot obliquis, albidis. Subtus villoſo-cinereus, abdomine linea media atra.

*Phleſus.* 184. R. femoribus dentatis, elytris baſi bituberculatis, albo-maculatis, maculaque magna ſuturali atra.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura plana, depreſſa R. Scrophulariae, at duplo maior. Roſtrum atrum, nitidum, baſi albidum. Thorax nigricans interdum punctis aliquot albis, interdum antice ferrugineus. Scutellum ferrugineum. Elytra tuberculis duobus baſeos eleuatis, obtuſis, ſubſtriata, nigro albo-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

que variegata maculaque magna media, communi, atra. Elytra apice emarginata. Subtus nunc cinereo, nunc ferrugineo-villoſus, femora elongata, clauata, dentata.

85. R. femoribus dentatis, obſcure-ferrugineus, elytris albo-punctatis, baſi bituberculatis, apice emarginatis. *Pleuronectes.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura praecedentis, paullo minor. Roſtrum elongatum, incuruum, atrum. Caput et thorax obſcure-brunnea, interdum nigro-maculata. Elytra punctato-ſtriata, brunnea, punctis paruis, albis, ſparſis, baſi tuberculis duobus eleuatis, apice emarginatis. Femora valde elongata, clauata.

186. R. femoribus dentatis, poſticis apice ſpinoſis, fuſcus, albo-punctatus. *planulus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Praecedentibus ſtatura plana, depreſſa conuenit. Roſtrum incuruum, atrum. Caput fuſcum, orbita oculorum alba. Thorax fuſcus: lateribus nigris, albo-punctatis. Elytra ſubſtriata, apice rotundata, fuſca, punctis plurimis ſparſis, albis. Pedes nigricantes, femoribus poſticis elongatis, clauatis, tridentatis apiceque ſpina valida, acuta, annuloque albo-cincta.

187. R. femoribus dentatis, niger, thorace rugoſo: linea laterali alba, coleoptris punctato-ſtriatis: maculis quatuor albis, duabus communibus. *lamella.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Roſtrum atrum, baſi parum carinatum. Thorax rugoſus, ater, linea laterali lata, alba. Elytra ro-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

rotundata, nigra: maculis tribus albis; duabus communibus. Pectus albo-maculatum. Pedes nigri, femoribus elongatis, tenuibus.

*temporarius.* 188. R. femoribus dentatis, niger, elytris fuscis margine nigro, albo-punctato.

Habitat in America meridionali. D. Smidt. Mus. D. de Seheſtedt.

Statura omnino praecedentium. Roſtrum arcuatum, incuruum, atrum. Caput et thorax laeuia, atra, orbita, oculorum thoracisque lateribus albis. Elytra laeuia, fuſca, baſi parum nigro-alba, lateribus nigris, albo-punctatis. Corpus cinereo-villoſum, nigro-maculatum. Pedes nigri, femoribus tridentatis: dentibus duobus poſticis minutis.

*Pipa.* 189. R. femoribus dentatis, griſeus, elytris ſtriatis nigro ferrugineoque variis.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura praecedentis. Roſtrum incuruum, atrum, baſi biſulcatum. Orbita oculorum albida. Thorax et elytra nigro ferrugineoque varia. Corpus griſeum.

*Vlula.* 190. R. femoribus dentatis, cinereus, thoracis macula media nigra: linea albida, elytris ſtriatis: faſcia nigra.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura omnino praecedentium. Roſtrum elongatum, arcuatum, atrum. Thorax cinereus, macula magna nigra, et in hac linea media cinerea. Elytra ſtriata cinerea: faſcia lata media nigra.

*Scrophulariae.* 191. R. femoribus dentatis, thorace albido, coleoptris maculis duabus atris, albae connatis.

*Cur-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Scrophulariae.* Ent. ſyſt. 2. 434. 167.
Linn. ſyſt. Nat. 2. 614. 61. Fn. Sv. 603.
Geoff. Inſ. 1. 296. 44.
Reaum. Inſ. 3. tab. 2. fig. 12.
Degeer Inſ. 5. 208. 3. tab. 6. fig. 17. 18. 19. 20.
Payk. Monogr. 23. 22.
- Fn. Sv. 3. 207. 24.
Herbſt. Col. 6. tab. 73. fig. 1.
Habitat in Europae Scrophulariis.

192. R. femoribus dentatis, coleoptris maculis duabus dorſalibus ſimplicibus atris. *Thapſus.*
*Curculio Thapſus* Ent. ſyſt. 2. 434. 168. *
Oliv. Inſ. 83. fig. 21.
Herbſt. Col. 6. tab. 73. fig. 2.
Habitat in Scrophulariis, Verbaſco.
*Varietatem* praecedentis credit D. de Paykull rite?

193. R. femoribus dentatis, niger, thoracis lateribus flaueſcentibus, elytris punctis albis nigrisque, alternis ſtriatis. *Verbaſci.*
*Curculio Verbaſci.* Ent. ſyſt. 2. 434. 169.
Habitat Kiliae. D. Daldorff.

194. R. femoribus dentatis, niger, thorace tuberculato, elytrorum futura macula atra albaque. *moeſtus.*
Habitat in Europa. Muſ. D. de Seheſtedt.
Statura et magnitudo praecedentium. Caput fuſcum, roſtro nigro. Thorax tuberculo vtrinque eleuato, niger, interdum puncto baſeos albido. Elytra ſtriata, nigra, futura puncto baſeos eleuato atro, apicis albo. Corpus cinereum, puncto paruo villoſo ferrugineo, ad baſin thoracis.

195. R. femoribus dentatis, albidus, coleoptris nigro variis: macula dorſali baſeos apicisque nigris. *Blattariae*
*Curculio Blattariae.* Ent. ſyſt. 2. 435. 170. *

*Cio-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Cionus Blattariae.* Ent. Heluet. 1. 66. 1. tab. 3 fig. 1.
Habitat in Italia. Dr. Allioni.

*olens.* 196. R. femoribus dentatis, cinereo-piloſus, coleo ptris macula dorſali atra.
*Curculio olens.* Ent. ſyſt. 2. 435. 171. ♀
Habitat in Gallia. Muſ. D. Boſc.

*caprimulgus.* 197. R. femoribus dentatis, villoſus, cinereus, an tennis pedibusque teſtaceis.
Habitat in Germania. D. Smidt.
Statura omnino et magnitudo R. olentis, at totu cinereus, immaculatus, antennis pedibusque te ſtaceis.

*foetidus.* 198. R. femoribus dentatis, elytris cinereis: macu la dorſali fuſca ſtriisque abbreuiatis punctorum nigrorum.
Habitat in Selandia Daniae. Muſ. D. de Seheſted
Statura praecedentium. Roſtrum baſi rufum, api ce nigrum. Thorax cinereus: macula dorſa nigra; linea media cinerea. Elytra cinere macula magna communi baſeos fuſca, lineisqu aliquot indiſtinctis abbreuiatis e punctis albi nigrisque.
Mas paullo minor magisque obſcurus.

*Solani.* 199. R. femoribus dentatis, obſcurus, elytris linei eleuatis nigris cinereo-punctatis.
*Curculio Solani.* Ent. ſyſt. 2. 435. 171. ♀
Habitat Halae Saxonum. Dom Hybner.

*Smidtii.* 200. R. femoribus dentatis, thorace carinato, dimi diato-rufo, elytris planis biſpinoſis fuſcis.
Habitat in America meridionali. D. Smidt. Mu D. Lund.
Statura praecedentium. Roſtrum arcuatum, cine reum, dorſo atro. Thorax valde carinatus, an tic

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tice rufus, poſtice niger. Elytra plana, angulo baſeos anteque apicem acute ſpinoſa, fuſca, baſi nigra. Corpus atrum, abdomine baſi albo-maculato. Pedes tenues, griſei, dorſo nigro.

1. R. femoribus dentatis, nigro-faſciatis, thorace elytrisque tuberculatis, femoribus anticis craſſiſſimis. *chiragra.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentium. Roſtrum longum, arcuatum, atrum. Thorax cinereus, nigro-faſciatus, tuberculis octo: 4. 4; duobus intermediis lineae anticae maioribus. Elytra plana, tuberculis plurimis, ſparſis apicis maiori obtuſo, cinerea, atro-faſciata. Pedes atri, femoribus anticis longioribus, craſſiſſimis, dentatis, tibiis incuruis.

2. R. femoribus dentatis, elytris tuberculo ſuturali eleuato anguloque apicis dilatato, plano. *Aeſopus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paruus. Roſtrum longiuſculum, cylindricum, ferrugineum, apice atrum. Thorax laeuis, ferrugineus, carina media eleuata, atra. Elytra ſtriata, ferruginea: angulo baſeos prominulo. In medio tuberculum, ad ſuturam eleuatum, acutum, at angulus apicis dilatatus, planus. Pedes ferruginei, femoribus anticis longioribus, incraſſatis, dentatis.

3. R. femoribus dentatis, niger, elytris ferrugineo-variis, femoribus canaliculatis. *grauis.*

*Curculio grauis.* Ent. ſyſt. 2. 435. 172. *

Oliv. Inſ. 83. fig. 177.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſpunctatus* 204. R. femoribus dentatis, elytris ſutura puncti que duobus albis, femoribus canaliculatis.
*Curculio ſpunctatus.* Ent. ſyſt. 2. 435. 173.
Linn. ſyſt. Nat. 2. 614. 64. Fn. Sv. 618.
Schaeff. Icon. tab. 1. fig. 12.
Herbſt. Arch. tab. 24. fig. 18.
Coleopt. 6. tab. 81. fig. 3.
Payk. Monogr. 15. 14.
Fn. Sv. 3. 201. 18.
Habitat in Europae plantis.

*guttula.* 205. R. femoribus dentatis, thorace tuberculato, ger, elytris ſtriatis: puncto poſtico albo.
*Curculio guttula.* Ent. ſyſt. 2. 436. 174. *
Habitat Halae Saxonum. Dom. Hybner.

*abbreuiatulus.* 206. R. femoribus dentatis, fuſcus, atomis griſei irroratis, elytris abbreuiatis.
*Curculio abbreuiatulus.* Ent. ſyſt. 2. 436. 175.
Habitat Halae Saxonum. Dom. Hybner.

*Echii.* 207. R. femoribus dentatis, niger, thorace elytr que albo-lineatis.
*Curculio Echii.* Ent. ſyſt. 2. 436. 176. *
*Curculio geographicus.* Oliv. Inſ. 83. fig. 56.
Panz. Fn. Germ. 17. tab. 12.
Habitat in Germaniae Echio. Dom. Prof. Helwi

*ſanguini-roſtris.* 208. R. femoribus dentatis, viridi-cinereus, vil fus, roſtro rufo, baſi albo.
Habitat in America meridionali. D. Smidt. Mu D. de Seheſtedt.
Paullo maior R. didymo, totus viridi-cinere villoſus. Roſtrum arcuatum, baſi atrum, ap rufum. Antennae cinereae, claua rufa. Ely ſtriata.

*didymus.* 209. R. femoribus dentatis, ſupra fuſcus, ely ſtriatis: macula laterali transuerſa, alba.

C

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio didymus.* Ent. ſyſt. 2 437. 177.
*Curculio vrticarius.* Ent. Helv. 1. 98. 11. tab. 11. fig. 1. 2.
Geoff. Inſ. 1. 299. 47. *IV. 74.*
Payk. Monogr. 29. 28.
Fn. Sv. 3. 213. 30.
Herbſt Col. 6. tab. 91 fig. 6.
Habitat in Germaniae Vrtica vrente. D. de Hattorff.

210. R. femoribus dentatis, niger, thorace tuberculato, elytris cinereo-variis, roſtro arcuato, atro. *Lamii.*
*Curculio Lami* Ent. ſyſt. 2. 437. 178. * *E. 57.*
Panz. Fn. Germ. 17. tab. 13.
Habitat in Germaniae Lamio. Dom. Prof. Helwig.

211. R. femoribus dentatis, cinereo-villoſus, roſtro arcuato, atro. *Braſſicae.*
*Curculio Braſſicae.* Ent. ſyſt. 2. 437. 179. *
Habitat in Kiliae Braſſica.

212. R. femoribus dentatis, cinereo-villoſus, tarſis rufis. *Borraginis.*
*Curculio Borraginis.* Oliv. Inſ. 83. fig.
Habitat in Galliae Borragine. Dom. Olivier.

213. R. femoribus dentatis, niger, roſtro pedibusque ferrugineis: femoribus anticis incraſſatis. *rufiroſtris.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Paruus. Roſtrum planum, flauum. Thorax niger. Elytra ſtriata, nigra. Pedes flaui: antici elongati, femoribus incraſſatis.

214. R. femoribus dentatis, cinereus, thorace tuberculis duobus, elytris plurimis eleuatis. *multituberculatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paruus. Roſtrum obſcurum. Thorax cinereus, tuberculis duobus. Elytra cinerea, tuberculis plurimis: interiori maiore, compreſſo. Corpus nigrum.

*haemorrhoidalis.* 215. R. femoribus dentatis, thorace fuſco: lateribus cineraſcentibus, elytrorum futura apice ferruginea.

*Curculio haemorrhoidalis.* Ent. ſyſt. 2. 437. 181. *

Habitat in noua Hollandia. Muſ. Dom. Banks.

*trimaculatus.* 216. R. femoribus dentatis, coleoptris nigris: maculis tribus cinereis; poſtica communi lunata.

*Curculio 3 maculatus.* Ent. ſyſt. 2. 438. 182.

Payk. Monogr. 26. 24.

Fn. Sv. 3. 209. 26.

Herbſt. Col. 6. 471. 482.

Habitat in Alſatia. Prof. Hermann.

*litura.* 217. R. femoribus dentatis, albo nigroque variis, roſtro atro.

*Curculio litura.* Ent. ſyſt. 2. 438. 183. *

Payk. Monogr. 26. 25.

Fn. Sv. 3. 210. 27.

Herbſt. Col. 6. tab. 91. fig. 4.

Habitat in Europae borealis Carduis.

*villoſus.* 218. R. femoribus dentatis, villoſus, griſeus, ſcutello faſciaque elytrorum poſtice albis.

*Curculio villoſus.* Ent. ſyſt. 2. 438. 184. *

Herbſt. Arch. tab. 24. fig. 19.

Habitat in Germania. Dom. de Hattorff.

*Sciurus.* 219. R. femoribus dentatis, niger, elytris ſtriatis punctis duobus ſuturalibus, albis.

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Pa

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paruus. Caput et thorax laeuia, atra, immaculata. Elytra ſtriata, atra: punctis duobus communibus, albis.

220. R. femoribus dentatis, ater, elytris ſtriatis: plaga rufa. *plagiatus.*

*Curculio plagiatus.* Ent. ſyſt. ſuppl. 166. 184. *
Habitat in Carolina. Muſ. D. Boſc.

221. R. femoribus dentatis, nigricans, thorace tuberculato, roſtro arcuato, atro. *Raphani.*

*Curculio Raphani.* Ent. ſyſt. 2. 438. 185. *
Habitat in Germaniae Raphano. D. Smidt.

222. R. femoribus dentatis, niger, elytris rufis: margine omni nigro. *varians.*

*Curculio varians.* Ent. ſyſt. 2. 439. 186. *
Payk. Monogr. 16. 15.
Fn. Sv. 3. 202. 19.
Herbſt. Coleopt. 6. tab. 80. fig. 8.
Habitat in Sueciae Pino. Dom. de Paykull.

223. R. femoribus dentatis, fuſcus, thorace linea dorſali cinerea, elytris pedibusque teſtaceis. *Troglodytes.*

*Curculio Troglodytes.* Ent. ſyſt. 2. 439. 187. *
Payk. Monogr. 30. 29.
Fn. Sv. 3. 214. 31.
Herbſt. Col. 6. 473. 484.
Habitat Kiliae. D. Daldorff.

224. R. femoribus dentatis, oblongus, ater, antennarum claua villoſa, elytris crenato-ſtriatis. *carbonarius.*

*Curculio carbonarius.* Ent. ſyſt. 2. 439. 188.
Linn. ſyſt. Nat. 2. 617. 42. Fn. Sv. 614.
Panz. Fn. Germ. 42. tab. 18.
Payk. Fn. Sv. 3. 194. 11.
Herbſt. Col. 6. tab. 64. fig. 3.
Habitat in Europa.

IV. 72



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*aterrimus.* 225. R. femoribus dentatis, ater, thorace apice bidentato, elytris nitidis.
*Curculio aterrimus* Ent. ſyſt. 2. 439. 189.
Linn. ſyſt. Nat. 2. 607. 10. Fn. Sv. 582.
Habitat in Europa boreali.

*Ceraſi.* 226. R. femoribus dentatis, ater, thorace apice bidentato, elytris oblongis opacis.
*Curculio Ceraſi.* Ent. ſyſt. 2. 440. 190.
Linn. ſyſt. Nat. 2. 607. 11. Fn. Sv. 583.
Ent. Heluet. 1. 90. 7. tab. 9. fig. 1. 2.
*W. 72.* Panz. Fn. Germ. 42. tab. 19.
Geoff. Inſ. 1. 299. 48.
Friſh. Inſ. 11. 31. tab. 23. fig. 1. 3.
Payk. Monogr. 8. 7.
Fn. Sv. 3. 193. 10.
Herbſt. Col. 6. tab. 64. fig. 1. 2.
Habitat in Ceraſi foliis.

*violaceus.* 227. R. femoribus dentatis, niger, elytris ſtriis violaceis.
*Curculio violaceus.* Ent. ſyſt. 2. 440. 191.
Linn. ſyſt. Nat. 2. 614. 63. Fn. Sv. 579.
Geoff. Inſ. 1. 271. 5.
Degeer Inſ. 5. 213. 5.
*W. 73* Panz. Fn. Germ. 42. tab. 20.
Payk. Monogr. 9. 8.
Fn. Sv. 3. 195. 12.
Herbſt. Col. 6. tab. 64. fig. 4.
Habitat in Europae borealis Pinis.

*Nucum.* 228. R. femoribus dentatis, corpore griſeo, longitudine roſtri.
*Curculio Nucum.* Ent. ſyſt. 2. 440. 192.
Linn. ſyſt. Nat. 2. 673. 59. Fn. Sv. 616.
Roſſ. Inſ. 3. tab. 67. fig. 67.
*W. 73* Oliv. Inſ. 83. fig. 47.
Panz. Fn. Germ. 42. tab. 21.

Schaf-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Schaeff. Elem. tab. 55. fig. 1.
Icon. tab. 50. fig. 4.
Degeer Inſ. 5. 205. 2. tab. 6. fig. 14. 15. 16.
Geoff. Inſ. 1. 295. 42.
Sulz. Hiſt. Inſ. tab. 3. fig. 22.
Payk. Monogr. 21. 20.
Fn. Sv. 3. 204. 20.
Herbſt. Col. 6. tab. 73. fig. 10.
Habitat in Coryli Auellanae nucleis.
Forte cum ſequentibus proprii generis. Differt parum ſtatura, at roſtrum longiſſimum, tenuiſſimum, cylindricum, parum incuruum, et antennae articulo ſecundo longiſſimo, vsque ad oculos attingente, clauato, tertio ſequentibus longiore, vltimis tribus craſſioribus, claua obtuſa, in medio roſtri inſertae, proprium genus indicare videntur.

29. R. femoribus dentatis, griſeus, roſtro corpore duplo longiore. *proboſcideus.*
*Curculio proboſcideus.* Ent. ſyſt. 2. 440. 193. *
Oliv. Inſ. 83. fig. 127.
Habitat in America boreali. Muſ. D. Banks.

30. R. femoribus dentatis, brunneus, pedibus teſtaceis. *guloſus.*
*Curculio guloſus.* Ent. ſyſt. 2. 441. 194. *
Habitat Pariſiis. Dom. Grandchamp.

31. R. femoribus dentatis, ferrugineus, cinereo-irroratus, femoribus macula nigra. *paraſita.*
*Curculio paraſita.* Ent. ſyſt. 2. 441. 195. *
Oliv. Inſ. 83. fig. 181.
Habitat in Cayenna. Muſ. D. Geoffroy.

32. R. femoribus dentatis, griſeus, ſcutello elytrorumque macula poſtica albis. *eſuriens.*
*Curculio eſuriens.* Ent. ſyſt. 2. 441. 196. *

Ent.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Ent. Helv. 1. 112. 12. tab. 11. fig. 3. 4.
Habitat in Gallia.

*auarus.* 233. R. femoribus dentatis, obſcure ferrugineus, thorace linea dorſali, elytris faſcia poſtica albis.
*Curculio auarus.* Ent. ſyſt. ſuppl. 170. 196.
Habitat in Gallia. Muſ. D. Boſc.

*C. album.* 234. R. femoribus dentatis, ater, coleoptris duplici albo notatis.
*Curculio C. album.* Ent. ſyſt. ſuppl. 170. 197.
Habitat in India orientali. Muſ. D. de Seheſtedt.

*naſutus.* 235. R. femoribus dentatis, brunneus, elytris ſtriatis cinereis, apice brunneis.
*Curculio naſutus.* Ent. ſyſt. 2. 441. 197.
Oliv. Inſ. 83. fig. 138.
Habitat - - - Muſ. Britann.

*hauſtellatus.* 236. R. femoribus dentatis, caſtaneus, elytris ſtriatis, roſtro corpore quadruplo longiore.
*Curculio hauſtellatus.* Ent. ſyſt. 2. 442. 198.
Oliv. Inſ. 83. fig. 171.
Habitat ad Cap Bon. Spei. Muſ. D. Lee.

*varius.* 237, R. femoribus dentatis, niger, cinereo-irroratus, roſtro pedibusque rufis.
*Curculio varius.* Ent. ſyſt. 2. 442. 199.
Habitat in Cayenna. D Mallet.
Pedes rufi, apice nigricantes, antici elongati.

*Ceraſorum.* 238. R. femoribus dentatis, fuſcus, ſcutello elytrorumque faſciis obſoletis cinereis.
*Curculio Ceraſorum* Ent. ſyſt. 2. 442. 200.
Panz. Fn. Germ. 42. tab. 22.
*ff.* 73 Payk. Monogr. 22. 21.
Fn. Sv. 3 206. 22.
Herbſt. Col. 6. tab. 73. fig. 10.
Habitat in Anglia. D. Lee.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

139. R. femoribus dentatis, ater, thorace punctis duobus, coleoptris quinque niueis. *amoenus.*

*Curculio amoenus.* Ent. ſyſt. 2. 442. 201. *
Oliv. Inſ. 83. fig. 143.
Habitat in noua Hollandia. Muſ. D. Banks.

140. R. femoribus dentatis, nigro-aeneus, thorace punctato, elytris ſtriato-punctatis. *metallinus*

*Curculio metallinus.* Ent. ſyſt. 2. 442. 202. *
Habitat in Americae meridionalis Inſulis. Dom. Smidt.

141. R. femoribus dentatis, griſeus, capite bidentato. *bicornis.*

*Curculio bicornis.* Ent. ſyſt. 2. 443. 203. *
Habitat in noua Zelandia. Muſ. D. Banks.

142. R. femoribus dentatis, niger, elytris albo-ſubfaſciatis, antennis rufis. *tenuiroſtris.*

*Curculio tenuiroſtris.* Ent. ſyſt. 2. 443. 204. *
Habitat in Anglia. Muſ. D. Banks.

143. R. femoribus dentatis, elytris teſtaceis, obſolete faſciatis. *Druparum.*

*Curculio Druparum.* Ent. ſyſt. 2. 443. 205.
Linn. ſyſt. Nat. 2. 614. 62. Fn. Sv. 617.
Geoff. Inſ. 1. 296. 43.
Degeer Inſ. 5. 214. 7.
Schaeff. Icon. tab. 1. fig. 11.
Sulz. Hiſt. Inſ. tab. 3. fig. 21.
Payk. Monogr. 11. 10.
Fn. Sv. 3. 197. 14.
Herbſt. Col. 6. tab. 70. fig. 9.
Habitat in Europae Pruno Pado.

144. R. femoribus dentatis, ferrugineo-fuſcus, capite dorſoque nigris. *Fraxini.*

*Curculio Fraxini.* Ent. ſyſt. 2. 443. 206. *
Degeer Inſ. 5. 212. 4.

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Monogr. 25. 23.
Fn. Sv. 3. 209. 25.
Herbſt. Col. 6. tab. 73. fig. 5.
Habitat in Sueciae Fraxino. D. de Paykull.

*melanocephalus.* 245. R. femoribus dentatis, brunneus, capite fuſco, roſtro atro.

II. 57. *Curculio melanocephalus.* Ent. ſyſt. 2. 444. 207. *

Panz. Fn. Germ. 18. tab. 12.
Habitat Halae Saxonum. Dom. Hybner.

*pectoralis.* 246. R. femoribus dentatis teſtaceis, pectore nigro.

*Curculio pectoralis.* Panz. Fn. Germ.
Habitat in Auſtria. D. de Meyerle.

III. 62 Statura parua R. melanocephali. Caput et thorax teſtacea, immaculata. Elytra punctato-ſtriata, teſtacea, magis nebuloſa. Pectus nigrum. Pedes teſtacei.

*flauipes.* 247. R. fuſcus, elytris cinereo-variis, pedibus flauis.

*Curculio flauipes.* Panz. Fn. Germ.
Habitat in Auſtria. D. de Meyerle.

VI. 38 Statura parua praecedentis. Caput et thorax fuſca, immaculata. Elytra haud ſtriata, cinereo fuſcoque variegata. Pedes omnes flaui.

*arcuatus.* 248. R. femoribus dentatis, pallidus, elytris cineraſcentibus.

II. 64. *Curculio arcuatus.* Panz. Fn. Germ.
Habitat in Auſtria. D. de Meyerle.
Paruus. Roſtrum elongatum, parum rufeſcens. Caput et thorax laeuia, pallida, immaculata. Elytra ſtriata, magis cineraſcentia. Pedes acute dentati, pallidi.

*depreſſus.* 249. R. femoribus dentatis, thorace depreſſo: lateribus obtuſe angulatis.

*Curculio depreſſus.* Ent. ſyſt. 2. 444. 208. *
Linn.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Linn. syst. Nat. 2. 612. 53. Mus. Lud. Vlr. 49.
Seb. Mus. 4. tab. 95. fig. 4.
Habitat in America meridionali.

50. R. femoribus anticis dentatis, corpore griseo nebuloso. ***Pomorum.***
*Curculio Pomorum.* Ent. syst. 2. 444. 209.
Linn. syst. Nat. 2. 612. 46. Fn. Sv. 612.
Frisch. Inst. 1. 32. tab. 8. III. 63.
Payk. Monogr. 13. 12.
Fn. Sv. 3. 199. 7.
Degeer Inf. 5. 215. 8. tab. 6. fig. 26. 27.
Herbst. Col. 6. tab. 70. fig. 11.
Habitat in Pomonae floribus.

251. R. femoribus dentatis, griseo fuscoque maculatus, rostro arcuato atro. ***vorax.***
*Curculio vorax.* Ent. syst. 2. 444. 210.
*Curculio longimanus.* Forst. Cent. Inf. 1. 24.
Payk. Monogr. 3. 3.
Fn. Sv. 3. 189. 5.
Herbst. Col. 6. tab. 71. fig. 4. II. 57
Panz. Fn. Germ. 18. tab. 13.
Habitat in Italia. Dr. Allioni.

252. R. femoribus dentatis, corpore testaceo, pectore fusco. ***tortrix.***
*Curculio tortrix.* Ent. syst. 2. 444. 211. *
Linn. syst. Nat. 2. 615. 67. Fn. Sv. 622.
Geoff. Inf. 1. 300. 51.
Degeer Inf. 5. 214. 6.
Ent. Helv. 1. 92. 8. tab. 9. fig. 3. 4.
Panz. Fn. Germ. 18. tab. 19. II. 58
Payk. Monogr. 7. 6.
Fn. Sv. 3. 192. 9.
Herbst. Col. 6. tab. 7. fig. 8.
Habitat in Populo, folia contorquens.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Tremulae.* 253. R. femoribus dentatis, nigricans, elytris ſtriatis griſeo-irroratis.
*Curculio Tremulae.* Ent. ſyſt. 2. 445. 212.
Payk. Monogr. 4. 4.
Fn. Sv. 3. 189. 6.
Habitat in Sueciae Populo. D. de Paykull.

*aeſtuans.* 254. R. femoribus dentatis, fuſcus, cinereo-lineatus, pedibus pallidis.
*Curculio aeſtuans.* Ent. ſyſt. 2. 446. 213.
Habitat in Americae meridionalis Inſulis. Dr. Smidt.

*taeniatus.* 255. R. femoribus dentatis, thorace nigro: margine antico poſticoque rufis, elytris pallidis, nigro maculatis.
*Curculio taeniatus.* Ent. ſyſt. 2. 445. 214.
Payk. Monogr. 6. 5.
Fn. Sv. 3. 191. 8.
Habitat in Anglia. Muſ. D. Banks.

*** *Femoribus ſaltatoriis.*

*Alni.* 256. R. pedibus ſaltatoriis, niger, coleoptris teſtaceis: maculis duabus obſcuris.
*Curculio Alni.* Ent. ſyſt. 2. 445. 216.
Linn. ſyſt. Nat. 2. 611. 42. Fn. Sv. 608.
Geoff. Inſ. 1. 286. 20.
Degeer Inſ. 5. 262. 49.
Payk. Monogr. 20. 19.
Fn. Sv. 3. 220. 39.
Habitat in Betula, Alno.

*craſſus.* 257. R. pedibus ſaltatoriis, ſupra fuſcus, thorace quadri-tuberculato, elytris ſtriatis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Magnus in hac familia. Thorax fuſcus, tuberculis quatuor eleuatis. Elytra ſtriata, fuſca, obſcura.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

ſcura. Corpus brunneum, femoribus poſticis incraſſatis.

58. R. pedibus ſaltatoriis, niger, cinereo - variegatus. *piloſus.*
*Curculio piloſus.* Ent. ſyſt. 2. 446. 217. *
Payk. Monogr. 18. 17.
Fn. Sv. 2. 318. 36.
Habitat in Anglia. Muſ. D. Banks.

59. R. pedibus ſaltatoriis, ater, elytrorum faſcia *hortorum.*
ſesquialtera pedibusque teſtaceis.
*Curculio hortorum.* Ent. ſyſt. 2. 446. 218. *
*Curculio mutilatus.* Laichart. Auſtr.
Habitat in Germania. Muſ. D. de Seheſtedt.

60. R. pedibus ſaltatoriis, niger, obſcurus, roſtro *rufeſcens.*
tibiisque rufis.
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Paullo maior, quam R. calcar. Antennae rufae, claua faſcia fuſca. Caput fuſcum, roſtro arcuato, rufo. Thorax et elytra laeuia, nigra, obſcura. Pedes nigri, femoribus poſticis incraſſatis, ſubdentatis.

261. R. pedibus ſaltatoriis, niger, antennis plantisque teſtaceis, femoribus dentatis. *calcar.*
*Curculio calcar.* Ent. ſyſt. 2. 446. 219. * Ill. 62.
Payk. Monogr. 18. 17.
Habitat Kiliae. D. Daldorff.

262. R. pedibus ſaltatoriis, niger, tibiis teſtaceis. *Saliceti.*
*Curculio Saliceti.* Ent. ſyſt. 2. 446. 220. *
Payk. Monogr. 66. 64.
Fn. Sv. 3. 271. 94.
Herbſt. Col. 6. 430. 418.
Habitat in Sueciae Salicetis.

263. R. pedibus ſaltatoriis, coleoptris ante apicem rufis. *Beccabungae.*

*Cur-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Beccabungae.* Ent. syst. 2. 447. 221.
Linn. syst. Nat. 2. 611. 41. Fn. Sv. 604.
Payk. Fn. Sv. 3 254. 72.
Habitat in Sueciae Veronica Beccabunga. D. de Paykull.
Haud saltat teste D. de Paykull.

*Salicis.* 264. R. pedibus saltatoriis, elytris atris: fasciis duabus albis.
*Curculio Salicis.* Ent. syst. 2. 447. 222.
Linn. syst. Nat. 2. 611. 43. Fn. Sv. 610.
Degeer Inf. 5. 246. 51.
Panz. Fn. Germ. 18. tab. 15.
Payk. Monogr. 64. 62.
Fn. Sv. 3. 269. 91.
Herbst. Col. 6. tab. 93. fig. 2.
Habitat in Salicis floribus.

*viminalis.* 265. R. pedibus saltatoriis, corpore testaceo.
*Curculio viminalis.* Ent. syst. 2. 447. 223.
*Curculio Quercus.* Linn. syst. Nat. 2. 609. Fn. Sv. 596.
*Curculio Vlmi.* Degeer Inf. 5. 260. 48. tab. 8. fig. 5.
Geoff. Inf. 1. 286. 19.
Reaum. Inf. 3. 31. tab. 3. fig. 17. 18.
Payk. Monogr. 19. 18.
Fn. Sv. 3. 219. 38.
Herbst. Col. 6. tab. 93. fig. 1.
Habitat in Europae Salice.

*Ilicis.* 266. R. pedibus saltatoriis, nigricans, elytris striatis, cinereo-variis: sutura basi alba.
*Curculio Ilicis.* Ent. syst. 2. 447. 224.
Payk. Monogr. 18. 17.
Fn. Sv. 3. 219. 37.
Herbst. Col. 6. 431. 420.
Habitat in Vplandiae Quercu.

267

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

267. R. pedibus ſaltatoriis, teſtaceus, elytris femoribusque faſcia nigra. *Loniceraе.*

Habitat in Italiae Lonicera. Muſ. D. de Seheſtedt.

Statura et magnitudo omnino R. viminalis. Caput et thorax rufa, immaculata. Scutellum album. Elytra ſtriata, teſtacea: faſcia media fuſca. Pedes teſtacei, femoribus poſticis incraſſatis: faſcia fuſca.

268. R. pedibus ſaltatoriis, teſtaceus, ſcutello niueo. *ſcutellaris.*

Habitat in Germania. Muſ. D. Lund.

Affinis certe R. viminali, at minor et ſcutellum omnino niueum.

269. R. pedibus ſaltatoriis, ater, elytris ſtriatis: ſutura baſi alba. *Iota.*

*Curculio Iota* Ent. ſyſt. 2. 448. 225.
Panz. Fn. Germ. 18. tab. 16.
Payk. Monogr. 66. 63.
Fn. Sv. 3. 271. 93.
Herbſt. Col. 6. 432. 421.

Habitat in Vplandiae arboribus. D. de Paykull.

270. R. pedibus ſaltatoriis, corpore atro, femoribus albis. *Fagi.*

*Curculio Fagi* Ent. ſyſt. 2. 448. 226.
Linn. ſyſt. Nat. 2. 64. 44. Fn. Sv. 609.
Payk. Monogr. 64. 61.

Habitat in Fagi ſyluatici foliis.

271. R. pedibus ſaltatoriis, faſciis, antennis tarſisque teſtaceis. *Fragariae*

*Curculio Fragariae.* Ent. ſyſt. 2. 448. 227.

Habitat in Germaniae Fragariis. D. Smidt.

272. R. pedibus ſaltatoriis, ater, ſcutello albo, antennis pedibusque teſtaceis. *Populi.*

*Curculio Populi.* Ent. ſyſt. 2. 448. 228. II. 59.
*Rhynchaenus Populi.* Ent. Helv. 1. 72. 2.

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 18. tab. 17.
Habitat in Selandiae Populo. Muſ. D. de Seheſtedt.

160. COSSONVS. *Mandibula* incurua, acutiſſima.
*Antennae* claua ſolida, vniarticulata.

*linearis.* 1. C. niger, antennis pedibusque piceis.
*Curculio linearis* Ent. ſyſt. 2. 419. 110.
*Coſſonus linearis.* Ent. Helv. 1. 60. 1. tab. 1. fig. 1. 2.
1156. Panz. Fn. Germ. 18. tab. 7.
Payk. Monogr. 51 48.
Fn. Sv. 3. 242. 60.
Herbſt. Col. 6. 53. 15.
Habitat Strasburgi. D. Prof. Hermann.
*Variat* paullo minor, thorace elytrisque ferrugineis, at vix diſtinctus.
*Coſſonus ferrugineus.* Ent. Helv. 1. 60. 2. tab. 1. fig. 3. 4.

*elongatulus.* 2. C. ferrugineus, thorace canaliculato, femoribus compreſſis.
*Curculio elongatulus.* Ent. ſyſt. ſuppl. 68. 111.
Habitat ad Cap. Bon. Spei. D. Daldorff.

*canaliculatus.* 3. C. ater, abdomine ruſo, thorace punctato: dorſo canaliculato.

Habi-

---

*Coſſoni* corpus elongatum, cylindricum, glabrum, immarginatum, tardum; *capite* paruo, ſubrotundo, antice porrecto in roſtrum elongatum, rectum, apice incraſſatum; *antennis* articulo ſecundo longiſſimo, vltimo craſſiori, clauato, ante apicem incraſſatum inſertis; *oculis* rotundatis, lateralibus; *thorace* planiuſculo, latitudine elytrorum; *elytris* rigidis, foraminatis, longitudine abdominis, includentibus: *pedibus* brevibus, validis, compreſſis; *tarſis* quadriarticulatis; *colore* obſcuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Sumatra. D. Daldorff.

Statura praecedentium. Antennae nigrae, claua cinerea. Thorax punctatus, in medio glaber, planus, canaliculatus: canali basi dilatato. Elytra punctato-striata, nigra. Abdomen obscure-rufum. Pedes breues: femoribus clauatis.

Character generis naturalis adhuc desideratur.

C. ater, nitidus, elytris striato-punctatus. *corticalis.*

Habitat in America meridionali.

Corpus totum depressum, planum. Antennae nigrae: claua cinerascente. Thorax postice subtrilobus. Elytra punctato-striata. Corpus nigrum.

C. thorace punctato, postice retuso, ater, elytris macula apicis ferruginea. *squama.*

Habitat in America meridionali. D. Smidt. Musc. D. Lund.

Statura vtique praecedentis, at omnino distinctus. Thorax punctatus, apice retusus. Elytra macula magna versus apicem ferruginea.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

161. LIXVS. *Palpi* quatuor, aequales, breuiſſimi: articulo vltimo ſubulato.

*Maxilla* cornea, acuta, integra.

*Antennae* clauatae: articulo primo longiſſima, inſidentes roſtro craſſiuſculo.

* *femoribus muticis.*

*paraplecticus* 1. L. ſubcinereus, elytris mucronatis.

*Curculio paraplecticus*. Ent. ſyſt. 2. 414. 91.
Linn. ſyſt. Nat. 2. 610. 24. Fn. Sv. 604.
Scan. 184.
Oliv. Inſ. 83. fig. 106.
Ent. Helv. 1. 94. 9. tab. 10. fig. 1. 2.
Geoff. Inſ. 1. 279.
Schreb. Saml. 9. 336. tab. 4.
I. 48 Degeer Inſ. 5. 224. 18. tab. 7. fig. 8.
Sulz. Hiſt. Inſ. tab. 4. fig. 1.
Schaeff. Icon. tab. 44. fig. 1.
Payk. Monogr. 53. 50.
Fn. Sv. 3. 244. 63.
Herbſt. Col. 6. tab. 62. fig. 1.

Hi-

---

*Lixi* corpus elongatum, cylindricum, immarginatum, tardum; *capite* paruo, inſerto, antice porrecto in roſtrum rectum, cylindricum, craſſiuſculum, vtrinque ſulco obliquo exaratum, ad receptionem articuli ſecundi antennarum; *oculis* paruis, ob longis, lateralibus, ad baſin roſtri; *antennis* thorace breuioribus: articulo ſecundo longiſſimo, vltimis clauatis: clava acuta; *thorace* breui, rotundato, antice parum angustato; *elytris* rigidis, includentibus, longitudine abdominis, interdum apice acuminatis; *pedibus* longiuſculis, curſoriis; *tarſis* quadriarticulatis; *colore* obſcuro, ſaepius polline adſpe-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat Larua intra caulem Vmbellatarum aquaticarum. Paraplegiam equorum cauſſare fertur.

2. L. cylindricus, flauo-puluerulentus, elytris acuminatis. *puluerulentus.*

Habitat in Auſtria. Dom. de Meyerle.

Statura omnino praecedentis, paullo minor. Corpus totum atomis luteis, puluerulentis tectum, thorace vix linea laterali fuſca. Elytra acuminata.

3. L. cylindricus, cinereus, thorace elytrisque fuſco-ſtriatis, elytris mucronatis. *elegans.*

Habitat in Germania. D. Smidt.

Duplo minor L. paraplectico. Caput cinereum, roſtro apice nigro. Thorax cinereus, ſtriis duabus dorſalibus, latis, nigris. Elytra ante apicem vnidentata, apice valde mucronata, cinerea, ſtriis tribus diſtinctis, fuſcis.

4. L. cylindricus, canus, fuſco-lineatus. *anguinus.*

*Curculio anguinus* Ent. ſyſt. 2. 415. 92.
Linn. ſyſt. Nat. 2. 610. 35.
Ent. Helv. 1. 96. 70. tab. 10. fig. 3. 4.
Oliv. Inſ. 33. fig. 168. *IV. 70*
Panz. Naturf. 24. 19. 26. tab. 1. fig. 26.
Panz. Fn. Germ. 42. tab. 11.

Habitat in Germania.

*Os* maxillis palpisque. *Palpi* quatuor, aequales, breuiſſimi, filiformes; *anteriores* vix longiores, quadriarticulati: articulis rotundatis, aequalibus; vltimo ſubulato, adhaerentes maxillae dorſo; *poſteriores* breuiſſimi, triarticulati: articulis rotundatis, aequalibus, adnati labii baſi. *Mandibula* magna, fornicata, cornea, lata, incurua, apice multidentata. *Maxilla* breuiſſima, cornea. *Ligula* nulla? *Labium* corneum, ro-



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

tundatum, integrum. *Antennae* fractae, clauatae: claua acuta, ante apicem roſtri inſertae: articulo ſecundo in canalem flexuoſum roſtri recondendo.

*ſtriatellus.* 5. L. obſcurus, elytris cinereis, fuſco-ſubſtriatis.
*Curculio ſtriatellus.* Ent. ſyſt. 2. 415. 93. [3]
Habitat in Barbariae plantis. Dom. Desfontaines.

*Vmbellatarum.* 6. L. thorace nigro, cinereo, lineato, elytris mucronatis, griſeis.
*Curculio Vmbellatarum.* Ent. ſyſt. 2. 415. 94. [4]
Habitat in Barbariae vmbellatis. Dom. Prof. Vahl.

*mucronatus.* 7. L. cylindricus, cinereus, fuſco-lineatus, elytris mucronatis.
*Curculio mucronatus.* Ent. ſyſt. 2. 415. 95. [5]
Habitat in Barbariae plantis.
Affinis L. anguino et forte idem.

*ferrugatus* 8. L. niger, ferrugineo-villoſus, elytris obtuſis.
*Curculio ferrugatus* Ent. ſyſt. 2. 416. 96. [6]
Habitat in Hungaria. Dom. Hybner.
Varietas duplo minor mas.

*gages.* 9. L. ater, nitidus, elytris ſtriatis, obſcuris, roſtro truncato.
*Curculio gages.* Ent. ſyſt. 2. 416. 97. [7]
Habitat in Aquapim Guineae. Dr. Iſert.

*8 lineatus.* 10. L. cylindricus, niger, thorace elytrisque lineis quatuor albis.
*Curculio 8 lineatus.* Ent. ſyſt. 2. 416. 98.
Oliv. Inſ. 83. fig. 89.
Habitat ad Cap. Bon. Spei. Muſ. Gallicum.

*ſemipunctatus.* 11. L. cylindricus, niger, thorace lineis, elytris punctis albis.
*Curculio ſemipunctatus.* Ent. ſyſt. 2. 416. 99. [8]
Oliv. Inſ. 83. fig. 141.

Hi-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in noua Hollandia. Muſ. D. Banks.

2. L. niger, elytris maculis duabus ferrugineis. *4 puſtulatus.*
*Curculio 4 puſtulatus.* Ent. ſyſt. 2. 416. 100. *
Oliv. Inſ. 83. fig. 117.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

3. L. cylindricus, albo fuſcoque nebuloſus, elytris ſubmucronatis, antennarum claua teſtacea. *mixtus.*
*Curculio mixtus.* Ent. ſyſt. 2. 417. 101. *
Habitat in Barbariae plantis. Muſ. Dom. Desfontaines.

4. L. ater, thorace elytrisque ſcabris, fuſcis, femoribus anticis incraſſatis. *melanocephalus.*
*Curculio melanocephalus.* Ent. ſyſt. ſuppl. 168. 101. *
Habitat in Isle de France. D. Daldorff.
Roſtrum breue. Forte potius Curculio. C. brunnipe affinis videtur.

5. L. cylindricus, ſubcinereus, thorace lineis tribus fuſcis. *filiformis.*
*Curculio filiformis.* Ent. ſyſt. 2. 417. 102. *
Habitat in Italia. Dr. Allioni.

6. L. cylindricus, ſupra niger, elytris acuminatis: faſcia ~~nigra~~ *pallida.* *cylindricus.*
*Curculio cylindricus.* Ent. ſyſt. 2. 417. 103. *
Oliv. Inſ. 83. fig. 123.
Habitat in Sibiria. Muſ. D. Banks.

7. L. thorace fuſco: punctis quatuor albis, elytris fuſcis: faſcia ſesquialtera teſtacea. *notatus.*
*Curculio notatus.* Ent. ſyſt. 2. 417. 104.
Habitat in Barbariae plantis. Dom. Prof. Vahl.

8. L. niger, roſtro barbato, tibiis anticis tridentatis. *barbiroſtris.*
*Curculio barbiroſtris.* Ent. ſyſt. 2. 418. 105.
Oliv. Inſ. 83. fig. 37.

Seb.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Seb. Muſ. 4. tab. 95. fig. 5.
Habitat in Indiis.

*amphora.* 19. L. cylindricus, griſeo-villoſus, roſtro, pedib abdomineque ferrugineis.
Habitat Algirae. Dom. de Rehbinder.
Statura et magnitudo L. anguſtati. Roſtrum i curuum, cylindricum, ferrugineum. Thorax g ſeo-villoſus. Elytra obſcuriora, hinc inde g ſeo-villoſa. Abdomen et pedes ferrugin Elytra apice rotundata.

*anguſtatus* 20. L. cylindricus, ater, elytris obtuſis, punctatis
*Curculio anguſtatus* Ent. ſyſt. 2. 418. 106.
Herbſt. Arch. tab. 24. fig. 7.
Roſſ. Fn. Etr. 1. 111. 283.
*IV. 70* Panz. Fn. Germ. 42. tab. 12.
Habitat in Anglia.

*Pica.* 21. L. cylindricus, niger, albo-variegatus, pedib anticis elongatis.
*Curculio Pica.* Ent. ſyſt. ſuppl. 197. 106.
Habitat in Isle de France. Dom. Daldorff.

*Bardanae.* 22. L. cylindricus, griſeo-tomentoſus, pedibus a ticis elongatis.
*Curculio Bardanae.* Ent. ſyſt. 2. 418. 107.
*II. 55.* Panz. Fn. Germ 18 tab. 3.
Habitat in Saxoniae Bardana. Muſ. Dom. Roma

*truncatulus.* 23. L. cylindricus, fuſcus, obſcurus, elytris punct to-ſtriatis, roſtro truncato.
*Curculio truncatulus* Ent. ſyſt. ſuppl. 167. 107
Habitat in India orientali. Dom. Daldorff.

*volvulus.* 24. L. cylindricus, fuſcus, elytris caſtaneis, griſe variis.
*Curculio volvulus.* Ent. ſyſt. ſuppl. 167. 107.
Habitat in India orientali. Dom. Daldorff.

2

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. L. cylindricus, ſupra ferrugineus, ſubtus cinereus, abdomine nigro, punctato. *algirus.*

Habitat Algirae. Dom. de Rehbinder.

Statura praecedentium. Caput ferrugineum, roſtro cylindrico, apice nigro. Thorax ſcaber, ferrugineus. Elytra ferruginea, atomis vix conſpicuis, griſeis, ſtriata, puncto tamen baſeos maiori, diſtincto. Subtus cinereus, abdomine punctis numeroſis, ſubocellaribus, nigris.

6. L. cylindricus, ater, lateribus ſubcoeruleſcentibus. *Aſcanii.*

*Curculio Aſcanii.* Ent. ſyſt. 2. 418. 108.
Linn. ſyſt. Nat. 2. 610. 36.
Oliv. Inſ. 83. fig. 83.
Panz. Fn. Germ. 42. tab. 13.
Herbſt. Arch. tab. 24. fig. 8.
Habitat in Europa auſtraliori.

7. L. ater, thoracis lateribus elytrorumque ſtriis albis. *alboſtriatus.*

Habitat in Guinea. D. Heſſe. Muſ. D. Lund.

Statura et magnitudo L. Aſcanii. Roſtrum arcuatum, vti thorax laeuis, ater, lateribus late cinereis. Elytra ſtriata, ſtriis tribus approximatis, in medio marginali abbreuiata albis. Corpus atrum, albo-varium. Pedes picei.

8. L. cylindricus, niger, elytris vitta teſtacea. *lineola.*

*Curculio lineola.* Ent. ſyſt. 2. 419. 109. *

Oliv. Inſ. 83. fig. 185.

Habitat in noua Hollandia. Muſ. D. Banks.

9. L. elongatus, ater, thorace linea dorſali alba, pedibus elongatis piceis: femoribus eleuatis. *clauipes.*

Habitat in noua Cambria. D. Billardiere.

Statura praecedentium. Caput atrum, roſtro incuruo. Thorax punctatus, ater: linea dorſali alba.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

alba. Elytra punctato-ſtriata, atra. Corpus la teribus albo-tomentoſis. Pedes ſimplices, elon gati, picei, femoribus clauatis.

*craſſipes.* 30. L. femoribus anticis ſubclauatis, corpore atro.
*Curculio craſſipes.* Ent. ſyſt. 2. 419. 110. *
Habitat Strasburgi. Muſ. D. Prof. Hermann.

*Atriplicis.* 31. L. elongatus, ater, thorace nitido, elytris ſtria tis obtuſis.
*Curculio Atriplicis.* Ent. ſyſt. 2. 419. 112. *
*Curculio T. album.* Linn. ſyſt. Nat. 2. 609. 23
Fn. Sv. 594.
Payk. Monogr. 52. 49.
Fn. Sv. 3. 243. 62.
Herbſt. Col. 6. 67. 31.
Habitat in Atriplice littorali Florae Kilonienſis
Muſ. D. Lund.

*Lymexylon.* 32. L. elongatus, griſeus, thorace ſcabro, elytris ſtriatis.
*Curculio Lymexylon.* Ent. ſyſt. 2. 420. 113. *
Panz. Fn. Germ. 18. tab. 11.
Payk. Monogr. 41. 38.
Fn. Sv. 3. 240. 59.
Herbſt. Col. 6. tab. 76. fig. 9.
Habitat in Germaniae ligno quercino. D. Prof. Helwig.

** *Femoribus dentatis.*

*caliginoſus* 33. L. femoribus dentatis, ater, elytrorum ſtriis punctatis approximatis.
*Curculio caliginoſus.* Ent. ſyſt. 2. 427. 140. *
Habitat in Anglia. Muſ. Britt.

*dubius.* 34. L. femoribus dentatis, niger, thorace laeui, elytris ſtriatis ſcabris.
*Curculio dubius.* Ent. ſyſt. 2. 427. 141. *
Habitat - - Dr. Schulz.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. L. femoribus dentatis, fuluo-irroratus, pedibus anticis elongatis. *roreus.*

*Curculio roreus.* Ent. ſyſt. 2. 427. 142. *

Habitat in Cayenna. D. v. Rohr.

6. L. femoribus dentatis, brunneus, roſtro fuſco, elytris teſtaceis punctato-ſtriatis. *brunneus.*

*Curculio brunneus.* Ent. ſyſt. 2. 427. 143.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

7. L. femoribus dentatis, oblongus, piceus, elytris reticulatis: faſciis obliquis pallidis, tibiis anticis ſpinoſis. *reticulatus.*

*Curculio reticulatus.* Ent. ſyſt. 2. 429. 148. *

Habitat Tranquebariae. Dr. Koenig.

8. L. femoribus dentatis, niger, thorace trilobo ſcabro, elytris punctato-ſtriatis brunneis. *trilobus.*

Habitat in Guinea. D. Meyer. Muſ. D. Lund.

Statura et magnitudo praecdentis. Roſtrum breuiuſculum, ſubincuruum, nigrum. Thorax trilobus, ſcaber, niger, linea vna alteraue, obſoleta alba. Elytra punctato-ſtriata, brunnea. Corpus cinereo-villoſum, pedibus nigris: femoribus dentatis.

9. L. femoribus dentatis, niger, albo-lineatus, abdomine prominente acuto. *acutus.*

Habitat in Sumatra. D. Daldorff. Muſ. D. de Seheſtedt.

Statura cylindrica L. filiformis. Roſtrum porrectum, rectum, cylindricum, obſcure ferrugineum. Antennae fractae, ferrugineae, apice parum albicantes. Thorax planus, laeuis. niger, lineis duabus dorſalibus, lateralique albis. Elytra punctato-ſtriata, abbreuiata, obſcura: linea ad ſuturam atomisque plurimis albis. Corpus albo-villoſum, abdomine valde prominente,

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nente, acuto. Pedes nigri, femoribus acute dentatis.

*attenuatus* 40. L femoribus dentatis, ater, nitidus, elytris femoribusque apice ſpinoſis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Corpus valde elongatum, tenue. Roſtrum cylindricum, longitudine corporis; atrum, antennis articulo ſecundo reliquis omnibus longiore, in medio roſtri inſerto. Caput minutum. Thorax cylindricus, laeuis, ater, nitidus. Elytra cylindrica, laeuia, atra, nitida, pilis rarioribus adſperſa, apice ſpina valida, acuta. Pedes elongati, nigri, femoribus dentatis, apice vniſpinoſis.

*Antilope.* 41. L. femoribus dentatis, ater, nitidus, thorace varioloſo, femoribus clauatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentis, at paullo minor et diſtinctus. Roſtrum elongatum, longitudine corporis incuruum. Thorax globoſus, varioloſus, piloſus. Elytra laeuia, atra, nitida, apice inermia. Pedes elongati, femoribus dentatis, clauatis apice muticis.

*Gazella.* 42. L. femoribus dentatis, piloſus, ater, nitidus, roſtro thorace breuiore.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentium. Corpus elongatum, piloſum, atrum. Roſtrum thorace breuius, cylindricum, rectum. Thorax piloſus, gibbus. Elytra gibba, piloſa, laeuia, immaculata. Corpus atrum, abdominis margine baſi albido. Pedes elongati, piloſi, nigri.

Alius

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Alius roſtro thorace longiori ſimillimus, nullo modo differt.

3. L. femoribus dentatis, thorace elongato, tibiis anticis bidentatis. *elongatus.*

*Curculio elongatus.* Ent. ſyſt. 2. 445. 215.
Habitat in Iamaica. D. Drury.
Species mihi obſcura.

## 62. CVRCVLIO.

*Palpi* quatuor, breuiſſimi: articulo vltimo longiori, ſubulato.
*Maxilla* breuis, apice dilatata, truncata.
*Antennae* fractae, roſtro breui, inſidentes.

* *Femoribus muticis.*

1. C. corpore viridi, argenteo auroque maculato. *argyreus.*
Ent. ſyſt. 2. 448. 229.
*Curculio argyreus.* Linn. ſyſt. Nat. 2. 615. 74. Muſ. Lud. Vlr. 54.
Habitat in Indiis.

2. C. elytris ſtriatis, auro-maculatis, baſi gibbis, apice acuminatis. Ent. ſyſt. 2. 448. 230. *ſplendidus.*
Habitat in Braſilia. Muſ. D. Banks.

3.

---

*Curculionis* corpus oblongum, craſſum, immarginatum, tardum; *capite* paruo, inſerto, antice porrecto in roſtrum breue, craſſum, obtuſum vtrinque ſulco obliquo, ad receptionem ſecundi antennarum articulo: *oculis* paruis, oblongis, lateralibus ad baſin roſtri; *antennis* thorace breuioribus: articulo ſecundo longiſſimo, reliquis ſubaequalibus, vltimis craſſioribus: claua acutiuſcula; *thorace* rotundato, antice parum attenuato; *ſcutello* nullo; *elytris* rigidis, includentibus, longitudine abdominis; *pedibus* validis, curſoriis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*imperialis.* 3. C. elytris ſtriis eleuatis, atris ſulcisque punctatis viridi-aureis alternis, baſi gibbis, apice acuminatis. Ent. ſyſt. 2. 449. 231.
Drury Inſ. 2. tab. 33. fig. 1.
Oliv. Inſ. 83. fig. 1.
Hubner. Naturf. 10. tab. 2. fig. 1.
Habitat in America meridionali et Braſilia.

*ſumptuoſus* 4. C. elytris vireſcentibus: punctis eleuatis atris baſi gibbis. Ent. ſyſt. 2. 449. 232.
*Curculio ſumptuoſus.* Oliv. Inſ. 83. tab. 1. f. 15.
Habitat in Cayenna.

*Chryſis.* 5. C. albidus, elytris acuminatis: faſciis duabus punctisque duobus aureis. Ent. ſyſt. 2. 449. 233.
*Curculio Chryſis.* Oliv. Inſ. 83. tab. 1. fig. 6.
Habitat in India orientali. Muſ. Gallic.

*regalis.* 6. C. corpore viridi-ſericeo: faſciis aureis repandis. Ent. ſyſt. 2. 449. 234.
*Curculio regalis.* Linn. ſyſt. Nat. 2. 616. 75.
Oliv. Inſ. 83. tab. 1. fig. 8.
Habitat in Indiis.

*marginatus.* 7. C. fuſcus, elytrorum futura abdominisque marginibus aureis. Ent. ſyſt. 2. 449. 235.
Oliv. Inſ. 83. fig. 82.
Habitat in Cayenna. Muſ. Dr. Hunter.

*aeneus.* 8. C. aeneus, elytris laeuiſſimis, poſtice virdi-maculatis.
Habitat in Amboina. D. Billardiere.
Paullo minor, quam C. regalis. Antennae nigrae, claua cinerea. Caput aeneum, nitidum: roſtro breui nigro. Thorax gibbus, globoſus, laeuiſſimus, aeneus, nitidus. Elytra laeuia, aenea, nitida, poſtice punctis variis lineaque marginali viridibus. Corpus atrum, vtrinque viridi-maculatum. Pedes atri, cyaneo-nitiduli. 9.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

. C. caneſcens, thorace punctis nigris quatuor, coleoptris nouemdecim. Ent. ſyſt. 2. 450. 236. * *19punctatus.*
Oliv. Inſ. 83. fig. 25.
Habitat in Cayenna. D. Mauduit.

o. C. coeruleſcens, thorace punctis nigris quatuor, coleoptris ſedecim. Ent. ſyſt. 2. 450. 237. * *16punctatus.*
*Curculio 16punctatus.* Linn. ſyſt. Nat. 2. 618. 92. Muſ. Lud. Vlr. 58.
Oliv. Inſ. 83. fig. 17.
Habitat in Braſilia.

1. C. ſupra fuſcus, thorace coleoptrisque vittis duabus viridi-aureis. Ent. ſyſt. 2. 450. 238. * *decorus.*
*Curculio decorus.* Oliv. Inſ. 83. fig. 152.
Habitat in Braſilia. Muſ. D. Banks.

2. C. varioloſus, niger, atomis aureis numeroſiſſimis. Ent. ſyſt. ſuppl. 170. 238. * *prodigus.*
Habitat in Tanger. D. Schousboe.

3. C. viridi-ſquamoſus, elytris punctatis. Ent. ſyſt. 2. 451. 239. * *nitidulus.*
*Curculio nitidulus.* Oliv. Inſ. 83. fig. 38.
Habitat in Cayenna.

4. C. elytris ſpinoſis albis: macula laterali fuſca. Ent. ſyſt. 2. 451. 240. * *candidus.*
Oliv. Inſ. 83. tab. 13. fig. 135.
Habitat in Cayenna. D. Mallet.

5. C. capite albido, thorace rugoſo, elytris puncato-ſtriatis viridibus. *pauperatus*
Habitat in Sumatra. D. Daldorff. Muſ. D. Lund.
Statura omnino C. candidi, at duplo maior. Antennae breues, craſſae, fractae, albidae: articulo vltimo craſſiori, nigro. Caput canaliculatum, albidum, fuſco purum irroratum. Thorax cylindricus, canaliculatus, rugis transuerſis inaequalis, viridis, ſtriis tribus latis, fuſcis.
Elytra

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Elytra punctato-ſtriata, viridia: ſutura fuſca Corpus viridi-ſquamoſum.

*auarus.* 16. C. cinereus, thorace rugoſo, elytris ſtriatis: vitta media obſoleta lateribusque viridi-albis.
Habitat in Iaua. Muſ. D. Boſc.
Affinis videtur C. pauperato. Corpus craſſum ouatum. Roſtrum breue, craſſum, canaliculatum, apice late emarginatum. Thorax rugis transuerſalibus inaequalis, cinereus, lineis quatuor viridi-albidis. Elytra ſtriata: ſtriis punctatis, cinereus: vitta parua media, obſoleta margineque late viridi-albis. Corpus cum pedibus cinereum.

*niueus.* 17. C. niueus, thoracis dorſo roſtroque fuſcis, elytris ſpinoſis acuminatisque. Ent. ſyſt. 2. 451. 241. *
Oliv. Inſ. 83. fig. 173.
Habitat in Cayenna. Muſ. Dr Hunter.

*lacteus.* 18. C. albus, auro-nitidulus, elytris ſulcatis, acuminatis. Ent. ſyſt. 2. 451. 242. *
Oliv. Inſ. 83 fig. 172.
Habitat in Iamaica. Muſ. Dr. Hunter.

*pulueruleutus.* 19. C. villoſus, vireſcens, elytrorum margine pulverulento, ferrugineo, capite plano, canaliculato. Ent. ſyſt. 2. 452. 243. *
Habitat Tranquebariae. Muſ. D. de Seheſtedt.

*ſquamoſus.* 20. viridi-ſquamoſus, roſtro thoraceque canaliculatis. Ent. ſyſt. 2. 452. 244 *
Habitat in China, in India orientali. Muſ. D. de Seheſtedt.

*ruſticus.* 21. C. villoſus, cinereus, roſtro ſubcarinato.
Habitat in Sumatra. Dom. Daldorff.
Affinis omnino C. vnicolori, at alius. Corpus totum villoſum, cinereum. Antennae breues, cinerea

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nereae. Roſtrum haud canaliculatum, ſed parum carinatum.

2. C. cinereo-pubeſcens, roſtro canaliculato, antennarum claua apice acuminata, atra. *vnicolor.*
Habitat in Sumatra. D. Daldorff.
Paullo maior C. ſquamoſo, totus pube cinerea tectus, immaculatus. Antennae breues, claua apice valde acuminata, atra. Roſtrum canaliculatum. Elytra punctato-ſtriata.

3. C. cylindricus, niger, elytris puncto baſeos apicisque viridi. *moratus.*
*Curculio moratus.* Ent. ſyſt. ſuppl. 171. 244.*
Habitat in Isle de France.

4. C. albus, elytris ſulcatis, vniſpinoſis acuminatisque. Ent. ſyſt. 2. 452. 245.* *cretaceus.*
Habitat in Inſula Guadeloupe. D. de Badier.

5. C. cinereus, elytris puncto baſeos duobusque apicis albis. Ent. ſyſt. 2. 452. 246.* *cinerascens.*
Habitat in Cayenna. Muſ. D. Gigot d'Orcy.

6. C. vireſcens, elytris punctatis: ſpina antica poſticaque erecta valida. Ent. ſyſt. 2. 452. 247.* *ſmaragdulus.*
Habitat in Cayenna. D. v. Rohr.

7. C. cinereo fuſcoque varius, coleoptris punctatis poſtice gibbis: tuberculis octo. Ent. ſyſt. 2. 453. 248. *ſtuberculatus.*
Habitat in Cayenna. D. v. Rohr.

8. C. capite thoraceque obſcure ferrugineis, elytris fuſcis: punctis maculisque albis. Ent. ſyſt. 2. 453. 249.* *ferruus.*
Habitat in Cayenna. D. v. Rohr.
Simillimus in Muſeo D. Lund, totus ater, punctis elytrorum albidis, vix diſtinctus.

29. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*innocuus.* 29. C. albidus, thorace rugoſo, elytris punctato-ſtriatis.
Habitat in Mogador. D. Schousboe. Muſ. D. de Seheſtedt.
Corpus totum albidum, immaculatum. Thorax rugoſus. Elytra punctis impreſſis ſtriata, nec ſpinoſa, nec acuminata.

*modeſtus.* 30. C. cinereus, thorace elytrisque fuſco-maculatis. Ent. ſyſt. 2. 453. 250. *
*Curculio modeſtus.* Oliv. Inſ. 83. fig. 178.
Habitat in noua Zelandia. Muſ. D. Banks.

*flaueſcens.* 31. C. obſcurus, thoracis elytrorumque acuminatorum lateribus flauis. Ent. ſyſt. 2. 454. 251.
Habitat in America meridionali. D. Schreber.

*pollinoſus.* 32. C. ſquamoſo-flaueſcens, elytris acuminatis. Ent. ſyſt. 2. 454. 252. *
Panz. Fn Germ. 19. tab. 4.
Habitat in Auſtria. D. Schneider.

*lateralis.* 33. C. viridi-ſquamoſus, thoracis elytrorumque lateribus flauis. Ent. ſyſt. 2. 454. 253. *
Habitat in India orientali. Muſ. D. Lund.

*viridis.* 34. C. vireſcens, thoracis elytrorumque lateribus flauis. Ent. ſyſt. 2. 454. 254. *
Linn. ſyſt. Nat. 2. 616. 76.
Oliv. Inſ. 83 fig. 18.
Degeer Inſ. 5. 256. 45.
Sulz. Inſ. tab. 3. fig. 24.
Schaeff. Icon. tab. 53. fig. 6.
Payk. Monogr. 107. 103.
Fn. Sv 3. 301. 24.
Bonsd. Curc. 22. 6.
Herbſt. Col. 6. tab. 69. fig. 4.
Habitat in Germaniae Pomona.

35. C.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. C. virescens, thoracis dorso fuluo. Ent. syst. suppl. 171. 255. * *sellatus.*
Habitat in Russia meridionali. Dom. Boeber.

6. C. virescente-albus, elytris basi vnidentatis, postice gibbosis. Ent. syst. 2. 454. 255. * *gibber.*
*Curculio gibber.* Oliv. Inf. 83. fig. 189.
Pallas Icon. tab. B. fig. 54.
Habitat in Cayenna. Mus. Dr. Hunter.

7. C. fuluus, capite basi pedibusque fuscis, rostro emarginato. Ent. syst. 2. 455. 256. * *fuluus.*
Habitat in Saxonia. Dom. Hybner.

8. C. fuscus, thoracis elytrorumque margine cinereo. Ent. syst. 2. 455. 257. * *palliatus.*
Panz. Fn. Germ. 19. tab. 5.
Habitat in Germania. Dom. Schneider.

9. C. corpore ferrugineo, auro-maculato. Ent. syst. 2. 455. 258. * *aurifer.*
*Curculio aurifer.* Oliv. Inf. 83. fig. 124.
Brown. Jamaic. 429. tab. 43. fig. 3.
Drury. Inf. 1. tab. 32. fig. 1.
Habitat in Americae meridionalis Insulis.

10. C. albus, elytris vitta abbreuiata, dentata, pedibusque cyaneis. Ent. syst. 2. 455. 259. * *cyanipes.*
*Curculio cyanipes.* Oliv. Inf. 83. fig. 190.
Habitat in Cayenna. Mus. Dr. Hunter.

11. C. thorace fusco, albo-lineato, elytris cinereis, fusco-fasciatis: margine corporeque viridibus, nitidis. Ent. syst. 2. 456. 260. * *repandus.*
Habitat in Italia. Dom. de Schlangbush.

12. C. viridi-nitens, elytris viridi-ferrugineo, nigro cinereoque variis. Ent. syst. 2. 456. 261. * *Tamarisci*
*Curculio Tamarisci.* Oliv. Inf. 83. fig. 71.
Habitat in Africae Tamarisco. Dom. Prof. Vahl.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſplendidulus.* 43. C. viridi-nitens, coleoptrorum diſco cinereo nigro, faſciato. Ent. ſyſt. 2. 456. 262. *
Habitat in Sibiria. Muſ. D. Banks.

*curuipes.* 44. C. argento-ſquamoſus, tibiis poſticis arcuatis intus dentatis, ciliatis. Ent. ſyſt. 2. 456. 263.
*Curculio curuipes.* Oliv. Inſ 83. fig. 84.
Habitat in India orientali. Muſ. D. Lund.
Femina tibiis poſticis ſimplicibus.

*morbilloſus* 45. C. thorace elytrisque fuſco cinereoque variis punctis eleuatis, atris, numeroſis. Ent. ſyſt. 2. 457. 264.
Habitat in Gallia, Barbaria.

*nebuloſus.* 46. C. canus, elytris faſciis obliquis, nigris. Ent. ſyſt. 2. 457. 265.
Linn. ſyſt. Nat. 2. 617. 84. Fn. Sv. 635.
Geoff. Inſ. 1. 278. 1. tab. 4. fig. 8.
Degeer Inſ. 5. 241. 27.
Sahaeff. Icon. tab. 25. fig. 3.
Friſh. Inſ. 11. 32. tab. 23. fig. 5.
Payk. Monogr. 104. 161.
Fn. Sv. 3. 298. 122.
Bonsd. Curc. ſuec. 20. 3.
Herbſt. Coleopt. 6. tab. 64. fig. 8.
Habitat Larua in foliis Salicis, imago in collibus lapidoſis.

*roridus.* 47. C. oblongus, nigricans, elytris faſciis abbreuiatis punctisque albis. Ent. ſyſt. 2. 457. 266. *
*Curculio roridus.* Pall. Icon. tab. A.
Habitat in Auſtria. Dom. Schneider.

*cinereus.* 48. C. cinereus, thoracis lineis corporeque fuſcis.
*Curculio cinereus.* Schrank. Auſtr.
Habitat in Auſtria. Dom. de Meyerle.
Statura et magnitudo omnino C. nebuloſi. Antennae nigrae, claua apice cineraſcente. Roſtrum

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

breue, craſſum, ſupra cineraſcens: linea media parum eleuata. Thorax laeuis, cinereus: lineis tribus latis, fuſcis. Elytra cinerea, ſutura fuſca. Corpus et pedes fuſca.

9. C. fuſcus, elytris cinereo-variis, roſtro pedibusque nigris. *ſenilis.*

Habitat in Auſtria. Dom. de Meyerle.

Statura praecedentis. Antennae nigrae, claua cineraſcente. Roſtrum breue, craſſum, cylindricum, nigrum. Thorax vix ſcaber, fuſcus: linea laterali lineolisque tribus dorſalibus, obſoletis, cinereis. Elytra laeuia, fuſca, cinereo valde variegata. Corpus fuſcum, pedibus nigris.

0. C. thorace nigro, ſcabro, albo-lineato, elytris albis: maculis nigris, ſparſis, ſubquadratis. Ent. ſyſt. 2. 458. 267. *marmoratus.*

Habitat in Germaniae lapidoſis.

1. C. oblongus, cinereus, ſubnebuloſus, roſtro trifulcato. Ent. ſyſt. 2. 458. 268. *ſulciroſtris*

Linn. ſyſt. Nat. 2. 617. 85.
Degeer Inſ. 5. 240. 26.
Oliv. Inſ. 83. fig. 24.
Schaeff. Icon. tab. 25. fig. 111.
Payk. Monogr. 103. 100.
Fn. Sv. 3. 297. 121.
Bonsd. Curc. ſuec. 21. 4.

Habitat in Europae plantis.

2. C. nigricans, cinereo-varius, capite retuſo, roſtroque carinato. Ent. ſyſt. 2. 458. 269. * *porculus.*

*Curculio porculus.* Act. ſuec. 6. tab. 2. fig. 20.

Habitat in Hungaria. Dom. Hybner.

3. C. cinereus, thorace antice vtrinque retuſo, elytris faſcia fuſca, pedibus anticis elongatis. *retuſus.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Guinea. Dom. Meier. Muſ. D. de Seheſtedt.

Statura magna C. emarginati. Roſtrum planum cinereum, biſulcatum. Thorax cinereus, dorſo obſcuro, antice vtrinque valde retuſo. Elytra ſtriata, cinerea: faſcia pone medium nigra. Corpus cinereum, pedibus anticis elongatis.

*emarginatus.* 54. C. oblongus, fuſcus, coleoptrorum dorſo cinereo lineis duabus punctorum impreſſorum atrorum. Ent. ſyſt. 2. 458. 270. *

Habitat Halae Saxonum. Dom. Hybner.

*perlatus.* 55. C. niger, abdomine albo: punctis eleuatis, glabris, atris, roſtro ſulcato. Ent. ſyſt. 2. 459. 271. *

Habitat in China. Muſ. D. de Seheſtedt.

*diſtinctus.* 56. C. griſeus, elytris punctis duobus diſtinctis albidis: anteriore didymo. Ent. ſyſt. 2. 459. 272.

Habitat in Hungaria. Dom. Hybner.

*glaucus.* 57. C. roſtro carinato, thoraceque inaequali obſcuris, elytris glaucis: puncto poſtico eleuato. Ent. ſyſt. 2. 459. 273.

Panz. Naturf. 24. 22. 30. tab. 1. fig. 30.
Fn. Germ. 19. tab. 6.

Habitat Kiliae. D. Daldorff.

*obliquus.* 58. C. cinereus, elytrorum dorſo albo: ſtrigis obliquis punctoque poſtico fornicato nigris. Ent. ſyſt. 2. 460. 274.

Habitat in Germania.

*Os* maxillis palpisque. *Palpi* quatuor ſubaequales, breuiſſimi; *anteriores* vix longiores, quadriarticulati: articulis tribus prioribus, breuiſſimis, aequalibus, rotundatis, quarto longiori ſubulato, adhaerentes maxillae dorſo; *po-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*riores* vix breuiores, triarticulati, adnati ligulae baſi. *Mandibula* breuis, lata, fornicata, incurua, apice tridentata. *Maxilla* breuiſſima, cornea, integra, apice latior, truncata, ciliata. *Ligula* breuiſſima, membranacea, integra. *Labium* breue, transuerſum, corneum.

9. C. oblongus, fuſcus, elytris albidis: faſcia media lituraque baſeos apicisque fuſcis. Ent. ſyſt. 2. 460. 275. *albidus.*

*Curculio niueus.* Bonsd. Curc. ſuec. 21. 5.
*Curculio candidus.* Herbſt. Arch. 33. 75. tab. 24. fig. 31.
Panz. Fn. Germ. 19. tab. 7.
Payk. Monogr. 106. 102.
Fn. Sv. 3. 300. 123.
Herbſt. Col. 6. 80. 42.
Habitat in Germaniae plantis.

II. 60

10. C. oblongus, fuſcus, villis cinereis, maculatus. Ent. ſyſt. 2. 460. 276.* *diſſar.*

Habitat in India orientali. Prof. Abildgaard.

11. C. thorace tuberculis duobus compreſſis, elytris ſtriatis tuberculatisque apice cinereis. *bicriſtatus*

Habitat in noua Cambria. Dom. Billardiere.
Statura et magnitudo C. crenulati. Caput cinereum, roſtro breui, fuſco. Thorax punctatus, fuſcus, lateribus parum cineraſcentibus, criſtis duabus eleuatis, compreſſis, erectis. Elytra ſtriata: ſtriis punctatis, fuſca, apice cinerea, tuberculo duplici pari ad baſin, ſolitarioque vno ante apicem et altero in apice cinereo. Corpus cinereum, femoribus poſticis annulo fuſco.

12. C. griſeus, elytris rugoſis, nigris, cinereo-maculatis, roſtro atro. Ent. ſyſt. 2. 460. 477.* *Vetula.*

Habitat Tranquebariae. Muſ. D. Lund.

63.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*luſitanicus* 63. C. oblongus, ſupra fuſcus, elytris macula baſeos alba. Ent. ſyſt. 2. 460. 278. *
*Curculio luſitanicus*. Oliv. Inſ. 83. tab. 16. fig. 202.
Habitat in Luſitania. Dr. Gray.

*crenulatus*. 64. C. cinereus, elytris lineis eleuatis tribus crenulatis, atris. Ent. ſyſt. 2. 461. 279. *
*Curculio crenulatus*. Oliv. Inſ. 83. fig. 729.
Habitat in noua Hollandia. Muſ. D. Banks.

*punctatulus*. 65. C. niger, elytris punctis nigris albisque alternis holoſericeis, ſtriatis. Ent. ſyſt. 2. 461. 280. Oliv. Inſ. 83. fig. 119.
Habitat in Terra noua Americes. Muſ. D. Banks

*incanus*. 66. C. oblongus, fuſcus, thoracis dorſo plano. Ent. ſyſt. 2. 461. 281.
Linn. ſyſt. Nat. 2. 610. 81. Fn. Sv. 631.
Geoff. Inſ. 1. 282. 10.
Panz. Fn. Germ. 19. tab. 8.
Bonsd. Curc. ſuec. 41. 36.
Degeer Inſ. 5. 242. 28.
Payk. Monogr. 102. 99.
Fn. Sv. 3. 296. 120.
Herbſt. Col. 6. tab. 77. fig. 3.
Habitat in Europae plantis.

*articulatus* 67. C. ater, thorace punctato, roſtro baſi articulato. Ent. ſyſt. 2. 461. 282.
Habitat in Auſtria. Dom. Schneider.

*coſtatus*. 68. C. cinereus, thorace nigro: lineis quatuor cinereis. Ent. ſyſt. 2. 462. 283. *
Habitat in Gallia auſtraliori. Muſ. D. Banks.

*melancholicus*. 69. C. fuſcus, elytris tibiisque teſtaceis. Ent. ſyſt. 2. 462. 284.
Habitat in Germania. D. Smidt.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

C. obſcurus, coleoptrorum faſcia interrupta alba. Ent. ſyſt. 2. 462. 285. * *interruptus.*

*Curculio interruptus.* Oliv. Inſ. 83. fig. 122.
Habitat in noua Hollandia. Muſ. D. Banks.

C. ater, elytris crenato-ſtriatis, pedibus anticis elongatis. Ent. ſyſt. 2. 462. 286. * *ſcutellaris.*

*Curculio ſcutellaris.* Oliv. Inſ. 83. fig. 142.
Habitat in Terra Diemenii. Muſ. D. Banks.

C. fuſcus, thoracis margine punctisque duobus ferrugineis, pedibus anticis elongatis. Ent. ſyſt. 2. 462. 287. * *longimanus.*

*Curculio longimanus.* Oliv. Inſ. 83. fig. 114.
Habitat in Braſilia. Muſ. D. Banks.

C. cinereus, fuſco-irroratus, thorace acute ſpinoſo. Ent. ſyſt. 2. 463. 288. * *ſpinifex.*

*Curculio Inderienſis.* Pall. Icon. tab. B. fig. 5.
Habitat in Ruſſia meridionali. Dr. Schulze.

C. cinereus, thorace triſulcato, elytris gibbis, ſtriatis, multituberculatis. *Echinus.*

Habitat in Guinea. D. Meier. Muſ. D. Lund.
Magnitudo C. ſpinificis. Antennae piloſae, cinereae, claua parua, fuſca. Thorax planus, ſulcis tribus abbreuiatis, impreſſus. Elytra gibba, ſtriata, tuberculis plurimis, eleuatis; poſticis duobus maioribus. Corpus cinereum.

C. fuſco-aureus, pedibus ferrugineis. Ent. ſyſt. 2. 463. 289. * *micans.*

*Curculio Pyri.* Payk. Monogr. 85. 82.
*Curculio micans.* Payk. Fn. Sv. 3. 302. 125.
Bonsd. Curc. 24. 8.
Herbſt. Col. 6. tab. 78. fig. 3.
Panz. Fn. Germ. 19. tab. 9.
Schaeff. Icon. tab. 2. fig. 11.
Habitat in Daniae hortis. Muſ. D. de Seheſtedt.

II. 61.

76.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*murinus* 76. C. fuſcus, thorace trilineato, elytris pallido-ſtri tis nigroque punctatis. Ent. ſyſt. 2. 463. 29
Habitat in Germania. D. Smidt.

*Polygoni.* 77. C. thorace lineato, elytris cinereis: linea fuſcis ſuturaque nigro-punctata. Ent. ſyſt. 464. 291. *
Linn. ſyſt. Nat. 2. 609. 26. Fn. Sv. 597.
Oliv. Inſ. 83. fig. 41.
II. 61. Panz. Fn. Germ. 19. tab. 10.
Payk. Monogr. 42. 39. β.
Habitat in Europae Polygono.

*dilatatus.* 78. C. cinereus, thorace inaequali marginato, el tris ſulcatis, poſtice vnidentatis.
Habitat in Tanger. D. Schousboe. Muſ. D. Lar
Statura fere Opatri. Roſtrum breue, valde ſulca tum. Thorax cinereus, linea antica transuerſa lineolisque duabus transuerſis incuruis elev tis margineque dilatato. Elytra cinerea, ſulca ta: ſulcis ſtriis duabus punctorum impreſſ rum. Stria eleuata intermedia, abbreuiata p ſtice dentem format. Corpus cinereum.

*tomentoſus* 79. C. piloſus, obſcurus, roſtro plano, elytris pun ctato-ſtriatis.
Habitat in Guinea. D Meier. Muſ. D. de Schefer
Statura et magnitudo C. griſei. Antennae cin reae, claua parua, fuſca. Roſtrum breue, p num, obſcurum, villoſum. Thorax planus, v loſus, obſcurus, dorſo obſolete biſulcato. El tra hirta, obſcura, ſtriato-punctata, apice i acuminata. Corpus cinereum.

*griſeus.* 80. C. ſupra griſeo-fuſcus, ſubtus cinereus, ro canaliculato. Ent. ſyſt. 2. 464. 292. *
Habitat in Anglia et Italia.

*fritillum.* 81. C. griſeus, elytris nigro-punctatis. Ent. ſ
II. 64. ſuppl. 175. 292.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio fritillum.* Panz. Fn. Germ. 18. tab. 19.
Habitat in Germania. Dr. Panzer.

82. C. obſcure brunneus, thorace lineis duabus, elytris macula laterali vireſcentibus, pedibus anticis longioribus. *rubiginosus.*

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo C. incani. Antennae longitudine fere corporis nigrae. Roſtrum breue, planum. Thorax brunneus, lineis duabus latis, vireſcentibus. Elytra ſtriata, obſcure brunnea, litura parua in ipſa baſi maculaque oblonga in medio marginis exterioris vireſcentibus. Corpus cinereum. Pedes nigri, femoribus clauatis, antici longiores.

83. C. nigricans, elytris griſeis: maculis duabus albis; poſteriore maiore communi. Ent. ſyſt. 2. 464. 293. * *3guttatus.*

Petiv. Gazoph. tab. 50. fig. 8.
Schaeff. Icon. tab. 43. fig. 9.
Sulz. Inſ. tab. 4. fig. 11.
Habitat in Anglia.

84. C. faſciis, thorace elytrisque ſcabris cinereo-lineatis, elytris ante apicem vnidentatis. *Meyerlaei.*

Habitat in Auſtria. D. Scheidler.

Statura C. moerentis, at maior. Roſtrum fuſcum, ſubcarinatum. Thorax rotundatus, punctis eleuatis, ſcaber, fuſcus, lineis tribus laeuibus, cinereis. Elytra ſcabra, fuſca, lineis tribus laeuibus, cineraſcentibus, ante apicem decliuia, dente in medio vnico, eleuato. Pedes nigri.

85. C. luteſcens, thorace lineis duabus dorſalibus fuſcis. Ent. ſyſt. 2. 464. 294. *Arundinis*

*Curculio luteus.* Knoch. Beytr. 1.

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 19. tab. 11.
Habitat in Germaniae Arundine. D. Prof. Helwi

*planatus.* 86. C. obſcurus, elytris rugoſo-ſtriatis, femoribu clauatis rufis.
Habitat in Inſula Chorſo. D. de Meyerle.
Nimis affinis videtur C. 2ſulcato, at diſtinctus duplo fere maior. Caput atrum, roſtro breui craſſo dorſo biſulcato, villoſo. Thorax obſcu rus, vix ſcaber. Elytra rugoſa, ſtriata, obſcu ra, feminae depreſſa, plana, lateribus griſeis Pedes nigri, femoribus clauatis, rufis.

*Goerzenſis* 87. C. obſcurus, ſcaber, femoribus omnibus clauatis rufis.
Habitat in Auſtria. D. de Meyerle.
Statura omnino C. ſcabrati, at duplo maior. Caput atrum, roſtro breui, craſſo, biſulcato. Thorax et elytra obſcura, punctis eleuatis numeroſiſſimis, atris, ſcabra. Pedes nigri, femoribus clauatis, rufis.

*ſcabratus.* 88. C. cinereus, punctis eleuatis atris.
Habitat in noua Cambria. D. Billardiere.
Statura C. geminati, at duplo maior. Antennae nigrae, articulo primo valde elongato, incuruo. Caput et thorax cinerea, punctis plurimis eleuatis, atris. Elytra ſtriata, cinerea, punctis plurimis eleuatis, ſparſis, atris. Pedes cinerei, femoribus clauatis, inermibus.

*ſulphurifer* 89. C. ſcaber, cinereus, flauo-puluerulentus, pedibus nigris.
*Curculio ſulphurifer.* Herbſt. Col. 6.
Habitat in Auſtria. D. de Meyerle.
Statura omnino C. Goerzenſis, at paullo minor. Corpus totum punctis paruis, eleuatis ſcabrum, cinereum, puluere flaueſcente adſperſum. Pedes omnes nigri, femoribus clauatis.

90.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

1. C. laeuis, ater, immaculatus. *glabratus.*
*Curculio dignus.* Herbſt. Col. 6. *dirus*
Habitat in Auſtria. D. de Meyerle.
Corpus magnum. atrum, glabrum, omnino immaculatum. Roſtrum breue, craſſum, lineola parua, vtrinque impreſſa.

1. C. ater, thorace ſcabro, elytris punctato-ſtriatis. *orbicularis.*
*Curculio orbicularis.* Herbſt. Col. 6.
Habitat in Auſtria. D. de Meyerle.
Statura omnino praecedentium, totus ater. Thorax punctis elevatis, ſcaber. Elytra punctato-ſtriata, nullo modo acuminata.

2. C. ſupra fuſco-cinereus, ſubtus niger, roſtro ſubſulcato. Ent. ſyſt. 2. 465. 295. * *moerens.*
Habitat Halae Saxonum. Dom. Hybner.

3. C. griſeus, capite thoraceque nigris: linea dorſali alba. Ent. ſyſt. 2. 465. 296. * *griſſonus.*
Habitat in Italia. D. de Schlanbuſch.

4. C. cinereus, thorace trilineato, elytris ſtriis numeroſis fuſcis, per paria approximatis. Ent. ſyſt. 2. 465. 297. * *geminatus*
Habitat Kiliae. D. Daldorff.

5. C. niger, elytris ſtriatis, antennis pedibusque ferrugineis. Ent. ſyſt. 2. 465. 298. * *gramineus*
Habitat in Germaniae graminoſis. D. Smidt.

6. C. flaueſcente villoſus, roſtro apice retuſo atro. Ent. ſyſt. 2. 465. 299. * *diadema.*
Habitat in Cayenna. D. v. Rohr.

7. C. griſeus, thoracis dorſo fuſco: linea albida, elytris nigro-punctatis: ſutura ante apicem albida. Ent. ſyſt. 2. 466. 300. * *Meles.*
Habitat in Germania. D. Smidt.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Coryli.* 98. C. cinereo-fuſcoque varius, elytrorum futura dimidiata atra Ent. ſyſt. 2. 466. 301. *
II. 62. Panz. Fn. Germ. 19. tab. 12.
Habitat in Angliae Corylo.

*lineatus.* 99. C. fuſcus, thorace ſtriis tribus pallidioribus.
Ent. ſyſt. 2. 466. 302. *
Linn. ſyſt. Nat. 2. 616. 80. Fn. Sv. 630.
Ent. Heluet. 1. 80. 1. tab. 6. fig. 1. 2.
Geoff. Inſ. 1. 283.
Degeer Inſ. 5. 247. 35.
Schaeff. Icon. tab. 103. fig. 8.
Payk. Monogr. 112. 108.
Fn. Sv. 3. 307. 130.
Bensd. Curcul. 30. 16.
Herbſt. Col. 6. 497. 527.
Habitat in Europae plantis.

*lunatus.* 100. C. coleoptris eleuato-ſtriatis: faſcia ſemicirculari maculaque poſtica albis; lunula nigra.
Ent. ſyſt. 2. 466. 303. *
Habitat in Anglia in arenoſis Richmondenſibus.

*caninus.* 101. C. fuſcus, thorace, lineis elytrisque brunneis.
Ent. ſyſt. 2. 467. 304. *
Habitat in Germania. D. Smidt.

*eleuatus.* 102. C. niger, elytris lineis tribus eleuatis rufis: margine atro.
Habitat in Selandia. D. Roſenkilde. Muſ. D. Lund.
Paruus. Caput nigrum, puncto frontali impreſſo. Roſtrum breue, nigrum, antennis ferrugineis. Thorax canaliculatus, niger, dorſo punctis quatuor impreſſis. Elytra ſtriato-punctata, lineis tribus eleuatis rufis: margine atro punctoque atro margine connexo. Corpus cinereum.

*mus.* 103. C. cinereus: dorſo fuſco.
*Curculio cineraſcens.* Ent. ſyſt. 2. 467. 305. *
Ha-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Italia. Dr. Allioni.

104. C. tomentoſo-cineraſcens, pedibus teſtaceis. *fuluipes.*
Ent. ſyſt. 2. 467. 306. *
Payk. Monogr. 114. 109.
Fn. Sv. 3. 310. 133.
Degeer Inſ. 5. 235. 41.
Herbſt. Col. 6. 497. 526.
Habitat Halae Saxonum. Dom. Hybner.

105. C. teſtaceus, capite elytrisque ſtriatis, cinereo- *ruficollis.*
fuſcis. Ent. ſyſt. 2. 467. 307. *
Habitat Halae Saxonum. D. Hybner.

106. C. brunneus, elytris faſciis vndatis cinereis. *fuluicornis*
Ent. ſyſt. 2. 468. 308.
Payk. Monogr. 109. 105.
Fn. Sv. 3. 304. 127.
Bonsdorff. Curcul. 27. 13.
Herbſt. Col. 6. tab. 65. fig. 13.
Habitat in Europae arboribus.

107. C. ſubhiſpidus, cinereus, elytris punctato- *echinatus.*
ſtriatis.
Habitat in noua Cambria. Muſ. D. Billardiere.
Maior ſequentibus. Caput cinereum, roſtro breui apice fuſco. Thorax nullo modo canaliculatus, at punctis latis approximatis, cinereis. Elytra punctato-ſtriata. Pedes cinerei, femoribus clauatis.

108. C. cinereus, piloſus, antennis nigris. Ent. ſyſt. *piloſus.*
2. 468. 309. *
Habitat in Germania. D. Schneider.

109. C. fuſcus, elytris apice pallidis: ſtriga vndata *vndatus.*
fuſca, Ent. ſyſt. 2. 468. 310. *
*Curculio tereticollis.* Bonsd. Curc. 28. 14.
Degeer Inſ. 5. 246. 33.

Payk.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Monogr. 108. 104.
Fn. Sv. 3. 303. 126.
Herbſt. Col. 6. 499. 529.
Habitat in Europae Corylo Auellana. Dom. de Hattorff.

*lanatus.* 110. C. piloſus, cinereus, elytris ſtriatis: faſciis duabus abbreuiatis albis.
Habitat in noua Cambria. Muſ. D. Billardiere.
Statura parua praecedentium. Roſtrum breue, canaliculatum, cinereum. Thorax globoſus, laeuis, cinereus, immaculatus. Elytra piloſa, punctato-ſtriata, cinerea, faſciis duabus, in medio interruptis, albis. Pedes cinerei.

*hiſpidulus.* 111. C. fuſcus, thorace cinereo-lineato, elytris hiſpidis: punctis obſcurioribus ſtriatis. Ent. ſyſt. 2. 468. 311. *
Payk. Monogr. 110. 106.
Fn. Sv. 3. 305. 128.
Herbſt. Col. 6. tab. 87. fig. 14.
Habitat in Kiliae plantis aquaticis. Muſ. D. Lund.

*hirſutulus.* 112. C. cinereus, hiſpidus, elytris ſtriatis, antennis pedibusque flaueſcentibus. Ent. ſyſt. 2. 468. 312. *
*Curculio echinatus.* Bonsd. Curc. 33. 21.
Payk. Monogr. 92. 89.
*1.48.* Fn. Sv 3. 286. 109.
Herbſt. Col. 6. tab. 87. fig. 11.
Habitat in Germania. D. Smidt.

*holoſericeus.* 113. C. teſtaceus, elytris holoſericeis punctato-ſtriatis.
Habitat in Auſtria. D. Scheidler.
Paruus. Caput et thorax obſcure teſtacea. Elytra punctata, dilutiora, holoſericea. Pedes teſtacei.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

114. C. cinereus, capite thoraceque canaliculatis, elytris hiſpidis. Ent. ſyſt. 2. 469. 313. *ſcabriculus.*
Linn. Mant. 91. 88.
Bonsd. Curc. 32. 20.
Payk. Monogr. 91. 88.
Fn. Sv. 3. 285. 108.
Herbſt. Col. 6. tab. 78. fig. 10. tab. 77. fig. 14.
Habitat in Sueciae arenoſis.

115. C. obſcurus, thorace ſcabro, elytris piloſis ſulcatis, pedibus rubris. *ſetoſus.*
*Curculio ſcaber* longiroſtris, cinereus, pedibus rufis, elytris ſcabris. Linn. ſyſt. Nat. 2. 609. 22. Fn. Fv. 592.
Habitat in Suecia. Muſ. D. Lund.
Statura et magnitudo omnino C ceruini. Roſtrum breue, obſcurum, antennis rufeſcentibus. Thorax globoſus, fuſcus, punctis eleuatis ſcaber. Elytra hiſpida, ſulcata: ſulcis ſtriis duabus punctorum impreſſorum. Pedes rufi.

116. C. cinereus, capite thoraceque laeuibus, elytris ſubſtriatis, hiſpidis. *ſquamulatus.*
*Curculio ſquamulatus.* Herbſt. Col. t. 87. f. 12.
Habitat in Germania.
Nimis affinis videtur C. ſcrabriculo, at duplo fere minor et caput et thorax laud impreſſa.

117. C. ater, hiſpidus, antennis pedibusque rufis. *Erinaceus.*
Habitat in Auſtria. Dom. de Meyerle.
Statura et magnitudo omnino C. ſcrabiculi, at totus ater, pllis cinereis hiſpidus. Elytra crenato-ſtriata. Antennae et pedes rufi.

118. C. ater, elytris vitta marginali ſcutelloque inauratis. Ent. ſyſt. 2. 469. 314. *limbatus.*
*Curculio lateralis.* Payk. Monogr. 95. 92.
Fn. Sv. 3. 288. 111.
Habitat in Europa.

119.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*paruulus.* 119. C. viridis, antennis tibiisque teſtaceis. Ent. ſyſt. 2. 469. 315. *
Habitat in Italia. Dr. Allioni.

*viridicollis* 120. C. thorace viridi-ſquamoſo, elytris ſtriatis, nigris. Ent. ſyſt. 2. 469. 316.
Panz. Fn. Germ. 19. tab. 13.
Payk. Fn. Sv. 3. 309. 131.
Herbſt. Col. 6. tab. 75. fig. 12.
Habitat in Germaniae Quercu. Dom. Prof. Helwig

*ceruinus.* 121. C. griſeus, elytris puncto obſoleto baſeos albo, antennis baſi rufeſcentibus. Ent. ſyſt. 2. 470. 317. *
Linn. ſyſt. Nat. 2. 615. 70. Fn. Sv. 627.
Degeer Inſ. 5. 220. 13.
Habitat in Europae plantis.

*ſtriatus.* 122. C. fuſcus, elytris ſtriis cinereis, nigro-punctis. Ent. ſyſt. 2. 470. 318. *
Habitat in Barbaria. Dom. Prof. Vahl.

*lacerta.* 123 C. griſeus, elytris ſtriatis, antennis apice nigris. Ent. ſyſt. 2. 470. 319. *
Habitat in India orientali. Muſ. D. Banks.

*velox.* 124. C. piloſus, niger, elytris margine puncto baſeos, faſcia interrupta punctoque apicis albis. Ent. ſyſt. 2. 470. 320. *
Habitat in Cayenna. Dom. v. Rohr.

*ſeminulum* 125. C. niger, immaculatus, elytris ouatis, puncto-ſtriatis. Ent. ſyſt. 2. 470. 321. *
Habitat in Hungaria. Muſ. Dom. Hybner.

*ſinuatus.* 126. C. cineraſcens, elytris faſcia ſinuata, fuſca, antennis pedibusque teſtaceis.
Habitat in Auſtria. Muſ. Dom. de Meyerle.
Minutus. Antennae teſtaceae. Caput cinereum. Thorax cinereus, dorſo fuſco. Elytra ſtriata

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

villoſa, cinerea, dorſo ad ſuturam faſciaque media, valde ſinuata fuſcis. Pedes omnes teſtacei.

27. C. ater, antennis rufis, thorace vtrinque bituberculato. Ent. ſyſt. 2 471. 322. *ruficornis.*
Linn. ſyſt. Nat. 2. 616. 78. Fn. Sv. 636.
Habitat in Europae plantis.

28. C. niger, obſcurus, elytris punctis plurimis griſeo-albis. Ent. ſyſt. 2. 471. 325. * *multipunctatus.*
Habitat Halae Saxonum. Dom. Hybner.

29. C. niger, elytris ſulcatis cinereis. Ent. ſyſt. 2. 471. 326. * *triſtis.*
Habitat in Suecia et Anglia.

30. C. niger, elytris ſtriatis fuſcis, cinereo-maculatis. Ent. ſyſt. 2. 472. 327. * *raucus.*
*Curculio triſtis.* Bonsd. Curc. ſuec. 36. 27.
Payk. Monogr. 97. 94.
Fn. Sv. 3. 291. 114.
Herbſt. Col. 6. tab. 87. fig. 2.
Habitat in Europa boreali.

31. C. ater, elytris poſtice albo-maculatis. Ent. ſyſt. 2. 472. 328. * *adſperſus.*
*Curculio adſperſus.* Oliv. Inſ. 83. fig. 156.
Habitat in noua Hollandia. Muſ. D. Banks.

32. C. niger, coleoptris ſtriatis fuſcis: ſtriga baſeos niuea. Ent. ſyſt. ſuppl. 177. 328. * *ſtriga.*
Habitat in Isle de France. D. Daldorff.

33. C. fuſcus, elytris punctis eleuatis holoſericeis. Ent. ſyſt. 2. 472. 329. * *punctatus.*
Payk. Monogr. 111. 107.
Fn. Sv. 3. 306. 129.
Herbſt. Col. 6. 505. 540.
Habitat in Suecia.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*obſcurus.* 134. C. ouatus, ferrugineo - fuſcus. Ent. ſyſt. 472. 330. *

*Curculio nitidus.* Bonsd. Curc. ſuec. 37. 8
Payk. Monogr. 98. 95.
Fn. Sv. 3. 292. 116.
Herbſt. Col. 6. tab. 87. fig. 3.
Habitat in Suecia.

*Mercurialis.* 135. C. ouatus, fuſcus, elytrorum ſutura poſt eleuata compreſſa.

Habitat in Selandiae Mercuriali. Muſ. D. Lun
Statura et ſumma affinitas C. obſcuri, at pa minor. Thorax fuſcus, linea dorſali parum e uata. Elytra haud punctata, at lineis alig eleuatis coſtata et ſutura poſtice eleuata, co preſſa. Corpus fuſcum, femoribus clauatis.

*maxilloſus* 136. C. ouatus, niger, mandibulis exſertis arcua Ent. ſyſt. 2. 472. 331. *

Habitat in Hungaria. Dom. Hybner.

*niger.* 137. C. ouatus, ſcaber, niger, pedibus rufis. E ſyſt. 2. 473. 332. *

*Curculio niger.* Ent. Heluet. 1. 88. 5. tab. fig. 1. 2.
*Curculio clauipes.* Bonsd. Curcul. ſuec. 40.
Payk. Monogr. 101. 198.
Fn. Sv. 3. 295. 119.
Herbſt. Col. 6. 332. 306.
Habitat in Germania.

*elev coſtatus.* 138. C. niger, thorace ſcabro, elytris ſulcatis: ſ cis punctis impreſſis, duplici ſerie.

Habitat in Auſtria. D. de Scheidler.
Paullo minor, quam C. niger. Roſtrum bre laeue. Thorax punctis eleuatis ſcaber. E lineis tribus valde eleuatis et ſulcis pun duplici ſerie impreſſis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

39. C. ater, nitidus, roſtro emarginato. Ent. ſyſt. 2. 473. 333. * *laeuigatus.*

Habitat Kiliae ſub lapidibus.

40. C. niger, elytris punctato-ſtriatis; antennis tibiisque rufis. Ent. ſyſt. 2. 473. 334. * *rotundatus.*

Habitat in Germania. D. de Leuwenſkiold.

41. C. niger, thorace carinato-varioloſo, elytris ſtriatis. Ent. ſyſt. 2. 473. 335 * *varioloſus.*

*Curculio ſenex* Roſſ. Fn. Etr. 1. 337.

Habitat in Saxonia.

42. C. thorace elytrisque varioloſis nigris, albo-punctatis. Ent. ſyſt. 2. 473. 336. * *ſticticus.*

Habitat ad Cap. Bon. Spei.

43. C. obſcurus, niger, thorace ſubpunctato, elytris punctato-ſtriatis. *affinis.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Selheſtedt.

Statura omnino C. valgi, at paullo maior. Caput nigrum, antennarum claua acuta. Thorax ſubpunctatus, niger, obſcurus. Elytra nigra, obſcura, punctato-ſtriata, ſtriis ante apicem coeuntibus. Pedes nigri, tibiis poſticis ſimplicibus.

44. C. ater, nitens, elytris punctato-ſtriatis, femoribus annulo cinereo. Ent. ſyſt. 2. 474. 337.* *valgus.*

Habitat in America.

Affinis certe C. curuipedis, et forte mera eius varietas.

45. C. ater, elytris margine lineolisque duabus albis. Ent. ſyſt. 2. 474. 338. * *ſuccinctus.*

Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

46. C. vireſcens, thoracis dorſo elytrorumque ſutura vittaque dorſali obſcurioribus, roſtro emarginato. *exciſus.*



Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Moldauia. Muſ. D. Lund.
Magnitudo C. valgi. Roſtrum breue, dorſo pl
obſcuriore, apice emarginato. Thorax fla
virens, dorſo late obſcuriore. Elytra ſubſtri
flauo-vireſcentia, ſutura vittaque media
ſali obſcurioribus, apice acuminata. Co
vireſcens.

*Cameleon.* 147. C. aeneus, elytris ſutura vittaque abbreui
viridi-aureis, Ent. ſyſt. 2. 474. 339.
*Curculio Cameleon.* Oliv. Inſ. 83. fig. 166.
Drury Inſ. 2. tab. 33. fig. 4. 5.
Habitat in Iamaica. Muſ. Dr. Hunter.

*vittatus.* 148. C. ater, elytris ſtriis albis rubrisque abbre
tis. Ent. ſyſt. 2. 474. 340.
Linn. ſyſt. Nat. 2. 610. 33.
Oliv. Inſ. 83. fig. 192.
Sloan. Iamaic. 2. 210.
Habitat in Americae Inſulis. Deſtruit larua n
cis Citri medicae. Dr. Schwarz.

*Spengleri.* 149. C. elytris flauis: lineis glaberrimis abbre
tis atris. Ent. ſyſt. 2. 475. 341.
Linn. ſyſt. Nat. 2. 609. 32.
Oliv. Inſ. 83. fig. 82.
Act. Berol. 4. tab. 7. fig. 7.
Habitat in Americae Inſulis.

*biuittatus.* 150. C. niger, elytris punctato-ſtriatis: vitta e
ginali dorſalique interrupta flauis. Ent. ſyſ
475. 342.
Oliv. Inſ. 83. fig. 23.
Habitat in Americae Inſulis. Dr. Pflug.

*liuidus.* 151. C. griſeus, thoraçe elytrisque cinereo atro
maculatis. Ent. ſyſt. 2. 475. 343.
*Curculio liuidus.* Oliv. Inſ. 83. fig. 59.
Brown. Illuſtr. tab. 49. fig. 8.

Spe

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Sparm. Act. Holm. 1785. tab. 2. fig. 17.
Habitat in America meridionali.

52. C. coleoptris atris: ſtriis ſex dorſalibus approximatis, abbreuiatis albis. Ent. ſyſt. 2. 476. 344. * *pulcher.*

*Curculio pulcher.* Oliv. Inſ. 83. fig. 130.
Brown. Illuſtr. tab. 49 fig. 6.
Habitat in Iamaica. Muſ. D. Banks.

53. C. elytris flaueſcentibus: ſtriis abbreuiatis nigris, marginalique ſanguinea. Ent. ſyſt. 2. 476. 345. * *feſtiuus.*

Habitat in Americae meridionalis Inſulis. Dr. Iſert.

54. C. griſeus, thorace lineis lateralibus fuluis, elytris albis, maculis atris fuluisque. Ent. ſyſt. 2. 476. 346. * *biſtrioui-cus.*

Habitat ad Cap. Bon. Spei. Muſ. D. Lee.

55. C. elytris acuminatis griſeis: margine baſeos flaueſcente. Ent. ſyſt. 2. 476. 347. * *Rohrii.*

Habitat in Americae Inſulis. D. v. Rohr.

56. C. niger, thorace elytrisque punctis impreſſis albis. Ent. ſyſt. 2. 477. 348. * *impreſſus.*

*Curculio impreſſus.* Oliv. Inſ. 83. fig. 126.
Habitat in Iamaica. Muſ. D. Banks.

57. C. niger, thorace angulato, elytris tuberculatis baſi apiceque albidis. Ent. ſyſt. 2. 477. 349. * *albipes.*

*Curculio albipes.* Oliv. Inſ. 83. fig. 109.
Habitat in Madagaſcar. Muſ. Gallic.

58. C. niger, thorace rufo-maculato, elytris lineis tribus impreſſis cinereis. Ent. ſyſt. 2. 477. 350. * *riuuloſus.*

*Curculio riuuloſus.* Oliv. Inſ. 83. tab. 11. f. 133.
Habitat in India orientali.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*depreſſiro-ſtris.* 159. C. fuſco cinereoque varius, roſtro depreſſo pl no, apice atro. Ent. ſyſt. 2. 477. 351. *
Habitat Halae Saxonum. Dom. Hybner.

*marginel-lus.* 160. C. niger, thoracis elytrorumque margine al Ent. ſyſt. 2. 478. 352. *
Habitat in Africa aequinoctiali. Muſ. D. Banks.

*verrucoſus.* 161. C. aeneo niger, eleuato-punctatus, elytris p ſtice verrucoſis. Ent. ſyſt. 2. 478. 353. *
Linn. ſyſt. Nat. 2. 618. 90. Muſ. Lud. Vlr. 6
*Curculio verrucoſus.* Oliv. Inſ. 83. fig. 125.
Drury Inſ. 1. tab. 32. fig. 5.
Habitat ad Cap. Bon. Spei.

*Capenſis.* 162. C. ater, thorace eleuato-punctato, elytris ſtr to-crenatis. Ent. ſyſt. 2. 478. 354.
Linn. ſyſt. Nat. 2. 618. 82. Muſ. Lud. Vlr. 5
*Curculio Capenſis.* Oliv. Inſ. 83. fig. 52.
Habitat ad Cap. Bon. Spei.

*exſertus.* 163. C. ater, mandibulis exſertis, elytris cren ſtriatis, poſtice vnituberculatis.
Habitat in noua Cambria. D. Billardiere.
Statura et magnitudo C. Capenſis. Roſtrum b ue, ſulcatum, mandibulis exſertis, inermib Thorax planiuſculus, ſubpunctatus, ater. Sc tellum eleuatum. Elytra ſtriata: ſtriis punc tis, atra, ante apicem dente eleuato, obt Pedes tibiis anticis compreſſis, vix dentat

*6vittatus.* 164. C. niger, thorace ſcabro albo-lineato, ely acuminatis. Ent. ſyſt. 2. 478. 355. *
*Curculio 6vittatus.* Oliv. Inſ. 83. tab. 12. f. 1
Habitat - - Muſ. Dr. Hunter.

*vitulus.* 165. C. niger, fronte bidentata, elytris vniſpino Ent. ſyſt. 2. 479. 356. *
Habitat in Terra del Fuego. Muſ. D. Banks.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

66. C. griſeus, thorace inaequali, antice prominulo, elytris ſulcatis: poſtice bidentatis, roſtro triſulcato. Ent. ſyſt. 2. 479. 357. * *inaequalis.*

*Curculio inaequalis.* Oliv. Inſ. 83. fig. 164.

Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

67. C. albidus, thorace elytrisque nodulis eleuatis atris numeroſis glabris, elytris apice recuruo acuminatis. *recurvus.*

Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

Statura magna praecedentium. Roſtrum craſſum, albidum, nigro-punctatum: dorſo canaliculato: canali atro. Thorax albidus, nodulis plurimis, approximatis, atris, nitidis. Elytra puncrato-ſtriata: nodulisque plurimis, eleuatis, atris, nitidis, apice in ſpinam validam, acutam, recuruam exeunt. Corpus albidum, nigro-punctatum.

68. C. cylindricus, fuſcus, elytris apice acuminatis. Ent. ſyſt. 2. 479. 358. * *acuminatus.*

*Curculio acuminatus.* Oliv. Inſ. 83. fig. 139.

Habitat in noua Zelandia. Muſ. D. Banks.

69. C. fuſcus, elytris acuminatis, crenato-ſtriatis. roſtro ſulcato. Ent. ſyſt. 2. 479. 359. * *capiſtratus*

Habitat ad Cap. Bon. Spei. D. Prof. Leske.

70. C. niger, thorace elytrisque ſpinoſis, fronte excauata. Ent. ſyſt. 2. 480. 360. * *emeritus.*

Linn. ſyſt. Nat. 2. 617. 87. Muſ. Lud. Vlr. 35.

Habitat in Indiis.

71. C. ater, thorace punctato, elytris ſtriatis tuberculatisque. Ent. ſyſt. 2. 480. 361. * *tuberculatus.*

Habitat in India orientali. D. Prof. Abildgaard.

72. C. thorace mutico ſcabro, elytris lineis tribus eleuatis dentatis. Ent. ſyſt. 2. 480. 362. * *criſpatus.*

*Curculio criſpatus.* Oliv. Inſ. 83. fig. 160.

Habi-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat ad Cap. Bon. ſpei. Muſ. D. Banks.

*cultratus.* 173. C. niger, thorace tuberculis duobus compreſſis elytris tuberculatis, poſtice cinereis. Ent. ſyſt. 2. 480. 363. *

*Curculio cultratus.* Oliv. Inſ. 83. fig. 157.
Habitat in noua Hollandia. Muſ. D. Banks.

*tribulus.* 174. C. cinereus, thorace ſcabro, antice impreſſo elytris ſpinoſis. Ent. ſyſt. 2. 481. 364. *

*Curculio tribulus.* Oliv. Inſ. 83. fig. 161.
Habitat in noua Hollandia. Muſ. D. Banks.

*4dens.* 175. C. cinereus, thorace ſcabro, elytris ſpinoſis ſpinis quatuor, poſticis longioribus. Ent. ſyſt. 2. 482. 365. *

*Curculio 4dens.* Oliv. Inſ. 83. fig. 187.
Habitat in noua Hollandia. Muſ. D. Banks.

*4ſpinoſus.* 176. C. albidus, elytris quadriſpinoſis, roſtro fuſco. Ent. ſyſt. 2. 481. 366. *
Habitat ad Cap. Bon. Spei.

*clauus.* 177. C. albicans, thorace canaliculato, coleoptris ſpinoſis: lineolis tribus baſeos rubris. Ent. ſyſt. 2. 481. 367. *

*Curculio clauus.* Oliv. Inſ. 83. fig. 168.
Habitat in noua Hollandia. Muſ. D. Banks.

*noduloſus.* 178. C. thorace lineis ſex noduloſis, elytris ſpinoſis. Ent. ſyſt. 2. 482. 368. *

*Curculio noduloſus.* Oliv. Inſ. 83. fig. 188.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*rubifer.* 179. C. cinereus, thorace ſcabro, elytris ſpinis equatis ſanguineis. Ent. ſyſt. 2. 482. 369. *

*Curculio rubifer.* Oliv. Inſ. 83. fig. 159.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

*globifer.* 180. C. thorace ſcabro, elytris ſpinoſis, poſtice acuminatis. Ent. ſyſt. 2. 482. 370. *

Cu

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio globifer.* Oliv. Inſ. 83. fig. 135.
Habitat ad Cap. Bon. Spei. Muſ. D. Banks.

181. C. thorace vtrinque ſpinoſo noduloſo, elytrorum ſtriis ſubſerratis, apice acuminatis. *ſpectrum.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.
Magnitudo C. pilularii. Thorax planus, vtrinque ſpinoſus. Elytra griſea, obſolete maculata, ſtriis quatuor eleuatis, ſerratis.

182. C. thorace vtrinque ſpinoſo noduloſo, elytris tuberculato-ſtriatis, apice acuminatis. Ent. ſyſt. 2. 482. 371. * *pilularius.*
Habitat ad Cap. Bon. Spei. Muſ. D. Spengler.

183. C. obſcurus, thorace ſcabro, elytris lineis tribus eleuatis ſpinoſis. Ent. ſyſt. 2. 483. 372. *glandifer.*
Habitat ad Cap. Bon. Spei. Muſ. D. Lund.

** *Femoribus dentatis.*

184. C. femoribus dentatis, corpore viridi-variegato. Ent. ſyſt. 2. 483. 373. * *ſpectabilis.*
*Curculio ſpectabilis.* Oliv. Inſ. 83. fig. 180.
Habitat in noua Hollandia. Muſ. D. Banks.

285. femoribus anticis dentatis, gibbus, nigricans, thorace lineis, elytris ſtrigis duabus lineaque apicis flaueſcentibus. Ent. ſyſt. 2. 483. 374.* *pinguis.*
Habitat in Cayenna. D. v. Rohr.

186. C. femoribus dentatis, cinereus, elytris dentibus tribus apiceque emarginatis. Ent. ſyſt. 2. 483. 375. * *tridens.*
*Curculio tridens.* Oliv. Inſ. 83. fig. 154.
Habitat in noua Zelandia. Muſ. D. Banks.

187. C. femoribus ſubdentatis, ater, thorace elytrisque laeuibus fuſco-maculatis. Ent. ſyſt. 2. 484. 376. * *fuſco-maculatus,*
Habitat in Germania boreali.

188.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Liguſtici.* 188. C. femoribus dentatis, corpore obſcuro. Ent. ſyſt. 2. 484. 377.
Linn. ſyſt. Nat. 2. 615. 68. Fn. Sv. 621.
Oliv. Inſ. 83. fig. 77.
Degeer Inſ. 5. 218. 10.
Schaeff. Icon. tab. 2. fig. 12.
Payk. Monogr. 80. 77.
Fn. Sv. 3. 274. 97.
Herbſt. Col. 6. tab. 86. fig. 8.
Habitat in Liguſtico Leuiſtico.

*Zebra.* 189. C. femoribus dentatis, niger, elytris albo-variis. Ent. ſyſt. 2. 484. 378. *
Payk. Monogr. 83. 80.
Fn. Sv. 3. 278. 101.
Herbſt. Col. 6. tab. 88. fig. 7.
Habitat Halae Saxonum. Dom. Hybner.

*2 notatus.* 190. C. femoribus dentatis, obſcurus, elytris puncto poſtico cinereo. Ent. ſyſt. 2. 484. 379. *
Habitat in Germania. D. Smidt.

*nubilus.* 191. C. femoribus dentatis, griſeus, elytris punctis obſcurioribus numeroſis ſubquadratis. Ent. ſyſt. 2. 485. 380.
Schaeff. Icon. tab. 2. fig. 11.
Payk. Monogr. 79. 76.
Fn. Sv. 3. 273. 96.
Bonsd. Curcul. ſuec. 38. 31.
Herbſt. Col. 6. tab. 86. fig. 10.
Habitat Hamburgi. Dr. Schulz.

*punctum.* 192. C. femoribus anticis acute dentatis, niger, ſcutello niueo. Ent. ſyſt. ſuppl. 172. 381. *
Habitat ad Cap. Bon. Spei. D. Daldorff.

*brunnipes.* 193. C. femoribus anticis dentatis, fuſcus, thorace pedibusque obſcure brunneis. Ent. ſyſt. ſuppl. 172. 381. *
Habitat ad Cap. Bon. Spei. D. Daldorff.

194

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

194. C. femoribus obtuſe-dentatis, niger, elytris cinereo-maculatis, antennarum articulo ſecundo apice prominulo. Ent. ſyſt. ſuppl. 172. 381.* *denticornis.*

Habitat in India orientali. Dom. Dalderſt.

195. C. femoribus dentatis, thorace ſcabro, elytris ſulcatis: ſtriis ſerratis. *auſtriacus.*

Habitat in Auſtria. Dom de Meyerle.

Statura et magnitudo C. nigritae. Caput nigrum, roſtro breui, canaliculato. Thorax rotundatus, ſcaber, obſcurus, immaculatus. Elytra co[illegible]*nnata*, ſulcata, lineis elevatis, ſerratis. Pedes nigri, femoribus clauatis, dentatis.

196. C. femoribus dentatis, niger, obſcurus, thorace ſcabro, elytris crenato-ſtriatis. Ent. ſyſt. 2. 485. 381. * *nigrita.*

Habitat in Italia.

197. C. femoribus dentatis, ater, elytris ſtriatis, ferrugineo-maculatis Ent. ſyſt. 2. 485. 382. * *ſulcatus.*

Payk. Monogr. 81. 78.
Fn. Sv. 3. 275. 98.
Degeer Inſ. 5. 217. 9.
Bonsd. Curc. ſuec. 40. 34.
Herbſt. Col. 6. tab. 87. fig. 5.

Habitat in Saxonia.

198. C. femoribus acute-dentatis, antennis pedibusque rufis. Ent. ſyſt. 2. 485. 383. * *calcaratus*

*Curculio calcaratus.* Ent. Helv. 1. 91. 6. tab. 8. fig. 3. 4.

Habitat in Auſtria. Dom. Schneider.

199. C. femoribus dentatis, elytris punctis viridibus. Ent. ſyſt. 2. 485. 384. * *gemmatus*

*Curculio gemmatus.* Oliv. Inſ. 83. fig. 74.
Naturf. 6. tab. 4.

Habitat in Europae plantis.

200.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſquamiger.* 200. C. femoribus acute-dentatis, niger, elytris viridi-variegatis, pedibus rufis.
Habitat in Auſtria. Dom. Scheidler.
Affinis ſane C. gemmato. Roſtrum breue, apice incraſſatum, antennis nigris: articulo ſecundo longiſſimo, claua apice cinerea. Thorax globoſus, ſcaber. Elytra ſtriata, nigra, hinc vnde viridi-ſquamoſa. Pedes rufi, femoribus acute-dentatis.

*picipes.* 201. C. femoribus dentatis, griſeus, elytris nebuloſis: punctis ſubocellatis, ſtriatis. Ent. ſyſt. 2. 486. 385.
*Curculio notatus.* Bonsd. Curc. ſuec. 39. 33.
Geoff. Inſ. 1. 281. 9.
Payk. Monogr. 100. 97.
Fn. Sv. 3. 293. 117.
Herbſt. Col. 6. tab. 88. fig. 5.
Habitat Kiliae. Muſ. D. Lund.

*Morio.* 202. C. femoribus dentatis, ater, nitens, elytris glabris, roſtro biſulcato. Ent. ſyſt. 2. 486. 386.
Oliv. Inſ. 83. fig. 26.
Payk. Monogr. 99. 96.
Fn. Sv. 3. 294. 118.
Degeer Inſ. 5. tab. 7. fig. 22.
Herbſt. Col. 6. tab. 86. fig. 6.
Habitat in Germania. Dom. de Hattorff.

*biſulcatus.* 203. C. femoribus dentatis, niger, thorace ſcabro, elytris rugoſo-ſtriatis, roſtro biſulcato. Ent. ſyſt. 2. 486. 387. *
Habitat in Italia. Dr. Allioni.

*collaris.* 204. C. femoribus dentatis, thorace cinereo, ſcabro, elytris nigris, roſtro carinato. Ent. ſyſt 2. 486. 388. *
Habitat Kiliae.

205.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

205. C. femoribus dentatis, obſcure-cinereus, fuſco-maculatus, elytris piloſis. *aſperatus.*

*Curculio aſperatus.* Payk. Monogr. 84. 81.

Habitat in Europa boreali. Muſ. D. Lund.

Antennae rufeſcentes. Thorax ſquamoſus, fuſcus, lineis tribus cineraſcentibus. Elytra ſtriata, pilis breuiſſimis denſioribus irrorata, cinerea, fuſco-maculata. Pedes rufeſcentes.

206. C. femoribus dentatis, cinereus, elytris teſſellatis. Ent. ſyſt. 2. 486. 389. * *Iris.*

Habitat Kiliae.

207. C. femoribus acute dentatis, niger, elytris viridi-ſquamoſis, pedibus rufis. Ent. ſyſt. 4. App. 454, 390. * *lepidopterus.*

*Curculio lepidopterus.* Payk. Fn. Sv. 3. 276. 99. *

Herbſt. Col. 6. tab. 88. fig. 8.

Habitat in Auſtria. D. Schneider.

208. C. femoribus anticis acute dentatis, viridis, nigro-irroratus. Ent. ſyſt. ſuppl. 174. 390. * *roralis.*

Habitat in Isle de France.

209. C. femoribus anticis dentatis, vireſcens, elytris punctato-ſtriatis: vitta abbreuiata cyanea. *clarus.*

Habitat in America meridionali. D. Smidt. Muſ. D. de Scheſtedt.

Antennae vireſcentes, claua cinerea. Roſtrum breue, apice emarginatum, cinereum. Thorax parum punctatus, vireſcens. Elytra punctato-ſtriata, vireſcentia: vitta abbreuiata, cyanea, et in hac puncta impreſſa paullo maiora. Pedes ſubcyanei, femoribus anticis elongatis, acute dentatis.

210. C. femoribus dentatis, viridi-aeneus, pedibus rufis. Ent. ſyſt. 2. 487. 390. *Pyri.*

*Cur-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Curculio Pyri.* Linn. ſyſt. Nat. 2. 615. 72. Fn. Sv. 623.
*Curculio aeruginoſus.* Bonsd. Curc. ſuec. 23. 7.
Geoff. Inſ. 1. 282.
Oliv. Inſ. 33. fig. 30.
Degeer Inſ. 5. 246. 34.
Sulz. Inſ. tab. 3. fig. 23.
Payk. Monogr. 88. 85.
Fn. Sv. 3. 280. 103.
Herbſt. Col. 6. tab. 79. fig. 2.
Habitat in Pyro communi frequens.

*Alneti.* 211. C. femoribus dentatis, niger, coeruleo-ſquamoſus, antennis piceis. Ent. ſyſt. 2. 487. 391.
Payk. Fn. Sv. 3. 281. 103. β.
Herbſt. Col. 6. tab. 79. fig. 3.
Habitat in Germaniae Alno.
Praecedentis varietatem credit D. de Paykull, at vix rite.

*veſpertinus.* 212. C. femoribus dentatis, cinereo-ſubuilloſus, ſcutello albo. Ent. ſyſt. 2. 487. 392. *
*Curculio cerinus.* Payk. Monogr. 87. 84.
Fn. Sv. 3. 282. 105.
Habitat in Germania. D. Smidt.

*Mali.* 213. C. femoribus dentatis, pubeſcens, fuſcus, antennis pedibusque teſtaceis. Ent. ſyſt. 2 487. 393. *
*Curculio Padi.* Bonsd. Curc. ſuec. 26. 11.
Payk. Monogr. 86. 83.
Fn. Sv. 3. 281. 104.
Herbſt. Col. 6. tab. 79. fig. 5.
Habitat Lipſiae. D. Prof. Leske.

*pallidus.* 214. C. femoribus dentatis, ſquamoſo-pallidus, antennis tibiisque rufis. Ent. ſyſt. 2. 488. 394. *
Habitat in Germania. D. Smidt.

215.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

215. C. femoribus anticis dentatis, viridi-cinereus, elytris punctato-ſtriatis. Ent. ſyſt. 2. 488. 395. * *arboreti.*

Habitat in Cayenna. D. v. Rohr.

216. C. femoribus clauatis dentatis nigris, viridi-ſquamoſus, antennis ruſis. *Betulae.*

Habitat in Auſtriae Alno. Dom. de Meyerle.

Statura et magnitudo praecedentium. Antennae rufeſcentes. Caput et thorax viridi-ſquamoſa. Elytra ſtriata, villoſa, viridi-ſquamoſa. Pedes rufeſcentes, femoribus valde clauatis: claua nigra.

217. C. femoribus dentatis, cinereus, roſtro antice canaliculato, femoribus poſticis tridentatis. Ent. ſyſt. 2. 488. 396. *dentifer.*

Habitat in India orientali. D. Prof. Abildgaard.

218. C. femoribus dentatis, griſeus: lateribus albis, femoribus intermediis bidentatis. Ent. ſyſt. 2. 488. 397. * *curuicornis.*

Habitat in India orientali. D. Prof. Abildgaard.

219. C. femoribus dentatis, flaueſcens, capite thoraceque trilineatis, coleoptrorum diſco fuſco. Ent. ſyſt. ſuppl. 173 397. * *dorſatus.*

Habitat in India orientali. D. Daldorff.

220. C. femoribus dentatis, corpore viridi-argentato. Ent. ſyſt. 2. 489. 398. * *argentatus*

*Curculio argentatus.* Linn. ſyſt. Nat. 2. 615. 73.
Geoff. Inſ. 1. 293. 38.
Oliv. Inſ. 83. fig. 96.
Degeer Inſ. 5. 219. 12.
Sulz. Inſ. tab. 4. fig. 9.
Payk. Monogr. 89. 86.
Fn. Sv. 3. 281. 106.
Bonsd. Curc. ſuec. 27. 12.

Herbſt.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Col. 6. tab. 79. fig. 4.
Habitat in Europae Betula, Vrtica, Quercu.

*viridanus.* 221. C. femoribus dentatis, viridis, ore antennisque nigris, oculis atris. Ent. ſyſt. 2. 489. 399. *
Habitat Tranquebariae. Dr. Koenig.

*oblongus.* 222. C. femoribus dentatis, antennis, elytris pedibusque ferrugineis. Ent. ſyſt. 2. 489. 400. *
*Curculio oblongus.* Linn. ſyſt. Nat. 2. 615. 21. Fn. Sv. 625.
Panz. Fn. Germ. 19. tab. 15.
Herbſt. Arch. 7. tab. 45. fig. 9.
Payk. Monogr. 90. 87.
Fn. Sv. 3. 284. 107.
Bonsd. Curc. ſuec. 25. 9.
Geoff. Inſ. 1. 294. 39.
Oliv. Inſ. 83. fig. 30.
Habitat in Europae plantis.

*muricatus* 223. C. femoribus dentatis, griſeus, elytris ſtriis eleuatis hiſpidis, antennis rufis. Ent. ſyſt. 2. 489. 401. *
Habitat in Germania. D. Daldorff.

*ouatus.* 224. C. femoribus dentatis, niger, thorace ſcabro. Ent. ſyſt. 2. 490. 402. *
*Curculio ouatus.* Linn. ſyſt. Nat. 2. 615. 69. Fn. Sv. 626.
*Curculio Roſae.* Degeer Inſ. 5. 219. 11.
Payk. Monogr. 82. 79.
Fn. Sv. 3. 277. 100.
Bonsd. Curc. ſuec. 35. 25.
Herbſt. Col. 6. tab. 88. fig. 2.
Habitat in Europae arbuſtis.

*albo-lineatus.* 225. C. femoribus dentatis, cinereus, thorace elytrisque albo-lineatis, roſtro ſubtus atro. Ent. ſyſt. 2. 490. 403. *

Hab-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat Halae Saxonum. Dom Hybner.

226. C. femoribus dentatis, niger, elytris albo-maculatis. Ent. ſyſt. 2. 490. 404. * *Picus.*
Habitat in Germaniae Betula. D. Smidt.

227. C. femoribus dentatis, roſtro elytrisque vnituberculatis. Ent. ſyſt. 2. 490. 405. * *attelaboides.*
*Curculio attelaboides.* Oliv. Inſ. 83. fig. 174.
Habitat in Braſilia. Muſ. D. Banks.

## 163. BRENTVS.

*Palpi* inaequales, filiformes.
*Antennae* moniliformes, inſertae, roſtro recto, porrecto, cylindrico.

* *femoribus ſimplicibus.*

1. B. roſtro longiſſimo, ſubtus barbato, elytris elongato-clauatis. Ent. ſyſt. 2. 491. 103. * *barbicornis.*
*Brentus barbicornis.* Oliv. Inſ 84. tab. 1. fig. 5. tab. 2. fig. 5.
Habitat in noua Zelandia. Muſ. D. Banks.

2. B. thorace ſcabro, integro, elytris apice recurvato-ſpinoſis, antennis filiformibus. *curuidens.*
*Brentus curuidens.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 59. 1. *

*Bren-*

---

*Brenti* corpus elongatum, cylindricum, immarginatum, tardum; *capite* paruo, ouato, antice porrecto in roſtrum longiſſimum, rectum, cylindricum; *oculis* paruis, globoſis, prominulis, lateralibus, capitis medio inſertis; *antennis* breuibus, roſtri medio plerumque inſertis; *thorace* ſubcylindrico, antice parum attenuato, ſaepius canaliculato; *ſcutello* nullo; *elytris* rigidis, fornicatis, includentibus, abdomine longioribus; *pedibus* breuibus, validis; *femoribus* compreſſis: *tarſis* quadriarticulatis; *colore* nigro, ſaepius obſcuro.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Brentus myrmecophaga.* Herbſt. Col. tab. 107. fig. 6. mas.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Specimen noſtrum femina. Roſtrum elongatum, cylindricum, antennis verſus apicem inſertis mandibulisque exſertis. Antennae omnino filiformes: articulis elongatis, cylindricis: inferioribus ſex apice clauatis. Thorax oualis, ſcaber, dorſo integro vel absque canali dorſali. Elytra punctato-ſtriata, apice dente valido, recuruo, lineis duabus flauis, in medio coëuntibus. Praeterea lineae quinque glabrae. Pedes omnes inermes.

*aſſimilis.* 3. B. roſtro apice glabro, atro, elytris ferrugineo-ſubfaſciatis. Ent. ſyſt. 2. 491. 2. *

Habitat in noua Zelandia. Muſ. D. Banks.

*monilis.* 4. B. piceus, thorace poſtice canaliculato, elytris acuminatis, vniſtriatis. Ent. ſyſt. 2. 492. 3.

*Brentus monilis.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 61. 2. *

Habitat in Americae Inſulis. Muſ. D. de Seheſtedt.

*futuralis.* 5. B. viridi-aeneus, thorace canaliculato, elytris acuminatis, vniſtriatis: ſutura brunnea.

*Brentus futuralis.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 63. 3. *

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

B. monili paullo minor. Antennae moniliformes, brunneae: articulis tribus vltimis paullo maioribus, griſeis. Roſtrum cylindricum, baſi aeneum, apice ferrugineum. Thorax canaliculatus, aeneus. Elytra laeuia, aenea: ſutura late-brunnea, apice acuminata. Corpus viridi-aeneum.

6.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

6. B. thorace elytrisque vniſulcatis: ſulco villoſo, tibiis anticis ciliatis. *obtuſus.*

*Brentus obtuſus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 64. 4. *

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Magnitudine variat. Antennae omnino moniliformes: articulo vltimo acuto. Roſtrum elongatum, cylindricum, maris ante medium antenniferum, feminae vltra medium mandibulique exſertis. Thorax canaliculatus, canali cinereo, villoſo. Elytra rotundata, punctato-ſtriata, ſulcoque ſuturali cinereo, villoſo. Color vnicolor, nunc niger, nunc piceus. Pedes inermes.

7. B. ater, thorace elytrisque vniſulcatis: ſulco glabro, tibiis anticis glabris. *glabratus.*

*Brentus glabratus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 65. 5. *

Habitat in America meridionali. Muſ. D. de Seheſtedt.

Affinis praecedenti, at roſtrum paullo breuius, craſſius. Foſſula inter oculos impreſſa. Thoracis et elytrorum ſulcus glaber. Tibiae anticae nullo modo ciliatae.

8. B. thorace laeui, nitido, elytris ſubſtriatis, tibiis anticis vnidentatis. *triſtriatus*

*Brentus triſtriatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 66. 6. *

Habitat in Sumatra. D. Daldorff.

Corpus paruum. Roſtrum breue, planum, parum incuruum. Thorax breuiſſimus, obſcure-bruneus,

---

Variant Brenti ſexu. Femina roſtro longiore, cui ante apicem antennae inſertae, mandibulisque exſertis, vti pulcre expoſuit Dom. Lund in Actis ſocietatis hiſtoriae naturalis Hafnienſibus.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

neus, nitidus. Elytra ad ſuturam ſubſtriata. Corpus obſcure ferrugineum. Femora mutica, at tibiae anticae in medio vnidentatae.

*2ſulcatus.* 9. B. thorace roſtroque canaliculatis, elytris ad ſuturam biſtriatis.

*Brentus 2ſulcatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 67. 7. *

Habitat in Sumatra. Dom. Daldorff. Muſ. D. de Seheſtedt.

Statura praecedentis. Roſtrum cylindricum, canali medio impreſſo, qui tamen nec baſin, nec apicem attingit. Thorax canali medio lato impreſſus. Elytra ad ſuturam biſtriata. Color piceus.

*ferrugineus.* 10. B. brunneus, thorace canaliculato, elytris punctato-ſtriatis, ad ſuturam vniſulcatis.

*Brentus ferrugineus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 68. 8. *

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura praecedentium, at paullo minor. Antennae moniliformes, articulis tribus vltimis paullo craſſioribus. Caput inter oculos foſſula exaratum. Thorax valde canaliculatus. Elytra punctato-ſtriata, ad ſuturam vniſulcata. Corpus brunneum, pedibus inermibus.

*brunneus.* 11. B. brunneus, elytris integris immaculatus. Ent. ſyſt. 2. 492. 4. *

*Brentus brunneus.* Oliv. Inſ. 84. tab. 1. fig. 3.

Habitat in Senegallia. Muſ. Dom. Olivier.

*foueatus.* 12. B. ferrugineus, thorace ſubtus vtrinque foueato, elytris faſciis tribus atris.

*Brentus foueatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 69. 9. *

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Par-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Paruus. Antennae moniliformes, ferrugineae: articulis tribus vltimis craſſioribus. Caput atrum, nitidum, roſtro breuiore, ferrugineo, apice nigro. Thorax ferrugineus, nitidus: faſcia media atra, antice dentes duos exſerente. Dorſum integrum, at antice ſub lateribus fouea lata, impreſſa. Elytra truncata, laeuia, ſulco vnico ad futuram, brunnea: faſciis tribus atris.

13. B. ater, thorace ferrugineo. Ent. ſyſt. ſuppl. 174. 5. * *formicarius.*

*Brentus formicarius.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 71. 10. *

*Attelabus formicarius.* Ent. ſyſt. ſuppl. 163. *

Habitat Tranquebariae. Dom. Daldorff. Muſ. D. Lund.

Paruus in hoc genere. Roſtrum cylindricum, atrum, antennis ruſis, moniliformibus: articulo vltimo longiori, cylindrico, clauato. Thorax rufus, antice globoſus. Elytra laeuia, atra, nitida. Pedes ruſi, femoribus clauatis, at inermibus: annulo nigro.

In altero ſexu antennarum clava breuior, ouata.

** *femoribus dentatis.*

14. B. femoribus anticis dentatis, thorace poſtice canaliculato, elytris ſtria ſesquialtera flaua. Ent. ſyſt. 2. 492. 5. *anchorago*

*Brentus anchorago.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 72. 11.

*Curculio anchorago.* Linn. ſyſt. Nat. 2. 613. 56. Muſ. Lud. Vlr. 52.

*Curculio haſtile.* Pall. Icon. 25. 2. fem.

Herbſt. Coleopt. 7. tab. 108. fig. 4. fem.

Voet. Col. 2. tab. 34. fig. 1. 2. fem.

Sulz. Hiſt. Inſ. tab. 4. fig. 6. fem.

β. *Brentus peregrinus.* Herbſt. Col. 7. tab. 108. fig. 1. mas.

Degeer

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Degeer Inſ. 5. 273. 8. tab. 25. fig. 28. mas.
Oliv. Inſ. 84. tab. 1. fig. 2. mas.
Habitat in America meridionali.

*Os* palpis maxillisque. *Palpi* quatuor, inaequales, filiformes: *anteriores* longiores, quadriarticulati: articulo primo breuiſſimo, ſecundo longiori, cylindrico, tertio breui, orbiculato, quarto breui, obtuſo adhaerentes maxillae dorſo; *poſteriores* breuiſſimi, triarticulati: articulo ſecundo paullo craſſiori, tertio breui, cylindrico obtuſo adhaerentes labii apice. *Mandibula* elongata, cornea, inermis, acuta; femina exſerta arcuata. *Maxilla* longitudine palpi compreſſa, piloſa. *Ligula* nulla. *Labium* corneum, quadratum, integrum. *Antennae* in plurimis moniliformes, in paruis filiformes, roſtro inſidentes.

*voluulus.* 15. B. femoribus anticis poſticisque dentatis, thorace pectoreque canaliculo rufo, elytris lineolis flauis. Ent. ſyſt. 2. 493. 6.

*Brentus voluulus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 75. 12. *

*Brentus linearis.* Oliv. Inſ. 84. tab. 1. fig. 4.

*Brentus voluulus.* Herbſt. Col. 7. tab. 108. fig. 2. mas.

*Brentus canaliculatus.* Herbſt. Col. 7. tab. 109. fig. 7. fem.

Habitat in Inſula Guadeloupe.

In figura D. Herbſt femora omnia dentata, at vix merito.

Magnitudine variat.

*cylindrus.* 16. B. femoribus omnibus dentatis, thorace poſtice canaliculato, elytris ſtria interrupta, ferrugineis

*Brentus cylindrus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 77. 13. *

Ha

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino B. voluuli. Antennae extrorſum craſſiores, at vix piloſae. Thorax poſtice canaliculatus, ater. Elytra punctato-ſtriata, atra, ſtria media rufa, bis interrupta. Pectus haud canaliculatum. Femora omnia dentata. Tibiae a medio ciliatae.

7. B. femoribus omnibus tibiisque anticis dentatis, thorace elongato, aeneo, elytris acuminatis. Ent. ſyſt. 2. 493. 7. * *naſutus.*

*Brentus naſutus.* Oliv. Inſ. 84. tab. 2. fig. 7.

Habitat in Jamaica Inſula.

8. B. femoribus dentatis, capite thoraceque canaliculatis, tibiis anticis bidentatis. *canaliculatus.*

*Brentus canaliculatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 78. 14. *

*Brentus armiger.* Herbſt. Col. 7. tab. 108. fig. 5. fem.

*Brentus interruptus.* Herbſt. Col. 7. tab. 108. fig. 6. mas.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Magnitudine variat. Antennae piloſae, extrorſum craſſiores. Caput poſtice roſtrique baſis canaliculata. Thorax pectusque canaliculata. Elytra crenato-ſtriata, nigra, lineolis flauis; maris rotundata, feminae apice valde producta. Pedes atri, femoribus omnibus dentatis tibiisque anticis bidentatis.

9. B. femoribus omnibus dentatis, poſticis bidentatis, elytris rotundatis: ſtria interrupta, flaua. *2dentatus.*

*Brentus 2dentatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 80. 15.

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in America meridionali. D. Smidt. Mus. D. Lund.

Statura praecedentis. Antennae piloſae, extrorſum craſſiores. Roſtrum atrum, haud canaliculatum. Thorax poſtice canaliculatus. Elytra crenato-ſtriata, nigra, ſtria ſaepius interrupta flava, apice in omni ſexu rotundata. Pedes atri, femoribus vnidentatis, poſticis bidentatis.

*reticulatus* 20. B. femoribus dentatis, cinereus, thorace puncto canaliculato, elytris ſulcato-punctatis.

*Brentus reticulatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 81. 16. *

Habitat in Sumatra. D. Daldorff.

Medius. Roſtrum cylindricum, antennis brevibus, craſſis. Caput cinereum. Thorax punctatus, cinereus, canali medio lato exaratus. Elytra ſulcata, ſulcis punctatis; cinerea, ad futuram laeuia, nigra, apice rotundata. Abdomen atrum.

Mas minor, elytris poſtice minus rotundatis.

*depreſſus.* 21. B. femoribus omnibus dentatis, thorace canaliculato, rufus elytrorum futura atra.

*Brentus depreſſus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 83. 17.

Habitat in Guinea. Muſ. Dom. de Seheſtedt.

Statura et magnitudo omnino B. diſparis. Roſtrum cylindricum, atrum, lateribus a baſi ad antennas cinereo-villoſis. Antennae moniliformes. Thorax rufus, depreſſus, canali dorſali cinereo-pubeſcente lateribusque antice parum compreſſis, cinereo-pubeſcentibus. Elytra punctato-ſtriata: punctis impreſſis, cinereo-villoſis, rufa, futura late atra, apice rotundata. Pedes rufi, femoribus dentatis.

*calcar.* 22. B. femoribus omnibus dentatis, thorace integro, tibiis anticis bidentatis.

*Bre*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Brentus calcar.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 84. 18. ♀

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Caput et thorax atra nitida, nullo modo canaliculata. Elytra crenato-ſtriata, nigra, lineolis flauis, apice truncata. Pedes atri, femoribus omnibus dentatis. Tibiae anticae bidentatae.

23. B. femoribus dentatis, thorace integro, elytris apice vnidentatis, pedibus anticis elongatis. *longimanus.*

*Brentus longimanus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 86. 19. ♀

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura omnino praecedentium. Caput obſcure-aeneum, roſtro atro, antennisque filiformmibus. Thorax oblongus, nigro aeneus, laeuis, nullo modo canaliculatus. Elytra ſimpliciter ſtriata, nigro-aenea: lineolis flauis, apice truncata, vnidentata. Pedes nigri, femoribus omnibus dentatis, antici longiores, tibiis vnidentatis.

24. B. femoribus omnibus dentatis, thorace ſubintegro, roſtro vtrinque ſerrato. *ſerriroſtris*

*Brentus ſerriroſtris.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 86. 19. ♂

*Brentus cinnamomeus.* Herbſt. Col. 7. tab. 107. fig. 1. fem.

Habitat in Sumatra. D. Daldorff.

Specimen vnicum mas. Roſtrum apice parum incraſſatum, margine ſerrato. Thorax dorſo plano, vix canaliculato ater, nitidus. Elytra ad ſuturam ſtriata, nigra, flauo-maculata. Pedes ferruginei, femoribus valde incraſſatis, acute-dentatis.

25.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*diſpar.* 25. B. femoribus omnibus dentatis, thorace plano, integro, ferrugineo: vittis duabus nigris, elytris acuminatis, flauo-lineatis. Ent. ſyſt. 2. 493 8. *

*Brentus bifrons.* Ent. ſyſt. 2. 494. 10. *

*Brentus diſpar.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 88. 21. *

*Curculio diſpar.* Linn. ſyſt. Nat. 2. 613. 55. Muſ. Lud Vlr. 50.

α. *Brentus maxilloſus.* Herbſt. Col. 7. tab. 107. fig. 2. fem.

*Brentus Gnatho.* Herbſt. Col. 7. tab. 107. fig 3.

*Brentus diſpar.* Oliv. Inſ. 84. tab. 1. fig. 1. 6.

β. *Brentus ruficollis.* Herbſt. Col. 7. tab. 108. fig. 3.

Habitat in America meridionali. Dom. Drury.

Variat totus ater, lineolis tamen elytrorum eleuatis, glabris, diſtinctis.

*cylindricornis.* 26. B femoribus dentatis, thorace rotundato, nigro, aeneo, elytris ferrugineis, flauo-ſubſtriatis. Ent. ſyſt. 2. 494. 9. *

Habitat in noua Zelandia.

*tridentatus.* 27. B. femoribus dentatis, capite punctato, ſubcanaliculato, elytris ſulcatis, femoribus anticis, compreſſis.

*Brentus tridentatus.* Lund. Act. Hiſt. nat. Hafn. 5. 2. 91. 22. *

Habitat in Sumatrâ. D. Daldorff. Muſ. D. de Seheſtedt.

Caput magnum, cylindricum, punctatum, ſubcanaliculatum, mandibulis exſertis, validis, tridentatis. Thorax lateribus cinereo-villoſis, dorſo glabro, integro, atro. Elytra ſulcata, nigra, flauo-maculata. Corpus atrum. Femora intus bi- extus vnidentata, apice incraſſata, dente

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

dente ordinario, poſtica incraſſata, vnidentata.

164. COLYDIVM. *Palpi* quatuor, clauati: articulo vltimo maiori.
*Ligula* membranacea, emarginata.
*Antennae* perfoliatae.

1. C. ferrugineum, thorace ſulcato. Ent. ſyſt. 495. 1. * — *ſulcatum.*
Habitat in Germania ſub arborum corticibus. *ix.*

2. C. atrum, elytris ſtriatis, apice retuſis. — *retuſum.*
Habitat in Sumatra. Dom. Daldorff.
Sequentibus maius. Caput magnum, cylindricum, canaliculatum, atrum. Thorax cylindricus, vix punctatus, poſtice parum aguſtatus. Elytra ſtriata, poſtice retuſa, margine prominulo. Corpus atrum.

3. C. obſcure-brunneum, elytris teſtaceis. — *bicolor.*
Habitat in Auſtria Dom. de Meyerle ſub cortice Populi.
Corpus maiusculum. Caput obſcure-brunneum, Antennis teſtaceis. Thorax ſtriolis quatuor impreſſis obſcure-ferrugineus. Elytra ſubſtriata, teſtacea. Corpus cylindricum, obſcurum, elytris pedibusque teſtaceis.

4.

---

*Colydii* corpus paruum, cylindricum, elongatum, glabrum, immarginatum, tardum; *capite* ouato, plano, latitudine thoracis, inſerto; *oculis* globoſis, vix prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis; *thorace* elongato, plano: dorſo canaliculato; *ſcutello* triangulari; *elytris* rigidis, fornicatis, planis, longitudine elytrorum; *pedibus* breuibus, validis, compreſſis; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bidentatum* 4. C. nigrum, thorace antice denticulis duobus eleuatis, compreſſis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Maius et latius reliquis. Antennae omnino perfoliatae, nigrae. Thorax marginatus, glaber, niger, lineis duabus dorſalibus, eleuatis, quae antice in denticulos duos eleuatos, compreſſos excurrunt. Elytra laeuia, nigra. Pedes rufi.

*elongatum.* 5. C. atrum, pedibus ferrugineis. Ent. ſyſt. 2. 495. 2. *

*Colydium elongatum.* Payk. Fn. Sv. 3. 312. 1.
*Tritoma ſulcata.* Thunb. Muſ Vpſ. 3. 39.
*Ips linearis.* Oliv. Inſ. 2. 18. 53. tab. 2. fig. 17.
Roſſ. Fn. 50. 123. tab. 2. fig. 45.
Panz. Fn. Germ. 3. tab. 17.
Habitat in Europae ligno antiquo. Dom. de Paykull.

*ſtriatum.* 6. C. nigrum, thorace canaliculato, antice ſubbidentato, elytris crenato-ſtriatis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paruum. Caput planum, margine vtrinque reflexo. Thorax punctatus, niger, dorſo valde canaliculato. Elytra ſtriato-punctata, nigra, ſtriis apice parum eleuatis. Pedes ferruginei.

*filiforme.* 7. C. atrum, elytris ſulcatis: baſi pedibusque ferrugineis. Ent. ſyſt. 2. 496. 3. *

Habitat Halae Saxonum in truncis Quercus. Dom. Hybner.

*rufipes.* 8. C. nigrum, thorace laeui, elytris ſtriatis, pedibus ferrugineis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Sta-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura omnino C. ſtriati, at thorax laeuis, nec rugoſus, nec dentatus. Pedes ferruginei.

9. C. atrum, capite elytrorum baſi pedibusque ferrugineis. *erythrocephalum.*
Habitat in Hungaria. Dom Scheidler.
Statura et magnitudo omnino C. filiforme. Caput obſcure-rufum. Thorax laeuis, ater, immaculatus. Elytra laeuia, atra, baſi ferruginea. Pedes ferruginei.

10. C. ferrugineum, elytris laeuiſſimis. *cylindricum.*
Habitat in America meridionali. Muſ. D. Lund.
Corpus totum elongatum, filiforme, laeue, brunneum.

11. C. rufum, elytris punctato-ſtriatis. Ent. ſyſt. ſuppl. 175. 4 * *rufum.*
Habitat Pariſiis. Dom. Brogniart.

12. C. teſtaceum, thorace crenato: dorſo carinato. Ent. ſyſt. 2. 496. 4. * *frumentarium.*
*Colydium bidentatum.* Payk. Fn. Sv. 3. 313. 2.
*Dermeſtes ſurinamenſis.* Linn. ſyſt. Nat. 2. 562. 29.
*Tenebrio ſurinamenſis.* Degeer Inſ. 5. 54. 5. tab. 13. fig. 12.
Panz. Fn. Germ. 14. tab. 11.
Habitat in Americae meridionalis farina diutius aſſeruata. Muſ. D. Banks.
Meram varietatem Dermeſtae 6 dentati credit D. de Paykull an rite? — *Omnino ſec. Illig. IV. 162*

13. C. atrum, laeue, elytris, antennis pedibusque ferrugineis. *ruficorne.*
Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.
Statura parua, elongata praecedentium. Antennae perfoliatae, rufae. Caput et thorax laeuia, atra, niti-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

nitidula. Elytra ſtriata, nigra. Pedes ferruginei.

165. HYPOPHLOEVS. *Palpi* aequales, clauati.
*Ligula* elongata, integra.
*Antennae* extrorſum craſſiores, vtrinque ſerratae.

*rufipes.* 1. H. ater, nitidus, pedibus ferrugineis.
Habitat in America meridionali. D. Smidt. Muſ. D Lund.
Sequentibus maior. Antennae compreſſae, atrae. Caput et thorax laeuia, atra, nitida. Elytra ſtriata, nigra, immaculata. Corpus nigrum, pedibus ferrugineis.

*Boros.* 2. H. ſupra niger, obſcurus, ſubtus brunneus.
*Boros elongatus.* Herbſt. Col. tab. 110. fig. 7.
Habitat in Finlandia. Muſ. D. Lund.
Reliquis maior. Antennae breues, vtrinque ſerratae, nigrae. Caput punctatum, nigrum. Thorax planus, punctatus, niger, rufo purum micans. Elytra vix ſtriata. Corpus et pedes ferruginea.

*caſtaneus.* 3. H. laeuis, nitidus, caſtaneus, antennis nigris. Ent. ſyſt. 2. 500. 1. *

*Hy-*

---

*Hypophloei* corpus elongatum, cylindricum, glabrum, immarginatum, agile; *capite* ouato, transuerſo, inſerto; *oculis* globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſubter oculos inſertis; *thorace* elongato, planiuſculo; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis, compreſſis; *tarſis* quinquearticulatis; *colore* vario.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Hypophloeus caſtaneus*. Payk. Fn. Sv. 3. 321. 1.
*Ips taxicornis*. Oliv. Inſ. 2. 18. 11. 15. tab. 1. fig. 2.
Roſſ. Fn. Etr. 49. 122. tab. 4. fig. 2.
Panz. Fn. Germ. 12. tab. 13.
Herbſt. Arch. tab. 21. fig. 6.
Habitat ſub cortice ~~Vlmi~~ Quercus et Fagi. Dom. Prof. Helwig.

4. H. laeuis, ater, elytris antennis pedibusque teſtaceis. Ent. ſyſt. 2. 501. 2. * *linearis.*
Payk. Fn. Sv. 3. 325. 6.
Panz. Fn. Germ. 6. tab. 16.
Habitat ſub Pini ſylueſtris cortice Germaniae. Prof. Helwig.

5. H. laeuis, ater, elytris teſtaceis: faſcia atra. Ent. ſyſt. 2. 501. 3. * *faſciatus.*
Payk. Fn. Sv. 3. 324. 5.
Panz. Fn. Germ. 6. tab. 17.
Habitat ſub cortice Quercus Germaniae. Prof. Helwig.

6. H. ferrugineus, elytris ſubſtriatis. Ent. ſyſt. 2. 501. 4. * *depreſſus.*
*Ips vnicolor*. Oliv. Inſ. 2. 18. 12. 16. tab. 2. f. 18.
*Melinus*. Herbſt. Arch. tab. 21. fig. B.
Payk. Fn. Sv. 3. 325. 7.
Panz. Fn. Germ. 1. tab. 23.
Habitat ſub cortice Quercus Germaniae. Prof. Helwig.

7. H. oblongus, rufus, elytris teſtaceis, apice atris. Ent. ſyſt. 2. 501. 5. * *bicolor.*
*Ips bicolor*. Oliv. Inſ. 2. 18. 12. 15. tab. 2. f. 14.
Payk. Fn. Sv. 3. 323. 3.
Panz. Fn. Germ. 12. tab. 14.
Habitat in Gallia, ſub arborum corticibus.

8.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*bicornis.* 8. H. oblongus, rufus, capite bicorni. Ent. ſyſt. 2. 502. 6. *
Habitat in Gallia. Muſ. Dom. Boſc.

166. LYCTVS. *Palpi* breuiſſimi, filiformes.
*Maxilla* breuis, membranacea, bifida.
*Antennae* claua ſolida.

* *antennarum articulo vno craſſiori.*

*politus.* 1. L. niger, antennis pedibusque ferrugineis, thorace plano oblongo, punctato. Ent. ſyſt. 2. 502. 1. *
*Ips picipes.* Oliv. Inſ. 2. 13. 77. tab. 2. fig. 12.
*Tenebrio minutus* Linn. ſyſt. Nat. 2. 1675. 12.?
Panz. Fn. Germ. 4. tab. 18.?
Habitat in Europae Boletis arboreis.

*depreſſus.* 2. L. brunneus, pectore abdomineque nigricantibus, thorace oblongo, plano. Ent. ſyſt. 2. 503. 2. *
*Lyctus depreſſus.* Payk. Fn. Sv. 3. 327. 2.
Habitat Dresdae. D. Heſſe.
Varietatem praecedentis credit Dom. de Paykull, at vix merito.

*nitidulus.* 3. L. glaber, ater, nitidus, capite elytrorum baſi pedibusque ferrugineis. Ent. ſyſt. ſuppl. 175. 2. *
Habitat in Germania. Dom. Meyerhoff.

4.

---

*Lycti* corpus oblongum, depreſſiuſculum, glabrum, immarginatum, tardum; *capite* magno, ouato, plano, inferto; *oculis* paruis, vix prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis, thorace ouato, planiuſculo; *ſcutello* minuto; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis, compreſſis; *tarſis* quinquearticulatis; *colore* obſcuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. L. glaber, ater, elytris puncto ferrugineo. Ent. *2 puſtulatus.*
ſyſt. 2. 503. 3. *
*Lyctus diſpar.* Payk. Fn. Sv. 3. 328. 3. β.
*Ryzophagus clauicornis.* Herbſt. Col. 5. tab. 45. fig. 10. *Ol. bipunctatus tab. 45. f. 9.*
Habitat Halae Saxonum. Dom. Hybner.

5. L. niger, thorace ouato plano vtrinque dentato. *dentatus.*
Ent. ſyſt. 2. 503. 4. *
Habitat in America meridionali. Muſ. D. Lund.

6. L. ferrugineus, elytris laeuibus, teſtaceis. Ent. *brunneus.*
ſyſt. 2. 503. 5. *
Habitat in Americae Inſulis. Dr. Pflug.

7. L. oblongus, ferrugineus, elytris punctato-ſtria- *terebrans.*
tis. Ent. ſyſt. 2 503. 6. *
*Ips terebrans.* Oliv. Inſ. 2. 18. 5. 4. tab. 1. fig. 7.
*Lyctus cylindricus.* Panz. Fn. Germ. 35. t. 18.
Habitat in Germania, ſub arborum corticibus. D. Prof. Helwig.

8. L. obſcurus, elytris ſtriatis, antennis pedibusque *luglandis.*
teſtaceis. Ent. ſyſt. 2. 504. 7.
*Monotoma ſtriata.* Herbſt. Col. 5. tab. 46. f. 1.
Payk. Fn. Sv. 3. 330. 6.
Panz. Fn. Germ. 5. tab. 17.
Habitat Halae Saxonum. Dom. Hybner.

9. L. ater, nitidus, antennis pedibusque piceis. Ent. *biſteroides.*
ſyſt. 2. 504. 8. *
*Ryzophagus biſteroides.* Herbſt. Col. 5. t. 45. f. 2.
Payk. Fn. Sv. 3. 329. 5.
Panz. Fn. Germ. 5. tab. 16.
Habitat in Dania, ſub arborum corticibus. Muſ. D. Lund.

** *Antennarum articulis duobus craſſioribus.*

10. L. niger, thorace rugoſo, elytris ſtriato-crena- *crenatus.*
tis: maculis duabus rufis. Ent. ſyſt. 2. 504. 9. *



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Lyctus crenatus.* Payk. Fn. Sv. 3. 334. 11.
*Ips crenata* Oliv. Inſ. 2. 18. 6. 6. tab. 2. fig. 9.
*Monotoma crenata.* Panz. Fn. Germ. 1. tab. 24.
*Bitoma crenata.* Herbſt. Col. 5. tab. 46. fig. 6.
Herbſt. Arch. 4. tab. 20. fig. 20.
Habitat in Europa, ſub arborum corticibus.

*ruſipennis.* 11. L. niger, thorace rugoſo, elytris ruſis.
Habitat in Germania.
Affinis certe praecedenti, cuius varietatem olim credidi. Differt tamen thorace minus rugoſo elytrisque vix crenatis totis ruſis.

*obſcurus.* 12. L. depreſſus, niger, thorace lineis quatuor eleuatis, elytris obſcuris, eleuato-lineatis.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Minor L. crenato. Antennae ferrugineae: articulis duobus craſſioribus. Caput nigrum, obſcurum, linea vnica eleuata. Thorax depreſſus, niger, obſcurus: lineis quatuor eleuatis. Elytra depreſſa, fuſca: lineis tribus eleuatis.

*canaliculatus.* 13. L. obſcurus, thorace canaliculato, elytris ſtriatis brunneis. Ent. ſyſt. 2. 504. 11.
*Ips oblonga.* Oliv. Inſ. 2. 18. 7. 8. tab. 1. fig. 5.
*Bitoma vnipunctata.* Herbſt. Col. 5. tab. 46. f. 3.
*Dermeſtes linearis.* Thunb. Act. Vpſ. 4. 4. 4.
Payk. Fn. Sv. 3. 332. 9.
Panz. Fn. Germ. 4. tab. 16.
Geoff. Inſ. 1. 103. 9.
Habitat ſub arborum corticibus. Muſ. D. Banks.

*haematodes.* 14. L. obſcurus, thorace punctato-ſubcarinato, elytris crenato-ſtriatis.
Habitat in Carolina. Muſ. D. Boſc.
Statura parua L. canaliculati. Antennae breues, craſſae: articulis vltimis duobus, paullo craſſioribus, ſubferrugineis. Thorax niger, obſcurus.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

rus, depreſſus, ſubcarinatus, vel potius vtrinque parum impreſſus. Elytra ſtriata.

5. L. ater, glaber, nitidus, antennis pedibusque ferrugineis. Ent. ſyſt. 2. 505. 12. * *nitidus.*
*Lyctus nitidus.* Payk. Fn. Sv. 3. 331. 8. I. 55.
Habitat in Germania. Dom. Prof. Helwig.

6. L. oblongus, ferrugineus, elytris punctato-ſtriatis: ſutura nigra. Ent. ſyſt. 2. 505. 13. * *contractus*
*Ips contracta.* Oliv. Inſ. 2. 18. 6. 5. tab. 2. f. 10.
*Bitoma 2punctata.* Herbſt. Col. 5. tab. 46. f. 5.
Geoff. Inſ. 1. 103. 8.
Payk. Fn. Sv. 3. 333. 10.
Habitat in Gallia, ſub arborum cortice.

## 67. CATOPS. *Palpi* inaequales, ſubulati.

*Labium* membranaceum, bifidum: laciniis breuibus, acutis.

*Antennae* extrorſum craſſiores.

C. niger, elytris pedibusque griſeis. *rufeſcens.*
*Tritoma minuta.* Ent. ſyſt. 1. 2. 507. 9. *
*Chryſomela minuta.* Linn. ſyſt. Nat. 2. 593. 50. Fn. Sv. 533.
~~*Ptomaphagus rufeſcens.* Illig. Col. Bor. 1. 87. 1.~~
Habitat ſub corticibus arborum.

2. C.

---

*Catopis* corpus paruum, oblongum, gibbum, imnarginatum, agile,; *capite* latitudine fere thoracis, deflexo; *oculis* paruis, rotundatis, lateralibus poſticis; *antennis* extrorſum craſſioribus: articulo primo longiori, incuruo, ante oculos inſertis; *thorace* latitudine elytrorum, margine rotundato; *elytris* rigidis longitudine abdominis; *pedibus* longiuſculis; *femoribus* compreſſis; *tibiis* extus ſerratis; *tarſis* quinquearticulatis; *colore* nigro aut brunneo, obſcuro.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*fericeus.* 2. C. holoſericeus, nigricans, pedibus teſtaceis.
*Tritoma ſericea.* Ent. ſyſt. 1. 2. 507. 8. *
~~*Catops fericeus.*~~ Payk. Fn. Sv. 1. 342. 1.
*Cholera.* Latreille.
Herbſt. Col. 4. 196. 6.
Habitat in Germania ſub arborum corticibus.
*Os* palpis maxillisque. *Palpi* quatuor inaequales, filiformes: *anteriores* elongati, quadriarticulati; articulo primo breui, rotundato, reliquis ſubaequalibus; vltimo ſubulato, adhaerentes maxillae dorſo; *poſteriores* breues triarticulati: articulo vltimo ſubulato, paullo obtuſiore adnati labii laciniis. *Labrum* breue, transuerſum. *Mandibula* breuis, cornea, arcuata, acuta, inermis. *Maxilla* membranacea, bifida: laciniis tenuibus, obtuſis: exteriore paullo longiore. *Ligula* baſi cornea, apice membranacea, bifida: laciniis breuibus, linearibus, acutis. *Labium* breue, corneum, transuerſum, integrum.

*vittatus.* 3. C. rufus, elytris nigris: vitta nigra.
*Tritoma vittata.* Ent. ſyſt. 2. 506. 6. *
Habitat in India. Muſ. Dr. Hunter.

*morio.* 4. C. holoſericeus, ater, antennis pedibusque concoloribus.
*Tritoma morio.* Ent. ſyſt. 2. 507. 7. *
*Catops morio.* Payk. Fn. Sv. 1. 344. 2.
*Ptomaphagus fornicatus.* Illig. Col. Bor. 1. 89. 3
*Dermeſtes fornicatus.* Degeer Inſ. 4. 216. 9 tab. 8. fig. 15.
Panz. Fn. Germ. 8. tab. 1.
Habitat Kiliae. D. Daldorff.

*flauipes.* 5. C. fuſcus, ore, antennis pedibusque ferrugineis elytrorum ſtriis crenatis.
*Helops flauipes.* Ent. ſyſt. 1. 122. 27. *

Habi

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Americae meridionalis Inſulis. Dom. Smidt.

6. C. niger, elytris pedibusque teſtaceis. *agilis.*
*Tritoma dubia.* Ent. ſyſt. 2. 506. 5. *
~~*Ptomaphagus agilis.* Illig. Col. Bor. 1. 88. 2.~~ VIII. 34.
Habitat Kiliae. Muſ. D. Lund.

## 168. MYCETOPHAGVS.

*Palpi* inaequales.
*Maxilla* membranacea, vnidentata.
*Ligula* rotundata, integra,
*Antennae* extrorſum craſſiores.

1. M. niger, elytris ſtriatis: faſcia apiceque rufis, antennis pedibusque ferrugineis. Ent. ſyſt. ſuppl. 175. 1. * *fasciatus.*
Habitat Tranquebariae. D. Daldorff.

2. M. rufus, thorace elytrisque nigris: his maculis duabus rufis. *4 maculatus.*
*Silphoides Boleti.* Herbſt. Arch. tab. 61. f. 10.
*Chryſomela 4puſtulata.* Linn. ſyſt. Nat. 2. 592. 80. Fn. Sv. 549.
Panz. Fn. Germ. 12. tab. 9.
~~Geoff. Inſ. 1. 106. 16.~~ I. 49.
Payk. Fn. Sv. 3. 315. 1.
Habitat in Germaniae Boletis.

3. M.

---

*Mycetophagi* corpus oblongum, glabrum, immarginatum, agile; *capite* paruo, rotundato, inſerto; *oculis* prominulis, globoſis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis; *thorace* transuerſo, ouato: marginibus rotundatis; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, curſoriis, compreſſis; *tarſis* quinquearticulatis; *colore* plerumque rufo-nigro, vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ellipticus.* 3. M. niger, elytris laeuibus: macula baſeos elliptica ſanguinea.

*Tenebrio ellipticus.* Ent. ſyſt. ſuppl. 49. 15. *

Habitat in Carolina, ſub arborum cortice. Muſ. D. Boſc.

*Ianus.* 4. M. ſupra niger, ſubtus ferrugineus, elytris laeuibus.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Antennae extrorſum craſſiores nigrae, baſi apiceque ferrugineae. Caput, thorax, elytra laeuia, nigra, opaca, immaculata. Corpus et pedes ferruginea.

*bicolor.* 5. M. obſcurus, ſubtus antennis pedibusque ferrugineis, elytris ſtriatis. Ent. ſyſt. 2. 497. 2. *

Habitat in Carolina. Muſ. D. Boſc.

Affinis praecedenti, at diſtinctus videtur paullo minor, et elytra ſtriata.

*2 puſtulatus.* 6. M. niger, opacus, elytris macula ferruginea.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Antennae nigrae, articulo primo vltimoque ferrugineis. Caput et thorax nigra. Elytra laeuia, nigra, macula magna, ferruginea. Corpus piceum, pedibus teſtaceis.

*maculatus* 7. M. niger, elytris, macula baſeos, faſcia poſtica apiceque ferrugineis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura et magnitudo praecedentium. Antennae ferrugineae. Caput et thorax nigra, laeuia, obſcura, immaculata. Elytra laeuia, obſcura, macula baſeos, faſcia verſus apicem apiceque obſcure rufis. Corpus nigrum, pedibus rufis.

8. M.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

8. M. niger, obſcurus, pedibus teſtaceis. *rufipes.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Praecedente paullo minor. Corpus totum nigrum, obſcurum. Antennae baſi parum et pedes ferruginei.

9. M. niger, obſcurus, elytris faſcia media pedibusque rufis. *faſciatus.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura et magnitudo omnino M. 2 puſtulati, cuius forte mera varietas. Differt tantum faſcia elytrorum lata, loco puncti. Corpus nigricans, pedibus ferrugineis.

10. M. niger, thorace elytrisque ferrugineo nigroque variis. *varius.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura praecedentium, at paullo minor. Antennae nigrae, articulo primo vltimoque ferrugineis. Thorax elytraque laeuia, obſcura, ferrugineo valde variegata. Corpus nigrum, abdominis margine rufo. Pedes ferruginei.

11. M. niger, elytris ſtriatis, pedibus rufis. *flauipes.*
Habitat in Carolina. Muſ. D. Boſc.
Statura omnino M. bicoloris, at paullo minor, et antennae et corpus nigra.

12. M. ſupra niger, elytris laeuibus, baſi apiceque rufis. *glabratus.*
Habitat in Germania. D. de Meyerle.
Statura praecedentium, M. faſciato affinis. Caput nigrum. Thorax minus niger, margine in primis parum rufeſcente. Elytra laeuia, nigra, baſi apiceque late rufa. Corpus cum pedibus rufum.

13. M.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*dermeſtoides.* 13. M. fuſcus, abdomine pedibusque teſtaceis. Ent. ſyſt. 2. 498. 3.
Habitat in Germaniae Boletis.

*atomarius* 14. M. niger, elytris, punctis faſciaque poſtica fuluis. Ent. ſyſt. 2. 498. 4. *
*Dermeſtes atomarius.* Thunb. Inſ. Sv. 67. 78.
Payk. Fn. Sv. 3. 317. 3.
I. 19 Panz. Fn. Germ. 12. tab. 10.
Habitat Halae Saxonum. Dom. Hybner.

*nigrita.* 15. M niger, ſubtus piceus, elytris ſubſtriatis.
Habitat Tranquebariae.
Magnitudo fere M. dermeſtoides. Caput, thorax elytraque ſubſtriata, nigra, immaculata. Corpus piceum.

*multipunctatus.* 16. M. rufus, elytris ſubſtriatis: punctis rufis, numeroſis. Ent. ſyſt. 2. 498. 5. *
*Dermeſtes multipunctatus.* Thunb. Inſ. Suec. 679.
I. 50. Payk. Fn. Sv. 3. 320. 7.
Panz. Fn. Germ. 12. tab. 11.
Habitat in Sueciae Boletis. Dom. de Paykull.

*10 punctatus.* 17. M. niger, elytris punctis quinque pedibusque rufis.
Habitat in Ruſſia. Muſ. D. Lund.
Statura et magnitudo M. multipunctati. Antennae nigrae: articulo vltimo rufo. Caput et thorax laeuia, nigra, thoracis baſi punctis duobus impreſſis. Elytra ſubſtriata, nigra, punctis quinque 1. 1. 2. 1. rufis. Corpus nigrum, pedibus rufis.

*picicornis.* 18. M. ater, elytris ſtriatis, antennis pedibusque piceis. Ent. ſyſt. 2. 498. 6. *
Habitat in Americae meridionalis Inſulis. D. Smidt.

*lunaris.* 19. M. rufus, elytris ſubſtriatis, nigris: lunulis duabus punctisque duobus rufis.

Ha-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Germania.
Statura praecedentium. Antennae obſcurae, apice rufae. Caput nigrum, ore rufo. Thorax rotundatus, laeuis, rufus. Elytra ſubſtriata, nigra, lunula baſeos, puncto ad marginem, lunula transuerſa punctoque ante apicem rufis.

20. M. ater, elytris faſciis duabus lunatis punctoque apicis rufis. *finuatus.*
Habitat in Auſtriae Betula. Dom. de Meyerle.
Statura praecedentis, at duplo minor. Caput cum antennis nigrum. Elytra ſublaeuia, nigra, faſciis duabus latis, lunatis punctoque paruo apicis rufis. Pedes rufi.

21. M. niger, thorace rufo, elytris ſtriatis: margine maculisque duabus rufis. Ent. ſyſt. 2. 499. 8. * *fuluicollis.*
Payk. Fn. Sv. 3. 320. 6. VI. 39.
Habitat in Germania. Muſ. D. Lund.

22. M. piceus, elytris ſtriatis, nigris: baſi punctoque poſtico ferrugineis. Ent. ſyſt. 2. 499. 9. * *piceus.*
Payk. Fn. Sv. 3. 318. 4.
Panz. Fn. Germ. 1. tab. 22. I. 50.
Habitat in Germania. Dom. de Hattorff.

23. M. niger, elytris macula baſeos ferruginea. *humeralis.*
Habitat in America meridionali. D. Smidt. Muſ. Dom. de Seheſtedt.
Paruus, ſubuilloſus, ſupra niger, elytris macula magna baſeos ferruginea. Corpus ferrugineum.

24. M. flauus, antennis nigris. Ent. ſyſt. 2. 499. 11. *nigricornis.*
Habitat Halae Saxonum. Dom. Hybner.

25. M. ater, elytris ſtriatis, antennis pedibusque caſtaneis. Ent. ſyſt. 2. 499. 12. * *caſtaneus.*
Payk. Fn. Sv. 3. 316. 2. II. 66.
Habitat in Germania. Dom. Prof. Helwig.

26.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Populi.* 26. M. teſtaceus, capite elytrisque faſciis duabus fuſcis. Ent. ſyſt. ſuppl. 176. 12. *

Payk. Fn. Sv. 3. 319. 5.

Habitat in Sueciae Populo. Dom. de Paykull.

*metallicus.* 27. M. aeneus, pedibus ferrugineus. Ent. ſyſt. 2. 499. 13. *

Habitat Halae Saxonum. Dom. Hybner.

*teſtaceus.* 28. M. teſtaceus, immaculatus. Ent. ſyſt. 2. 499. 14. *

Habitat in Germaniae Boletis. D. Prof. Helwig.

## 169. TRITOMA. *Palpi* anteriores ſecuri-formes.

*Antennae* claua perfoliata.

*4guttatum* 1. T. ferrugineum, elytris macula baſeos apicisque flauis.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Paullo maior T. 2 puſtulata. Antennae ferrugineae: claua compreſſa. Caput et thorax laeuia, glabra, ferruginea, immaculata. Elytra punctato-ſtriata, ferruginea: macula baſeos apiceque late albidis. Corpus ferrugineum.

*faſciatum.* 2. T. pallidum, elytris puncto fuſcia punctoque apicis fuſcis.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Sta-

---

*Tritomatis* corpus ouatum, gibbum, glabrum, immarginatum, agile; *capite* paruo, ouato, inſerto; *oculis* globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis; *thorace* gibbo, transuerſo, ouato; *ſcutello* rotundato; *elytris* rigidis, ſornicatis, gibbis, longitudine abdominis, apice deflexis; *pedibus* breuibus, compreſſis; *tarſis* quinquearticulatis; *colore* nigro rufoque nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura et magnitudo praecedentis. Antennae pallidae, claua compreſſa, perfoliata, fuſca. Thorax marginatus, pallidus, interdum obſcurior. Elytra glabra, laeuia, pallida, puncto vno aut duobus baſeos, faſcia lata media, ad futuram dilata punctoque apicis ferrugineis. Corpus ferrugineum, pedibus pallidis.

3. T. atrum, elytris macula laterali coccinea, pedibus nigris. Ent. ſyſt. 2. 505. 1.* *2 puſtulatum.*

*Tritoma.* Geoff. Inſ. 1. 335. 1. tab. 6. fig. 2.
Panz. Naturf. 24. 12. 17. tab. 1. fig. 17.
Payk. Fn. Sv. 3. 335. 1.
Herbſt. Col. 4. 193. 1. tab. 43. fig. 11.
Habitat in Boletis arboreis Angliae, capta d. 17. Aprilis ad Hamſtead.

4. T. atrum, elytris macula laterali rufa, pedibus rufis. *humerale.*
Habitat in Carolina. Muſ. Dom. Boſc.
Statura omnino praecedentis, at diſtincta videtur. Antennae rufae, claua nigra. Differt inprimis corpore paullo minore, minus nitido, macula elytrorum laterali absque puncto nigro pedibusque totis rufis.

5. T. rufum, capite, thorace pedibusque atris. *ruſipenne.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Magnitudo T. 2 puſtulatae. Caput et thorax atra, nitida, immaculata. Elytra laeuiſſima, ſanguinea, immaculata. Corpus rufum, pedibus nigris.

6. T. pallide nitidum, antennis atris. *nigricorne*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus totum glabrum, pallidum, nitidum, antennis ſolis atris.

7.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*glabrum.* 7. T. glabrum, atrum, antennis pedibusque piceis. Ent. syst. 2. 506. 2. *
Payk. Fn. Sv. 3. 335. 2.
~~#. 66. Panz. Fn. Germ. 23. tab. 19.~~
Habitat in Suecia, sub arborum cortice. D. de Paykull.

*marginatum.* 8. T. atrum, abdominis margine sanguineo. Ent. syst. 2. 506. 3. *
Habitat in Americae meridionalis Insulis. Dom. Smidt.

*connatum.* 9. T. pubescens, nigrum, obscurum, thorace pedibusque rufis, elytris connatis.
*Tritoma connata.* Panz. Fn. Germ. 36. tab. 18.
*III. 64* Habitat in Austria. Dom. de Meyerle.
Statura parua praecedentium, at distinctum elytris connatis.

*collare.* 10. T. nigrum, thorace pedibusque rufis.
Habitat in Austria. D. de Meyerle.
Maior quam T. rufipes. Caput cum antennis nigrum. Thorax laeuis, rufus, nitidus. Elytra laeuia, obscura. Corpus nigrum, ano parum rufescente.

*rufipes.* 11. T. atrum, nitidum, capite, thorace pedibusque rufis. Ent. syst. 2. 506. 4. *
Habitat in Norwagiae, Germaniae fungis arboreis.

*pulchellum* 12. T. rufum, nitidum, elytris atris. Ent. syst. suppl. 177. 4. *
Habitat in India. Mus. D. de Seheſtedt.

*punctatum* 13. T. rufum, nitidum, thorace punctis quatuor atris.
Habitat in America meridionali. D. Smidt. Mus. D. Lund.
Media. Antennae rufae, claua perfoliata, atra. Caput rufum, nitidum, puncto frontali atro. Tho-

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Thorax laeuis, rufus, nitidus: punctis quatuor atris 2. 2. Elytra ſubpunctato-ſtriata, rufa. Corpus rufum, immaculatum.

14. T. rufum, nitidum, antennarum claua nigra. *rufum.*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Media. Caput et thorax laeuia, rufa, nitida, antennarum claua perfoliata, nigra. Elytra ſubtiliſſime ſtriato-punctata. Corpus rufum.

15. T. glabrum, atrum, pedibus ferrugineis. *atratum.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus medium totum atrum, nitidum, pedibus ſolis ferrugineis.

16. T. glabrum, ferrugineum, elytris baſi faſciaque media atris. *2faſciatum*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Media. Caput et thorax obſcure rufa et fere nigra, nitida. Elytra laeuia, rufa, baſi faſciaque pene medium atris. Corpus rufum.

17. T. nigrum, elytris fuſcis: ſtrigis vndatis albis. *vndatum.*
*Anthribus vndatus.* Ent. ſyſt. 2. 378. 7. *
*Bruchus vndatus.* Mant. Inſ. 1. 41. 9.
Habitat in Africae floribus. D. Prof. Vahl.
Species mihi haud rite nota. Vix ad hoc genus pertinere videtur.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

170. TETRATOMA. *Palpi* craſſiuſculi inaequales.
*Maxilla* membranacea, bifida.
*Ligula* breuiſſima rotundata, integra.
*Antennae* articulis quatuor vltimis clauato-perfoliatis.

*Fungorum.* 1. T. rufum, capite elytrisque nigris. Ent. ſyſt. 2. 507. 1. *
*Tetratoma dermeſtoides.* Herbſt. Col. 4. tab. 38. fig. 7.
Panz. Fn. Germ. 9. tab. 10.
Payk. Fn. Sv. 3. 337. 1.
Habitat in Fungis arboreis Germaniae. D. Daldorff.

*dimidiatum.* 2. T. atrum, capite, thorace pedibusque rufis.
Habitat in Carolina. Muſ. D. Boſc.
Media. Antennae huius generis nigrae, baſi rufae. Caput et thorax ſubſtriata, glabra, rufa, immaculata. Elytra laeuia, atra. Corpus atrum, pedibus rufis.

*ruficolle.* 3. T. nigrum, thorace pedibusque rufis. Ent. ſyſt. ſuppl. 176. 2. *
Habitat in Selandia. Muſ. D. Lund.

2.

---

*Tetratomatis* corpus oblongum. glabrum, immarginatum, tardum; *capite* ouato, inſerto; *oculis* magnis, globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis; *thorace* gibbo, transuerſo; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis, apice deflexis; *pedibus* tenuibus, longiuſculis; *tarſis* quinquearticulatis; *colore* atro, nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

4. T. capite thoraceque teſtaceis, coleoptris nigris: macula communi lobata alba. Ent. ſyſt. 2. 508. 2. * *ancora.*

Panz. Fn. Germ. 9. tab. 9.
Herbſt. Col. 4. tab. 38. fig. 8.
Habitat in Germania. Dom. Prof. Helwig.

I. 57.

171. SCAPHIDIVM. *Palpi* filiformes. *Maxilla* bifida. *Antennae* articulis quinque clauato-perfoliatis.

1. S. atrum, elytris truncatis: maculis duabus rufis. Ent. ſyſt. 2. 509. 1. * *4maculatum.*

*Scaphidium 4maculatum.* Oliv. Inſ. 2. 20. 1. tab. 1. fig. 1.
Payk. Fn. Sv. 3. 338. 1.
Panz. Fn. Germ. 12. tab. 11.
Herbſt. Col. 5. tab. 49. fig. 4.
Habitat in Germaniae truncis putreſcentibus.

I. 57

2. S. atrum, nitidum, thoracis margine elytrorumque macula ſanguineis. Ent. ſyſt. 2. 509. 2. * *4puſtulatum.*

*Scaphidium 4puſtulatum.* Oliv. Inſ. 2. 20. 4. 2. tab. 1. fig. 2.
Habitat in noua Hollandia. Muſ. D. Banks.

3.

---

*Scaphidii* corpus ouatum, antice poſticeque acuminatum, glabrum, immarginatum, agile; *capite* paruo, acuto, inſerto; *oculis* globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſub oculis inſertis; *thorace* transuerſo, antice parum anguſtato; *ſcutello* minuto, rotundato; *elytris* rigidis, fornicatis, abdomine acutiſſimo breuioribus; *pedibus* longiuſculis, teretibus, eurſoriis; *tarſis* quinquearticulatis; *colore* atro rufoque nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*immaculatum.* 3. S. atrum, nitidum, elytris immaculatis, punctat ſtriatis. Ent. ſyſt. ſuppl. 177. 3. *
*Scaphidium immaculatum.* Oliv. Inſ. 2. 20. 3. tab. 1. fig. 3.
Habitat in Gallia. Muſ. D. Boſc.

*agaricinum.* 4. S. atrum, nitidum, antennis pedibusque flau Ent. ſyſt. 2. 510. 3. *
*Scaphidium agaricinum.* Oliv. Inſ. 2. 20. 5. tab. 1. fig. 4.
*Silpha agaricina.* Linn. ſyſt. Nat. 2. 670. Fn. Sv. 448.
*Dermeſtes hemipterus.* Panz. Naturf. 24. tab. fig. 14.
Panz. Fn. Germ. 12. tab. 16.
Payk. Fn. Sv. 3. 339. 2.
Herbſt. Col. 5. tab. 49. fig. 5.
Habitat in Europae Agaricis.

*concolor.* 5. S. atrum, nitidum, elytris laeuiſſimis.
Habitat in America, in Carolina. Muſ. D. Boſc.
Omnino diſtinctum a S. immaculato. Corpus di plo minus, totum atrum, pedibus obſcure p ceis. Elytra laeuiſſima.

172. I

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

2. IPS. *Palpi* aequales, breuiſſimi: articulo vltimo ouato.
*Maxilla* bifida.
*Ligula* conica, emarginata.
*Antennae* perfoliatae.

. ferruginea, elytris atris: punctis ſex albis. *guttata.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Corpus oblongum. Antennae perfoliatae, nigrae. Caput et thorax ferruginea, immaculata. Elytra laeuia, atra, punctis ſex albis 2. 2. 2. Corpus ferrugineum.

. glabra, atra, elytris maculis duabus rufis. Ent. *grandis.*
ſyſt. 2. 511. 2. *
Habitat in Africa aequinoctiali. Muſ. D. Lee.

. glabra, atra, elytris faſciis duabus rufis: ante- *abbreuiata*
riore abbreuiata.
Habitat in Sumatra. D. Daldorff.
Statura praecedentium, at diſtincta. Caput et thorax laeuia. Elytra atra, faſciis duabus rufis. Anterior futuram haud attingit, at in medio antice poſticeque dentem emittit. Poſterior vix nec futuram, nec marginem attingit.

. atra, elytris limbo antico faſciaque poſtica ſinua- *ſinuata.*
tis, fuluis.
Habit t in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Me-

---

*is* corpus oblongum, gibbum, glabrum, immarginatum, agile; *capite* ouato, planiuſculo, inſerto; *oculis* globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, ſubter oculos inſertis; *thorace* transuerſo, ſubquadrato; *ſcutello* rotundato; *elytris* rigidis, fornicatis, longitudine abdominis; *pedibus* breuibus, validis; *tibiis* compreſſis; *tarſis* quadriarticulatis; *colore* plerumque atro, nitido.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Media. Antennae claua perfoliata. Caput et thorax laeuia, atra, nitida. Elytra fubpunctato-ſtriata, atra, nitida, baſi margineque vltra medium ſinuatis, fuluis. Pone medium faſcia vtrinque ſinuata, flaua, quae tamen ſuturam haud attingit. Corpus cum pedibus atrum.

*collaris.* 5. I. ferruginea, thorace atro: margine albo, puncto nigro.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura et magnitudo praecedentis. Antennae ferrugineae, claua nigra. Caput atrum, ore ferrugineo. Thorax ater, laeuis, nitidus, lateribus late albis: puncto nigro. Elytra punctato-ſtriata, ferruginea, macula humerali atra. Corpus ferrugineum, pectore atro.

*nigricornis.* 6. I. teſtacea, nitida, antennis nigris.

Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.

Statura praecedentium, at tota glabra, teſtacea, nitida, antennis ſolis nigris.

*2punctata.* 7. I. atra, elytris ruſis: macula magna, atra. Ent. ſyſt. 2. 511. 3. *

Habitat in Europa auſtraliori. Dom. Saldoner.

*2puſtulata.* 8. I. atra, elytris macula baſeos rufa. Ent. ſyſt. 2. 511. 4. *

*Vill. 55.* Habitat Halae Saxonum. Dom. Hybner.

*faſciata.* 9. I. pallida, thorace punctis elytris faſcia dentatu atris.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Statura Ipis 2punctatae, at minor. Caput pallidum, antennarum claua compreſſa, nigra. Thorax pallidus: punctis ſeptem nigris fere in annulum po

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

positis. Scutellum nigrum. Elytra laeuia, pallida, nitida, maculis duobus baseos obscuris, fascia media lata, dentata, atra, quae ad suturam dilatatur, fasciolaque ante apicem obscura. Corpus palliduu.

10. I. pallida, coleoptris macula media nigra. Ent. syst. 2. 512. 5. * *dorsalis.*

Habitat in Europa australi. Dom. Prof. Vahl.

11. I. nigra, elytris laeuissimis: punctis duobus ferrugineis. Ent. syst. 2. 512. 6. * *4pustulata.*

*Silpha quadripustulata.* Linn. syst. Nat. 2. 570. 5. Fn. Sv. 446.
Degeer Inf. 4. 185. 12. tab. 6. fig. 20. 21.
Naturf. 24. 12. tab. 1. fig. 18.
Oliv. Inf. 2. 12. 8. 9. tab. 3. fig. 22.
Payk. Fn. Sv. 3. 341. 1.
Herbst. Col. 4. tab. 42. fig. 1.
Habitat in Europa boreali.

12. I. nigra, elytris striatis: punctis quatuor rufis. Ent. syst. suppl. 179. 6. * *4notata.*

Payk. Fn. Sv. 3. 344. 4.
Habitat in Germaniae truncis.

13. I. picea, elytris subpunctatis, nigris, basi ferrugineis. *punctata.*

*Mycetophagus punctatus* Ent. syst. 2. 499. 10. * I. 51.
*Cryptophagus punctatus.* Payk. Fn. Sv. 3. 351. 1.
*Cryptophagus pilosus.* Herbst. Col. 4. tab. 42. fig. 15.
Panz. Fn. Germ. 12. tab. 12.
Habitat in Germaniae fungis. Dom. Prof. Helwig.

14. I. nigra, elytris fasciis duabus punctoque apicis ferrugineis. *bifasciata.*

*Mycetophagus 2fasciatus* Ent. syst. 2. 499. 11. * I. 51.
*Cryptophagus 2fasciatus.* Payk. Fn. Sv. 3. 352. 2. V. 18



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Herbſt. Col. 4. tab. 42. fig. 12.
Panz. Fn. Germ. 2. tab. 14.
Habitat Halae Saxonum. Dom. Hybner.

*6 puſtulata* 15. I. nigra, elytris ſtriatis: maculis tribus rufis. Ent. ſyſt. 2. 512. 7. *

Habitat in Europa boreali. Muſ. D. de Seheſtedt.

*4 guttata.* 16. I. atra, nitida, elytris maculis duabus albis: anteriore ſinuata. Ent. ſyſt. 2. 513. 8. *

*Nitidula 4guttata.* Oliv. Inſ. 2. 12. 10. 11. tab. 3. fig. 25.
Payk. Fn. Sv. 3. 343. 3.
Herbſt. Col. 4. tab. 42. fig. 3.
I. 59 Panz. Fn. Germ. 3. tab. 18.
Habitat in Europa.

*ferruginea* 17. I. ferruginea, elytris teſtaceis. Ent. ſyſt 2. 513. 9. *

*Dermeſtes ferrugineus.* Linn. ſyſt. Nat. 2. 564. 21. Fn. Sv. 433.
I. 55. Payk. Fn. Sv. 3. 344. 5.
Panz. Fn. Germ. 8. tab. 15.
Habitat in Germania. Muſ. Dom. de Loewenſkiold.

*haemorrhoidalis.* 18. I. rufa, elytris nigris, apice rufis. Ent. ſyſt. 2. 513. 11. *

*Diaperis haemorrhoidalis.* Payk. Fn. Sv. 3. 360. 4.
II. 67. Panz. Fn. Germ. 13. tab. 16.
Habitat in Suecia. Dom. de Paykull.
Differt ab hoc genere, numero articulorum tarſi, vix tamen Diaperis, mihi haud rite examinata.

*pallida.* 19. I. pallida, elytris laeuiſſimis, antennarum clava nigra.

Habitat in America meridionali. D. Smidt. Muſ. D. Lund.

Cor-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Corpus paruum, totum teſtaceum, glabrum, laeue, immaculatum, antennis ſolis, claua fuſca.

173 TRIPLAX. *Palpi* inaequales, ſecuriformes.
*Ligula* trifida.
*Antennae* claua perfoliata.

1. T. rufa, antennis, elytris pectoreque nigris. *nigripennis.*
*Ips nigripennis.* Ent. ſyſt. 2. 513. 10. *
*Silpha ruſſica.* Linn. ſyſt. Nat. 2. 570. 10. Fn. Sv. 449.
*Anthribus ruber.* Degeer Inſ. 5. 28. 30. tab. 8. fig. 12. *V. 19.*
*Triplax ruſſica.* Payk. Fn. Sv. 3. 346. 1.
*Triplax ruſſica.* Herbſt. Col. 6. tab. 49. fig. 12.
*Erotylus ruſſicus.* Oliv. Inſ 89. tab. 1. fig. 1.
Herbſt. Arch. tab. 43. fig. 9.
Habitat in Germaniae arboribus. D. Smidt.
*Os* maxillis palpisque. *Palpi* quatuor inaequales, *anteriores* longiores, quadriarticulati: articulo ſecundo longiſſimo, obconico, tertio breuiore, craſſiuſculo, quarto maximo, compreſſo, ſecuriformi, adhaerentes maxillae dorſo; *poſteriores* breuiores, triarticulati: articulo primo breui, ſecundo longiori, obconico, tertio magno, ſecuriformi, ligulae medio laterali adnati. *Mandibula* cornea, arcuata; acuta, apice fiſſa.

*Ma-*

---

*Triplacis* corpus oblongum, gibbum, glabrum, immarginatum, agile; *capite* paruo, ouato, inſerto; *oculis* magnis, prominulis, globoſis, lateralibus; *antennis* longitudine thoracis ante oculos inſertis; *thorace* ſubquadrato, antice, parum anguſtato, latitudine elytrorum; *ſcutello* paruo, rotundato; *elytris* rigidis, fornicatis, includentibus, longitudine abdominis; *pedibus* longiuſculis, compreſſis, curſoriis; *tarſis* quadriarticulatis; *colore* nigro rufoque nitidulo.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Maxilla* cornea, breuiſſima, bifida: laciniis ſubaequalibus, obtuſis. *Ligula* breuis, membranacea, triſida: laciniis exterioribus arcuatis, acutis, interiore breuiore, obtuſa, ciliata.

*ruſipes.* 2. T. atra, capite thorace pedibusque ferrugineis.
*Ips ruſipes.* Ent. ſyſt. 2. 514. 12. *
*Triplax ruſipes.* Payk. Fn. Sv. 3. 347. 2.
*Il. 67.* Panz. Fn. Germ. 13. tab. 17.
Habitat in Norwagiae Fungis.
Praecedenti valde affinis.

*aenea.* 3. T. coccinea, elytris aeneis, immaculatis.
*Ips aenea.* Ent. ſyſt. 2. 514. 13. *
*Triplax aenea.* Payk. Fn. Sv. 3. 348. 3.
*Silpha aenea.* Act. Hall. 1. 254.
*Cryptophagus aeneus.* Herbſt. Col. 4. tab. 42. fig. 9.
Habitat Halae Saxonum in Fungis arboreis. Dom. Hybner.

## 174. ENGIS. *Palpi* articulo vltimo obtuſo, ſubulato.
*Maxilla* biſida.
*Ligula* rotundata, integra.
*Antennae* claua perfoliata.

*faſciata.* 1. E. atra, elytris faſciis duabus ruſis: anteriore nigro-maculata.

*Ips*

---

*Engidis* corpus oblongum, gibbum, glabrum, immarginatum, agile; *capite* ouato, plano, inſerto; *oculis* magnis, oblongis, transuerſis, ſuperne inferneque obſeruantibus; *antennis* longitudine thoracis ante oculos inſertis; *thorace* quadrato, margine parum incraſſato, ſubreflexo; *ſcutello* latiori, rotundato; *elytris* thorace vix latioribus, rigidis, fornicatis, longitudine abdominis; *pedibus* longiuſculis, curſoriis; *tarſis* quinquearticulatis; *colore* nigro rufoque nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Ips fasciata.* Ent. syst. 2. 511. 1. *
Habitat in America boreali.

*Os* maxillis palpisque. *Palpi* quatuor subaequales, filiformes: *anteriores* vix longiores, quadriarticulati: articulis subaequalibus, quarto paullo longiori, attenuato, obtuso, adhaerentes maxillae dorso; *posteriores* paullo breuiores, triarticulati: articulis subaequalibus, tertio parum longiori, attenuato, obtuso, adnati ligulae medio exteriori. *Mandibula* cornea, arcuata, apice fissa. *Maxilla* membranacea, bifida: laciniis subaequalibus; exteriore parum longiore. *Ligula* breuis, membranacea, rotundata, integra. *Labium* breue, transuersum, corneum, integrum.

2. E. nigra, capite thorace elytrorum puncto baseos pedibusque rufis. *humeralis.*

*Ips humeralis.* Ent. syst. 2. 514. 14. *
*Engis humeralis.* Payk. Fn. Sv. 3. 349. 1.
*Dermestes 2 pustulatus.* Thunb. Act. Upsl. 4. 4. 6.
*Dermestes scanicus.* Panz. Fn. Germ. 4. tab. 9.
*Ips humeralis.* Herbst. Col. 4. tab. 42. fig. 4.
*Dacne.* Latreille Inst.
Habitat in Suecia. Dom. de Paykull.

3. E. nigra, elytris puncto baseos lunulaque postica ferrugineis. *lunata.*

*Ips lunata.* Ent. syst. 2. 514. 15. *
Habitat Kiliae Holsatorum.

4. E. atra, fronte maculis duabus elytrorum pedibusque ferrugineis. *rufifrons.*

*Ips rufifrons.* Ent. syst. 2. 514. 16. *
*Engis rufifrons.* Payk. Fn. Sv. 3. 350. 1. β.
Panz. Fn. Germ. 36. tab. 19.
Habitat in Anglia. Mus. D. Banks.

III. 64

5.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*ſanguini-collis.* 5. E. ater, thorace elytrorumque maculis duabus pedibusque rufis.
*Mycetophagus ſanguinicollis.* Ent. ſyſt. 2. 498. 7. *
*Silpha glabra.* Act. Hall. 1. 255.
*Engis ſanguinicollis.* Payk. Fn. Sv. 3. 351. 2.
*Dermeſtes 4puſtulatus.* Panz. Fn. Germ. 6. tab. 6.
Habitat Halae Saxonum. Dom. Hybner.

175. VPIS. *Palpi* inaequales, clauati.
*Maxilla* vnidentata.
*Ligula* breuis, emarginata.
*Antennae* extrorſum craſſiores.

*ceramboides.* 1. VPIS. Ent. ſyſt. 2. 515. 1. *
*Spondylis ceramboides.* Mant. Inſ. 1. 127. 2.
*Attelabus ceramboides.* Linn. ſyſt. Nat. 2. 621. 12. Fn. Sv. 643.
*Curculio maximus.* Vdm. Diſſ. 26. tab. 1. fig. 1.
*Tenebrio varioloſus.* Degeer Inſ. 5. 32. 2. tab. 2. fig. 1.
Panz. Naturf. 24. 24. 33. tab. 1. fig. 33.
Habitat in Europae borealis Boletis.

176. DIA-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

176. DIAPERIS. *Palpi* subfiliformes.
*Maxilla* bifida.
*Ligula* cylindrica, apice palpigera.
*Antennae* secundum totam longitudinem perfoliatae: foliolis compressis; vltimo ouato.

1. D. nigra, elytris fasciis tribus flauis repandis. *Boleti.*
Ent. syst. 2. 516. 1. *
*Diaperis.* Geoff. Inf. 1. 337. 1. tab. 6. fig. 3.
*Chrysomela Bolcti* Linn. syst. Nat. 2. 591. 36. Fn. Sv. 527.
*Tenebrio Boleti.* Degeer Inf. 5. 49. 9. tab. 3. f. 3.
Schaeff. Elem. tab. 58.
- Icon. tab. 77. fig. 6.
Oliv. Inf. 55. tab. 1. fig. 1.
Sulz. Hist. Inf. tab. 3. fig. 9.
Habitat in arborum Boletis.
*Variat* minor, nuper excluſa, capite thoraceque pallide ferrugineis elytrisque flauescentibus absque fasciis. D. Daldorff.

2. D. atra, elytris rufis: puncto futura fasciaque atris. Ent. syst. suppl. 178. 1. * *Hydni.*
Habitat in Carolinae Hydno candido. Mus. D. Bosc.

3.

---

*Diaperidis* corpus ouatum, gibbum, glabrum, immarginatum, tardum; *capite* paruo, rotundato, deflexo, inserto; *oculis* globosis, prominulis, lateralibus; *antennis* breuibus, distantibus, sub oculis insertis; *thorace* transuerso: margine rotundato, subreflexo; *scutello* minuto, triangulo; *elytris* rigidis, fornicatis, apice deflexis, longitudine abdominis; *pedibus* breuibus, validis; *tarsis* anticis quinque-, posticis quadriarticulatis; *colore* vario, nitido.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*violacea.* 3. D. nigro-cyanea, antennis apice ferrugineis. Ent. ſyſt. 2. 517. 2. *

Habitat in Germaniae Boletis. D. Prof. Helwig.

*viridipennis.* 4. D. thorace ferrugineo, elytris aeneis.

Habitat in Carolina. Muſ. D. Boſc.

Praecedentibus minor. Antennae omnino huius generis, nigrae, baſi rufae. Caput nigrum. Thorax laeuis, ferrugineus, immaculatus. Elytra ſubſtriata, viridi-aenea. Corpus obſcure ferrugineum, pedibus rufis.

*Mas* dentibus duobus clypei; cornubusque duobus erectis, validis nigris in occipite.

*aenea.* 5. D. nigro-aenea, antennarum baſi pedibusque ferrugineis.

Habitat in Boruſſia. Muſ. D. Lund.

Praecedentibus minor. Antennae nigrae, baſi ferrugineae. Caput et thorax laeuia, obſcure aenea, nitida. Elytra ſubſtriata, nigro-aenea. Pedes rufi.

*bicolor.* 6. D. atra, nitida, capite thoraceque obſcure rufis.

*Scaphidium bicolor.* Ent. ſyſt. ſuppl. 174. 4. *

*Diaperis aenea.* Payk. Fn. Sv. 3. 359. 3.

Panz. Fn. Germ. 8. tab. 2.

Habitat in Europae Boletis.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

177. MELOE. *Palpi* subfiliformes.
*Maxilla* recta, bifida.
*Labium* emarginatum.
*Antennae* moniliformes.

1. M. corpore violaceo. Ent. syst. 2. 517. 1. *proscarabaeus.*
~~*Meloe proscarabaeus.* Linn. syst. Nat. 2. 679. 1.~~ *baeus.*
Fn. Sv. 826.
~~Geoff. Inf. 1. 377. 1.~~
~~Frisch. Inf. 6. tab. 6. fig. 5.~~
~~Oliv. Inf. 45. tab. 1. fig. 1.~~
~~Schaeff. Icon. tab. 3. fig. 5.~~
Panz. Fn. Germ. 10. tab. 12. f. 59.
Degeer Inf. 5. 3. 1. tab. 1. fig. 1. ? 60
Sulz. Hist. Inf. tab. 7. fig. 54. 61.
Payk. Fn. Sv. 3. 361. 1. *var.* α.
Habitat in campis apricis arenosis Europae.
*Labrum* rotundatum, emarginatum. *Mandibula* cornea, crassa, acuta, edentula.
Animalculum molle, tardum, oleum limpidissimum flauicans e poris praesertim e pedum geniculis exsudat.

2. M. thorace quadrato nigro, elytris laeuissimis. *laeuigata.*
*Meloe laeuigata.* Oliv. Inf. 45. tab. 2. fig. 5.
Habitat in Marocco. Mus. D. de Seheſtedt.
Maior quam M. proscarabaeus. Caput thorace latius, orbiculatum, nigrum, obscurum. Thorax quadratus, niger, obscurus. Elytra breuia, laeuis-

---

*Meloes* corpus oblongum, molle, glabrum, inmarginatum, tardum; *capite* magno, ouato, exserto; *oculis* oblongis, lateralibus, haud prominulis; *antennis* mediocribus, inter oculos insertis; *thorace* eleuato, quadrato; *scutello* nullo; *elytris* coriaceis, mollibus, abdomine breuioribus; *alis* nullis; *pedibus* elongatis, compressis, cursoriis; *tarsis* antiçis quinque-, posticis quadriarticulatis; *colore* nigro, obscuro.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

uiſſima, nigra. Abdomen feminae longiſſimum, nigrum.

*majalis.* 3. M ſegmentis dorſalibus abdominis rubris. Ent. ſyſt. 2. 518. 2.
~~*Meloe majalis.* Linn. ſyſt. Nat. 2. 679 2.~~
Friſh. Inſ. 6 tab. 6. fig. 4.
Schaeff. Icon. tab. 3. fig. 6.
*I. Co.* Oliv. Inſ 45. tab. 1. fig. 4.
Panz. Fn. Germ. 10. tab. 13.
Habitat in Europa auſtraliori.

*limbata.* 4. M. nigra, thorace plano, elytris limbo ferrugineo.
Habitat in Hungaria. Muſ. D. Lund.
Statura et magnitudo M. proſcarabaei. Caput laeue, nigrum. Thorax quadratus, planus, laeuis. Elytra laeuia, nigra, limbo ferrugineo, qui tamen color marginis exterioris baſin haud attingit. Corpus atrum.

*marginata* 5. M. nigra, thoracis elytrorumque margine ferrugineo. Ent. ſyſt. 2. 518 3. *
*Meloe marginata.* Oliv. Inſ. 45. tab. 1. fig. 3.
*Meloe Hungarus.* Schrank. Beytr. 71. 19.
Habitat in Italiae Saluia verbenacea. Dr. Allioni.

*punctata.* 6. M. atra, thorace elytrisque varioloſo-punctatis. Ent. ſyſt. 2. 518. 4. *
*Meloe vralenſis.* Pall. Iter. 2. 722. 56.
Panz Fn. Germ. 10. tab. 16.
Habitat in Anglia. Muſ. Britt.

*breuicollis.* 7. M. nigra, thorace transuerſo, poſtice retuſo, elytris ſubrugoſis.
*Meloe breuicollis.* Panz. Fn. Germ. 10. tab. 15.
Payk. Fn. Sv. 3. 362. 2.
Habitat in Germania auſtraliore. D. de Meyerle.
Statura et magnitudo omnino M. proſcarabaei, at diſtincta videtur, tota nigra, obſcura. Caput tho-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

thorace latius. Thorax breuis, transuerſus, poſtice retuſus.

8. M. laeuis, cyanea, nitida. *cyanea.*
Habitat in Italia. Dr. Allioni.
Praecedentibus triplo minor. Caput et thorax laeuiſſima, cyanea, nitida. Elytra et abdomen paullo obſcuriora.

## 178. STAPHYLINVS. *Palpi* filiformes.
*Maxilla* vnidentata.
*Ligula* trifida.
*Antennae* moniliformes.

1. St. capite, thorace elytrisque villoſo-aureis. *aureus.*
Ent. ſyſt. 2. 519. 1. *
Oliv. Inſ. 42. tab. 1. fig. 2.
Habitat in Siam. Muſ. D. Banks.

2. St. hirſutus, niger, thorace abdomineque poſtice flauis. Ent. ſyſt. 2. 519. 2. * *hirtus.*
Linn. ſyſt. Nat. 2. 683. 1. Fn. Sv. 839.
Panz. Fn. Germ. 4. tab. 19.
Geoff. Inſ. 1. 362. 7.
Payk. Monogr. 8. 1.
Fn. Sv. 3. 363. 1.
Schaeff. Monogr. 1754. tab. 2. fig. 12.
- Icon. tab. 36. fig. 6.
Degeer Inſ. 4. 20. 5.
Oliv. Inſ. 42. tab. 1. fig. 6.
Sulz.

---

*Staphylini* corpus elongatum, glabrum, immarginatum, agile; *capite* magno, ouato, exſerto; *oculis* globoſis, lateralibus, vix prominulis; *antennis* breuibus, approximatis, in articulationem labii ſuperioris inſertis; *thorace* plano, poſtice rotundato, marginibus deflexis; *ſcutello* maiuſculo, triangulari; *elytris* dimidiatis, rigidis, margini deflexo, alas tegentibus; *pedibus* validis; *tarſis* quinquearticulatis; *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Sulz. Inſ. tab. 7. fig. 16.
Habitat in Europae locis arenoſis.

*nebuloſus.* 3. St. pubeſcens, cinereus, nigro-nebuloſus, abdomine nigro, tibiis teſtaceis. Ent. ſyſt. 2. 520. 3. *
Payk. Fn. Sv. 3. 375. 11.
Habitat in Germania. D. Smidt.

*murinus.* 4. St. pubeſcens, cinereus, nigro-nebuloſus, abdomine atro, pedibus nigris. Ent. ſyſt. 2. 520. 4. *
Linn. ſyſt. Nat. 2. 683. 2. Fn. Sv. 840.
Schaeff. Icon. tab. 4. fig. 11.
Degeer Inſ. 4. 12. 1. tab. 1. fig. 1.
Oliv. Inſ. 42. tab. 6. fig. 51.
Roſſ. Fn. Etr. 1. 247. 608.
Panz. Fn. Germ. 66. tab. 16.
Payk. Monogr. 14. 8.
Fn. Sv. 3. 376. 12.
Habitat in cadaueribus, fimo.
*Larua* hexapoda, nuda, pallida, capite abdominisque ſegmentis tribus anterioribus brunneis, ſubtus tuberculo cylindrico, pediformi. Stroem. Act. Hafn. 1.

*chloropterus.* 5. St. capite thoraceque cupreis, elytris viridibus.
*Staphylinus chloropterus.* Panz. Fn. Germ. 36. tab 20.
Habitat in Auſtria. D. de Meyerle.
Caput et thorax ſubpiloſa, cuprea, obſcure nitida, mandibulis atris, antennisque fuſco-teſtaceis. Elytra villoſa, viridia. Abdomen nigrum, ano ſubferrugineo. Pedes teſtacei, femoribus faſcia aut macula nigra.

*pubeſcens.* 6. St. pubeſcens cinereus, nebuloſus, abdomine ſubtus villoſo-argenteo. Ent. ſyſt. 2. 520. 5. *
*Staphylinus pubeſcens.* Payk. Monogr. 16. 9.

Geoff.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 303. 8.
Degeer Inf. 4. 17. 2.
Oliv. Inf. 42. tab. 2. fig. 15.
Payk. Fn. Sv. 3. 376. 13.
Habitat in Suecia. D. Com. de Souza.

7. St. obſcurus, elytris brunneis fuſco-maculatis. *viduatus.*
Habitat in Carolina. Muſ. D. Boſc.
Statura et magnitudo St. olentis. Antennae nigrae, articulo primo ferrugineo. Caput ferrugineum, mandibulis atris. Thorax punctatus, obſcurus. Elytra brunnea, fuſco obſolete maculata. Abdomen obſcurum, punctis dorſalibus duobus in ſingulo ſegmento obſcurioribus. Pectus pilis aureis tectum. Pedes rufi, femoribus nigris.

8. St. niger, opacus, immaculatus, capite thorace latiore. Ent. ſyſt. 2. 520. 6. * *olens.*
*Staphylinus olens.* Payk. Monogr. 9. 4.
Fn. Sv. 3. 371. 5.
Oliv. Inf. 42. tab. 1. fig. 1.
Panz. Fn. Germ. 27. tab. 1.
Geoff. Inf. 1. 360. 1. tab. 7. fig. 1.
Myll. Zool. Dan. 97. 1090.
Degeer Inf. 4. 22. 3.
Habitat in Europae cadaueribus.

9. St. ater, nitidus, elytris obſcuris, capite thorace latiore. Ent. ſyſt. 2. 521. 7. * *ſimilis.*
*Staphylinus ſimilis.* Payk. Monogr. 105.
Fn. Sv. 3. 371. 6.
Oliv. Inf. 42. tab. 5. fig. 42.
Geoff. Inf. 1. 361. 3.
Schaeff. Icon. tab. 30. fig. 12.
Habitat in Germania. D. Smidt.

10. St. ater, nitidus, elytris piceis. Ent. ſyſt. 2. 521. 8. * *picipennis.*

*Sta-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Staphylinus picipennis* Payk. Fn. Sv. 3. 373. 8.
Habitat in Germania. Muſ. D. de Seheſtedt.

*maxilloſus* 11. St. pubeſcens, niger: fafciis cinereis. Ent. ſyſt. 2. 521. 9.
*Staphylinus maxilloſus*. Linn. ſyſt. Nat. 2. 683. 3. Fn. Sv. 841.
Oliv. Inſ. 42. tab. 1. fig. 5.
*Ill. 65* Panz. Fn. Germ. 27. tab. 2.
Geoff. Inſ. 1. 360. 1. tab. 7. fig. 1.
Schaeff. Icon. tab. 20. fig. 1.
Degeer Inſ. 4. 18. 4. tab. 1. fig. 7. 8.
Payk. Monogr. 5. 2.
Fn. Sv. 3. 369. 2.
Habitat in Europae ſyluis, victitans rapina.
Adultior omnino niger et glaber euadit.

*oculatus.* 12. St. ater, nitidus, capite vtrinque macula fulua. Ent. ſyſt. 2. 521. 10. *
Oliv. Inſ. 42. tab. 2. fig. 19.
Habitat in noua Hollandia. Muſ. D. Banks.

*cyaneus.* 13. St. ater, opacus, thorace elytrisque coeruleſcentibus. Ent. ſyſt. 2. 521. 11. *
Geoff. Inſ. 1. 361. 2.
Oliv. Inſ. 42. tab. 1. fig. 4.
*Ill. 65* Panz. Fn. Germ. 27. tab. 3.
Payk. Monogr. 13. 7.
Fn. Sv. 3. 370. 4.
Habitat in Italia. Dr. Allioni.

*dilatatus.* 14. St. thorace marginato, atro, nitidulo, elytris fuſcis, antennis ſerratis. Ent. ſyſt. 2. 522. 12. *
Payk. Fn. Sv. 3. 389. 29.
Habitat Halae Saxonum. D. Prof. Leske.

*fulgens.* 15. St. cyaneus, nitidus, antennis apice nigris. Ent. ſyſt. 2. 522. 13. *
Habitat in Cayenna. Muſ. D. Boſc.

16.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

16. St. ater, elytris, antennarum basi pedibusque rufis. Ent. syst. 2. 522. 14. * *erythropterus.*
Linn. syst. Nat. 2. 683. 4. Fn. Sv. 842.
Oliv. Inf. 42. tab. 2. fig. 14.
Panz. Fn. Germ. 27. tab. 4.
Geoff. Inf. 1. 364. 9.
Frisch. Inf. 5. 49. tab. 29.
Degeer Inf. 4. 21. 6.
Schaeff. Icon. tab. 2. fig. 2.
Payk. Monogr. 7. 3.
Fn. Sv. 3. 369. 3.
Habitat frequens in fimo.
*Mandibula* cornea, medio dilatata, acutissima.
*Maxilla* cornea, vnidentata, in medio ciliata.
*Ligula* trifida: lacinia intermedia maiore, emarginata.

17. St. niger, aeneo-nitidus, elytris pedibusque testaceis, antennis fuscis. Ent. syst. 2. 522. 15.* *chalcocephalus.*
Oliv. Inf. 42. tab. 2. fig. 16.
Rossi. Fn. Etr. 1. 248. 612. tab. 7. fig. 13.
Panz. Fn. Germ. 27. tab. 5.
Degeer Inf. 4. 22. 8.
Payk. Monogr. 11. 6.
Fn. Sv. 3. 374. 10.
Habitat in Germania. D. Smidt.

18. St. niger, capite thoraceque obscure piceis, elytris aureis, tarsis anticis dilatatis. Ent. syst. 2. 523. 16. * *fossor.*
Panz. Fn. Germ. 27. tab. 6.
Habitat in Germania. Mus. D. de Seheſtedt.

19. St. ater, capite rufo. Ent. syst. 2. 523. 17. * *erythrocephalus.*
Oliv. Inf. 42. tab. 2. fig. 9.
Habitat in noua Hollandia. Mus. D. Banks.



Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*pilosus.* 20. St. pilosus, niger, abdomine fusco, elytris scabris. Ent. syst. 2. 523. 18. *
Oliv. Inf. 42. tab. 1. fig. 3.
Habitat in Cayenna. Dom. v. Rohr.

*splendens.* 21. St. capite thoraceque laeuissimis, atris, elytris aeneis, nitidulis, capite thorace latiore. Ent. syst. 2. 523. 19. *
Habitat in Germania. D. Smidt.

*politus.* 22. St. niger, thorace elytrisque nitidis: thorace lineis punctorum impressorum. Ent. syst. 2. 524. 20. *
Linn. syst. Nat. 2. 683. 5. Fn. Sv. 843.
Geoff. Inf. 1. 361. 3.
Degeer Inf. 4. 22. 7.
Schaeff. Icon. tab. 39. fig. 12.
Oliv. Inf. 42. tab. 2. fig. 10.
Panz. Fn. Germ. 27. tab. 7.
Payk. Monogr. 42. 31.
Fn. Sv. 3 391. 31.
Habitat in Europae cadaueribus.

*tristis.* 23. St. niger, thorace obscure-aeneo, elytris pubescentibus obscurioribus. Ent. syst. 2. 524. 21. *
Habitat in Dania. Mus. D. de Seheſtedt.

*femoratus.* 24. St. fuscus, obscurus, elytris subglaucis, ano rufescente, femoribus flauis.
Habitat in Carolina. Mus. Dom. Bosc.
Statura et magnitudo praecedentium. Antennae obscurae, articulo primo flauescente. Caput et thorax subpunctata, fusca, minime nitida. Elytra obscura, glauca. Abdomen fuscum, ano rufescente. Corpus obscurum, femoribus flauescentibus.

*varians.* 25. St. ater, nitidus, punctatus, alis albis. Ent. syst. 2. 524. 22. *

St...

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Staphylinus varians.* Payk. Monogr. 45. 33.
Fn. Sv. 3. 393. 33.
Oliv. Inf. 42. tab. 5. fig. 46.
Habitat in Europa boreali. Muſ. D. de Seheſtedt.

26. St. niger, pedibus, antennarum baſi apiceque ferrugineis. Ent. ſyſt. 2. 524. 23. * *brunnipes.*
Oliv. Inf. 42. tab. 1. fig. 7.
Habitat in Anglia. Muſ. D. Banks.

27. St. teſtaceus, abdomine atro: ano teſtaceo. *teſtaceus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et magnitudo St. brunnipedis. Caput planum, teſtaceum. Thorax planus, punctatus, teſtaceus. Elytra piloſa, teſtacea, ſubpunctata, immaculata. Abdomen piloſum, atrum, ano teſtaceo, appendiculis nigris. Pedes teſtacei.

28. St. ater, nitidus, antennarum articulo primo elytris abdominisque ſegmento penultimo rufis. *vlmineus.*
*Staphylinus Vlmi.* Roſſ. Fn. Etr. 1. 248. 611. tab. 5. fig. 6. *VIII. 35.*
Oliv. Inf. 42. tab. 4. fig. 37.
Habitat in Italia. Muſ. Dom. Boſc.
Statura St. nitidi, at maior et diſtinctus.

29. St. niger, nitidus, thorace, elytris pedibusque obſcure-brunneis. *obſcurus.*
Habitat in India orientali. D. Daldorff. Muſ. D. de Seheſtedt.
Statura omnino St. nitidi. Antennae nigrae, baſi apiceque brunneae: articulis vtrinque prominulis. Caput atrum, ore brunneo. Thorax et elytra ſubpunctata, fuſco-brunnea, immaculata. Abdomen atrum, ano brunneo. Pedes brunnei.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*nitidus.* 30. St. ater, nitidus, elytris teſtaceis. Ent. ſyſt. 2. 524. 24. *

Panz. Fn. Germ. 22. tab. 8.
Payk. Monogr. 40. 30.
Fn. Sv. 3. 390. 30.
Habitat Kiliae. Dom. Daldorff.

*brunnipennis.* 31. St. niger, obſcurus, elytris pedibusque brunneis.
Habitat in India orientali. D. Daldorff. Muſ. D. Lund.
Medius. Antennae nigrae, baſi ferrugineae. Caput et thorax ſcabra, nigra, obſcura. Elytra ſubpunctata, brunnea, immaculata. Abdomen atrum, ano ſiue vltimo ſegmento brunneo. Pedes brunnei.

*fulgidus.* 32. St. ater, nitidus, elytris plantisque teſtaceis, ano ſubferrugineo. Ent. ſyſt. 2. 525. 25. *
Oliv. Inſ. 42. tab. 4. fig. 34.
Habitat in Germania. D. Smidt.

*haemorrhoidalis.* 33. St. piloſus, ater, nitidus, ano brunneo.
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Statura praecedentis. Caput, thorax et elytra piloſa, atra, nitida, immaculata. Abdomen atrum, ano brunneo. Pedes atri.

*crenatus.* 34. St. nigricans, thorace marginato, elytris crenato-ſtriatis. Ent. ſyſt. 2. 525. 26. *
Payk. Monogr. App. Curc. 146. 45.
Fn. Sv. 3. 403. 47.
Habitat in Germania. D. Smidt.

*ſtriatulus.* 35. St. niger, thorace canaliculato, elytris ſtriatis. Ent. ſyſt. 2. 525. 27. *
Habitat in Germania. D. Smidt.

36.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

36. St. niger, thorace marginato, elytris teſtaceis. *ſtrumoſus.*
Ent. ſyſt. 2. 525. 28. *
Payk. Fn. Sv. 3. 402. 45.
Habitat Halae Saxonum. Dom. Hybner.

37. St. ater, nitidus, elytris cyaneis. Ent. ſyſt. 2. 525. 29. * *cyanipennis.*
*Staphylinus amoenus.* Oliv. Inſ. 42. t. 4. f. 36.
Habitat Pariſiis. Muſ. Dom. Boſc.
Varietatem St. politi credit Paykull, at vix rite.

38. St. ater, thoracis lateribus pedibusque flauis. *marginatus.*
Ent. ſyſt. 2. 526. 30. *
Oliv. Inſ. 42. tab. 3. fig. 29.
Act. Nidroſ. 4. 3321. 33.
Payk. Monogr. 44 32.
Fn. Sv. 3. 392. 32.
Habitat in Europa boreali.

39. St. ater, antennarum baſi, elytris pedibusque teſtaceis. *rufipennis.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Triplo maior St. alpino et alius. Antennae extrorſum craſſiores, nigrae, baſi ferrugineae. Caput et thorax punctata, atra, immaculata, nitida, ore ferrugineo. Elytra teſtacea. Corpus atrum, pedibus teſtaceis.

40. St. ater, nitidus, elytris pedibusque teſtaceis, thorace ouato. Ent. ſyſt. 2. 526. 31. * *fuluipes.*
Panz. Fn. Germ. 27. tab. 9. III. 67.
Habitat in Germania. D. Smidt.

41. St. ater, pedibus rufis, elytrorum margine flauo. Ent. ſyſt. 2. 526. 32. * *flaueſcens.*
Linn. ſyſt. Nat. 2. 684. 9. Fn. Sv. 830.
Habitat in Europa boreali.

42. St. ater, elytris fuſcis: linea media pedibusque teſtaceis. Ent. ſyſt. ſuppl. 180. 32. * *plagiatus.*

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Staphylinus plagiatus*. Payk. Fn. Sv. 3. 387. 26.
Habitat in Europa boreali.

*alpinus*. 43. St. depreſſus, ater, antennarum baſi, elytris pedibusque liuidis. Ent. ſyſt. 2. 526. 33. *
Oliv. Inſ. 42. tab. 6. fig. 55.
Payk. Fn. Sv. 3. 388. 27.
Habitat in Betula, Salice, in ſummis Alpibus Lapponiae. Dom. de Paykull.

*2puſtulatus* 44. St. niger, elytris puncto poſtico ferrugineo. Ent. ſyſt. 2. 526. 34. *
Linn. ſyſt. Nat. 2. 685. 16. Fn. Sv. 847.
III. 67. Panz. Fn. Germ. 27. tab. 10.
Habitat in Europa.

*analis*. 45. St. ater, nitidus, antennis, elytris, ano pedibusque teſtaceis. Ent. ſyſt. 2. 526. 35. *
Oliv. Inſ. 42. tab. 3. fig. 24.
Payk. Monogr. 47. 34.
Fn. Sv. 3. 394. 34.
Habitat Kiliae. Dom. Daldorff.

*merdarius* 46. St. ater, thorace elytris ano pedibusque teſtaceis.
*Oxyporus merdarius*. Ent. ſyſt. 2. 533. 11. *
III. 71. Panz. Fn. Germ. 27. tab. 18.
Oliv. Inſ. 42. tab. 5. fig. 45.
Payk. Monogr. 47. 34. β.
Fn. Sv. 3. 394. 34. β.
Habitat in Europae ſtercore.
Varietatem praecedentis credit D. de Paykull forte rite.

*fuſcipes*. 47. St. ater, thorace ſubrotundo, elytris pedibusque piceis. Ent. ſyſt. 2. 527. 38.
Linn. ſyſt. Nat. 2. 685. 23. Fn. Sv. 857.
III. 68. Panz. Fn. Germ. 27. tab. 12.
Payk. Fn. Sv. 3. 397. 39.
Habitat in Europae fimetis.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

48. St. niger, elytris, antennis pedibusque ferrugi- *flauipes.*
neis. Ent. syst. 2. 527. 39. *
*Staphylinus flauipes.* Linn. syst. Nat. 2. 685.
24. Fn. Sv. 856.
Habitat in Selandia. Mus. D. de Sehestedt.

49. St. thorace rufo, elytris fuscis: puncto baseos *atricapil-*
margineque postico albis. Ent. syst. 2. 527. 40.* *lus.*
Oliv. Ins. 42. tab. 4. fig. 39.
Payk. Monogr. 49. 35. II. 68
Fn. Sv. 3. 414. 60.
Habitat in Anglia. Mus. D. Banks.

50. St. filiformis, ater, elytris, apice pedibusque te- *angustatus*
staceis. Ent. syst. 2. 528. 41. *
*Paederus angustatus* Payk. Fn. Sv. 3. 431. 6.
Oliv. Ins. 42. tab. 2. fig. 18.
Panz. Fn. Germ. 34. tab. 18. I. 62.
Payk. Monogr. 36. 27.
Habitat in Suecia. Dom. de Paykull.

51. St. elongatus, ater, elytris, antennis pedibusque *linearis.*
fuscis. Ent. syst. suppl. 180. 41. *
*Staphylinus linearis.* Oliv. Ins. 42. tab. 4. f. 38.
Habitat in Gallia. Mus. Dom. Bosc.

52. St. flauus, capite abdominisque cingulo atris, *canalicula-*
thorace canaliculato. Ent. syst. 2. 528. 42. * *tus.*
Oliv. Ins. 42. tab. 3. fig. 31.
Panz. Fn. Germ. 27. tab. 13. III. 68
Payk. Monogr. 31. 23.
Fn. Sv. 3. 385. 22.
Habitat Kiliae. D. Daldorff.

53. St. ater, nitidus, thorace elytrorumque apicibus *tenuis.*
rufis. Ent. syst. 2. 528. 43. * III. 73
Habitat in Germania. Dom. de Paykull.

54. St. filiformis, flauus, ano fusco. Ent. syst. 2. *gracilis.*
528. 44. *

*Sta-*

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:26 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Staphylinus gracilis.* Payk. Monogr. 38. 28.
*Paederus anguſtatus.* Fn. Sv. 431. 7. β.
Habitat in Suecia. Dom. de Paykull.

*brunneus.* 55. St. brunneus, capite abdominisque medio atris. Ent. ſyſt. ſuppl. 180. 44. *
*Staphylinus brunneus.* Payk. Monogr. 63. 45. Fn. Sv. 3. 404. 47.
Habitat in Europa boreali.

*punctulatus.* 56. St. filiformis, ater, nitidus, capite ouato. Ent. ſyſt. 2. 528. 45. *
*Staphylinus punctulatus.* Payk. Monogr. 30. 22. Fn. Sv. 3. 380. 17.
Habitat in Suecia.

*emarginatus.* 57. St. depreſſus, brunneus, elytris pallidioribus, thoracis margine reflexo. Ent. ſyſt. 2. 528. 46. *
*Staphylinus emarginatus.* Payk. Monogr. 56. 40.
Payk. Fn. Sv. 3. 401. 44.
Oliv. Inſ. 42. tab. 2. fig. 12.
Habitat in Gallia. Muſ. Dom. Boſc.

*philanthus* 58. St. ater, nitidus, elytris pallidis: margine nigro. Ent. ſyſt. ſuppl. 181. 46. *
Habitat ad Cap. Bon. Spei. Dom. Daldorff.

*limbatus.* 59. St. niger, elytrorum medio pedibusque teſtaceis.
Habitat in Germania. Dom. Karſtens.
Paruus, ſtatura fere St. fuſcipedis, at margo elytrorum exterior et interior niger.

*bicolor.* 60. St niger, elytris pedibusque fuſcis. Ent. ſyſt. 2. 529. 47. *
*Staphylinus bicolor.* Payk Monogr. 21. 13. Fn. Sv. 3. 381. 18.
Habitat in Galliae fimetis. Muſ. Dom. Boſc.

*ruſipes.* 61. St. ater, nitidus, elytris obſcuris, pedibus ruſis. Ent. ſyſt. 2. 529. 48. *
Oliv. Inſ. 42. tab. 4. fig. 35.

Payk.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Payk. Fn. Sv. 418. 65.
Habitat in Germania. D. Smidt.

62. St. niger, elytris teſtaceis ad ſuturam nigris, capite tuberculato. Ent. ſyſt. ſuppl. 181. 49. * *tuberculatus.*
Habitat in India. Dom. Daldorff.

63. St. ferrugineus, capite atro. Ent. ſyſt. 2. 529. 49.* *melanocephalus.*
Oliv. Inſ. 42. tab. 4. fig. 32.
Payk. Monogr. 59. 42.
Fn. Sv. 3. 416. 43.
Habitat in Suecia. Dom. de Paykull.

64. St. niger, elytris liuidis, pedibus ferrugineis, antennis extrorſum craſſioribus. Ent. ſyſt. 2. 529. 50. * *craſſicornis.*
Habitat in Dania. Muſ. D. de Seheſtedt.

65. St. flauus, immaculatus. Ent. ſyſt. 2. 530. 53.* *caraboides*
*Staphylinus caraboides.* Linn. ſyſt. Nat. 2. 530. 53. Fn. Sv. 854.
*Staphylinus fuluus.* Degeer Inſ. 4. 25. 12.
Payk. Monogr. 19. 11.
Fn. Sv. 3. 385. 25.
Oliv. Inſ. 42. tab. 2. fig. 17.
Habitat in Europa.

66. St. niger, thorace elytrisque rugoſis. Ent. ſyſt. 2. 530. 54. * *rugoſus.*
Oliv. Inſ. 42. tab. 5. fig. 43.
Habitat in Anglia.

67. St. niger, thorace depreſſo: lineis eleuatis tribus, elytris piceis. Ent. ſyſt. 2. 530. 55. * *piceus.*
*Staphylinus piceus.* Linn. ſyſt. Nat. 2. 686. 85.
Geoff. Inſ. 1. 367. 16.
Oliv. Inſ. 42. tab. 3. fig. 30.
Panz. Fn. Germ. 27. tab. 14.
Payk. Monogr. 19. 12.
Fn. Sv. 3. 374. 22.
Habitat in Europae borealis fimo.

III. 69



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*porcatus.* 68. St. niger, obſcurus, thorace marginato, elytris eleuato-ſtriatis. Ent. ſyſt. 2. 530. 56. *

Oliv. Inſ. 42. tab. 4. fig. 33.

Payk. Monogr. 71. 50.

Fn. Sv. 3. 413. 59.

Habitat in Sueciae ſtercore bouino, varius. D. de Paykull.

Minutus, at diſtinctus, thoracis margine vtrinque prominulo elytrisque ſtriis tribus eleuatis.

## 179. STENVS. *Palpi* inaequales: *anteriores* longiſſimi. *Antennae* claua ſolida.

*2guttatus.* 1. St. niger, elytris puncto albido, oculis prominulis.

*Staphylinus 2guttatus.* Ent. ſyſt. 2. 527. 36. *

*Staphylinus Iuno.* Payk. Monogr. 34. 35. β.

*Staphylinus 2guttatus.* Linn. ſyſt. Nat. 2. 685. 15. Fn. Sv. 851.

*Paederus 2guttatus.* Oliv. Inſ. 44. tab. 1. fig. 3.

*I. 61.* *Stenus Iuno.* Payk. Fn. Sv. 3. 433. 1. β.

Geoff. Inſ. 1. 371. 24.

Habitat ad littora maris Baltici.

*Varietas* ſequentis teſte D. de Paykull.

*Iuno.* 2. St. niger, immaculatus, thorace rotundato laeui.

*Staphylinus clauicornis.* Ent. ſyſt. 2. 527. 37. *

*III. 68.* *Staphylinus Iuno.* Payk. Monogr. 34. 35. α.

*Stenus Iuno.* Payk. Fn. Sv. 3. 432. 1. α.

Geoff.

---

*Steni* corpus paruum, elongatum, glabrum, immarginatum, agile; *capite* ſubrotundo plano, exſerto, thorace latiore; *oculis* magnis, globoſis, prominulis, lateralibus; *antennis* breuibus, approximatis, frontalibus; *thorace* anguſto, rotundato, cylindrico; *ſcutello* vix vllo; *elytris* valde abbreuiatis, truncatis, rigidis, alas tegentibus; *abdomine* elongato, cylindrico, margine vtrinque eleuato; *pedibus* longiuſculis, tenuibus, curſoriis; *tarſis* quadriarticulatis; *colore* nigro, obſcuro.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY

Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 371. 24. *a.*
Panz. Fn. Germ. 27. tab. 11.
Habitat Kiliae. Muf. D. Lund.
*Variat* pedibus fufcis et flaueſcentibus.
*Os* maxillis palpisque. *Palpi* quatuor inaequales, *anteriores* longiſſimi, exſerti, quadriarticulati: articulo primo breuiſſimo, ſecundo tertioque longioribus, filiformibus, quarto longo, ouato, adhaerentes maxillae dorſo; *poſteriores* breuiores, triarticulati: articulo primo breuiſſimo, ſecundo longiori, filiformi, tertio ouato, craſſiori, adnati ligulae medio. *Labrum* reconditum, breue, transuerſum, obtuſum. *Mandibula* cornea, arcuata, acuta, apice bifida. *Maxilla* breuis, membranacea, bifida: laciniis aequalibus obtuſis. *Ligula* breuis, membranacea, trifida: lacinia media breuiſſima, lateralibus elongatis, diuaricatis, acuminatis. *Labium* breue, transuerſum, integrum.

. St. ater, nitidus, abdomine conico. tibiis teſtaceis. *ſpiſſicornis*
Habitat in America meridionali. D. Smidt. Muſ. D. de Seheſtedt.
Paruus, ater, nitidus. Antennae breues, claua ſolida. Abdomen poſtice attenuatum, ſubconicum. Pedes teſtacei, femoribus nigris.

. St. ater, nitidus, abdomine conico, elytris pedibusque teſtaceis. *conicus.*
Habitat in America meridionali. D. Smidt. Muſ. D. Lund.
Statura et ſumma affinitas praecedentis. Differt tantum elytris pedibusque totis teſtaceis.

. St. niger, elytris obſcuris: macula poſtica repanda pallida. *Ficus.*
Habitat in Oriente, lectus inter ficus fructus ſiccatos. D. Amthor.

Sta-

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Statura parua, craſſa, breuis. Antennae nigrae. Caput et thorax nigra. Elytra abbreuiata, obſcura: macula magna, repanda, pallida. Corpus nigrum, ano pedibusque ferrugineis.

*floralis.* 6. St. niger, pedibus flaueſcentibus.
*Staphylinus floralis.* Ent. ſyſt. 2. 530. 52.
*Staphylinus floralis.* Payk. Monogr. 67. 47.
I. 62 Fn. Sv. 3. 406. 49.
Panz. Fn. Germ. 34. tab. 20.
Habitat in Galliae floribus. Muſ. D. Boſc.

180. OXYPORVS. *Palpi* inaequales: *anteriores* filiformes; *poſteriores* ſecuriformes.
*Ligula* emarginata cum mucrone.
*Antennae* moniliformes.

*rufus.* 1. O. rufus, capite elytrorum abdominisque poſtico nigris. Ent. ſyſt. 2. 531. 1.
*Staphylinus rufus.* Linn. ſyſt. Nat. 2. 684. 6. Fn. Sv. 844.
Oliv. Inſ. 43. tab. 1. fig. 1.
II. 64. Panz. Fn. Germ. 16. tab. 19.
Geoff. Inſ. 1. 370. 22.
Schaeff. Icon. tab. 85. fig. 3.
Degeer Inſ. 4. 24. 10. tab. 1. fig. 11. 12.
Payk. Monogr. 18. 18.
Fn. Sv. 3. 425. 1. Ha-

---

*Oxypori* corpus paruum, elongatum, glabrum, immarginatum, agile; *capite* ouato, magno, exſerto; *mandibulis* porrectis, forficatis; *oculis* globoſis, prominulis, lateralibus; *antennis* breuibus, diſtantibus, infra oculos inſertis; *thorace* rotundato: margine deflexo; *ſcutello* paruo, rotundato; *elytris* dimidiatis, rigidis, fornicatis, alas tegentibus; *pedibus* longiuſculis, tenuibus; *tarſis* quadriarticulatis; *colore* vario.

Digitized by Google Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Habitat in Europae Boletis.

O. ater, elytris pallidis: angulo poſtico nigro, abdomine rufo: ano fuſco. Ent. ſyſt. 2. 531. 2. * *maxilloſus*

Panz. Fn. Germ. 16. tab. 20.

Habitat in Germania. D. Schneider.

O. flauus, elytris nigris, baſi apiceque pallidis. Ent. ſyſt. 2. 532. 3. * *lunulatus.*

*Staphylinus lunulatus.* Linn. ſyſt. Nat. 2. 689. 7. Fn. Sv. 845.

Payk. Monogr. 49. 35.
Fn. Sv. 3. 415. 61.

Habitat in Europae Boletis.

O. ater, laeuis, nitidus, elytris macula baſeos oblonga, teſtacea. Ent. ſyſt. 2. 532. 4. *ſubterraneus.*

*Staphylinus ſubterraneus.* Linn. ſyſt. Nat. 2. 690. 9.

Payk. Monogr. 52.
Fn. Sv. 3. 420. 66.

Habitat in Suecia. D. de Paykull.

O. ater, thorace vtrinque puncto baſeos rufo. Ent. ſyſt. 2 532. 5. * *cellaris.*

*Staphylinus littoreus.* Linn. ſyſt. Nat. 2. 685. 18. Fn. Sv. 852.

*Staphylinus pubeſcens.* Payk. Monogr. Curc. App. 138. 33.

*Staphylinus cellaris.* Payk. Fn. Sv. 3. 421. 76.

Habitat in Sueciae cellis ſubterraneis. D. de Payk.

O. ater, nitidus, thoracis margine pedibus elytrisque rufis: his futura maculaque marginali nigris. Ent. ſyſt. 2. 532. 6. * *marginatus.*

Panz. Fn. Germ. 27. tab. 17.

Habitat in Suecia. Muſ. D. de Seheſtedt.

O. thorace rufo, coleoptris palleſcentibus: maculis tribus nigris. Ent. ſyſt. 2. 533. 7. * *3 maculatus.*

*Sta-*

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Staphylinus littoreus.* Payk. Monogr. 80. 55.
*Staphylinus 3maculatus.* Payk. Fn. Sv. 3. 422. 68.
Habitat in Dania. Muſ. D. de Seheſtedt.

*3maculatus.* 8. O. niger, elytris macula baſeos pedibusque pallidis. Ent. ſyſt. 2. 533. 8. *
Habitat in Selandia. Muſ. D. de Seheſtedt.

*2puſtulatus* 9. O. ater, nitidus, elytris macula baſeos anoque rufis. Ent. ſyſt. 2. 533. 9.
II. 68 Panz. Fn. Germ. 16. tab. 21.
Habitat in Germania. D. Smidt.

*analis.* 10. O. flauus, elytris baſi anoque atris. Ent. ſyſt. 2. 533. 10.
II. 69. Habitat Kiliae. D. Daldorff.

*thoracicus* 11. O. nigricans, thorace rufo, elytris fuſcis, baſi albis. Ent. ſyſt. 2. 534. 12. *
Habitat in Holſatiae Boletis. Muſ. D. de Seheſtedt.

*pygmaeus.* 12. O. nigricans, thorace elytrisque pallidis. Ent. ſyſt. 2. 534. 13. *
III. 11. Panz. Fn. Germ. 27. tab. 19.
Habitat Kiliae in Boletis. Muſ. D. de Seheſtedt.

*flauipes.* 13. O. niger, elytris pedibusque fuſcis. Ent. ſyſt. 2. 534. 13. *
Habitat in Europa boreali.

*chriſomelinus.* 14. O. ater, thorace rufo, elytris teſtaceis: margine baſeos atro. Ent. ſyſt. 2. 534. 14. *
*Staphylinus chryſomelinus.* Linn. ſyſt. Nat. 2. 685. 2. Fn. Sv. 855.
Geoff. Inſ. 1. 368. 18.
Oliv. Inſ 42. tab. 3. fig. 22.
I. 63. Panz. Fn. Germ. 9. tab. 14.
Payk. Monogr. 77. 54.
Fn. Sv. 3. 423. 69.
Habitat in Europae Boletis.

15.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

5. O. ater, nitidus, thoracis margine poſtico, elytris abdominis ſegmentorum marginibus pedibusque teſtaceis. Ent. ſyſt. ſuppl. 182. 15. * *dimidiatus*

Habitat in Germania. D. Daldorff.

5. O. teſtaceus, capite abdomineque atris. Ent. ſyſt. 2. 534. 16. * *melanocephalus.*

Habitat in Daniae Selandia. Muſ. D. de Seheſtedt.

7. O. rufus, abdomine atro nitido. Ent. ſyſt. 2. 535. 17. * *abdominalis.*

Habitat in Norwagia. Muſ. D. Lund.

3. O. teſtaceus, abdomine obſcuriore. Ent. ſyſt. 2. 535. 18. * *teſtaceus.*

Habitat in Germania. D. Smidt.

9. O. brunneus, capite thoracisque medio atris. Ent. ſyſt. 2. 535. 19. * *brunneus.*

Habitat in Germania. D. Smidt.

0. O. nigricans, elytris pallidis. Ent. ſyſt. 2. 535. 20. * *minutus.*

Habitat in Selandia. Muſ. D. de Seheſtedt.

1. O. ater, nitidus, pedibus rufis. Ent. ſyſt. 2. 535. 21. *rufipes.*

*Staphylinus rufipes.* Linn. ſyſt. Nat. 2. 685. 21.
Fn. Sv. 855.
Geoff. Inſ. 1. 368. 18.
Panz. Fn. Germ. 27. tab. 20.

III. 71

Habitat in Europa boreali.

2. O. ater, nitidus, thoracis margine elytris pedibusque teſtaceis. Ent. ſyſt. 2. 535. 22. *hypnorum.*

Habitat in Anglia.

3. O. ater, nitidus, thoracis elytrorumque marginibus ferrugineis. Ent. ſyſt. 2. 536. 23. * *marginellus.*

*Staphylinus rufipes.* Payk. Fn. Sv. 3. 419. 65. *e.*
Monogr. 76. 53. β.

I. 63

Panz.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Panz. Fn. Germ. 9. tab. 13.
Habitat Kiliae.

181. PAEDERVS. *Palpi* inaequales:
*anteriores* clauati;
*posteriores* filiformes.
*Ligula* linearis, integra.
*Antennae* moniliformes.

*riparius.* 1. P. rufus, elytris coeruleis, capite abdominisque apice nigris. Ent. syst. 2. 536. 1. ♀
*Staphylinus riparius.* Linn. syst. Nat. 2. 684. 8. Fn. Sv. 846.
Geoff. Inf. 1. 369. 21.
Oliv. Inf. 44. tab. 1. fig. 2.
Schaeff. Icon. tab. 71. fig. 3.
Degeer Inf. 4. 28. 14. tab. 1. fig. 18.
Panz. Fn. Germ. 9. tab. 1.
I. 63 Payk. Monogr. 27. 19.
Fn. Sv. 427. 2.
Habitat in Europa, in America meridionali et in India orientali.

*ruficollis.* 2. P. niger, thorace rufo, elytris cyaneis. Ent. syst. 2. 537. 2.
*Staphylinus ruficollis.* Payk. Monogr. 26. 18.
Oliv. Inf. 44. tab. 1. fig. 1.
III. 72 Panz. Fn. Germ. 27. tab. 22.
Geoff.

---

*Paederi* corpus elongatum, filiforme, glabrum, immarginatum, agile; *capite* subgloboso, exserto; *oculis* globosis, prominulis, lateralibus; *antennis* mediocribus, distantibus, infra oculos insertis; *thorace* cylindrico: margine deflexo; *scutello* minuto; *elytris* dimidiatis, rigidis, fornicatis, alas tegentibus; *pedibus* breuibus, cursoriis; *tarsis* quadriarticulatis: *colore* vario.

Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

Geoff. Inf. 1. 370. 23.
Payk. Fn. Sv. 3. 427. 1.
Herbſt. Arch. tab. 48. fig. 5.
Habitat Kiliae, ſub lapidibus. Muſ. D. de Seheſtedt.

P. niger, elytris poſtice pedibusque fuluis. Ent. *elongatus.*
ſyſt. 2. 537. 3.
*Staphylinus elongatus.* Linn. ſyſt. Nat. 2. 685. 14.
Payk. Monogr. 25. 17.
Fn. Sv. 428. 2.
Habitat in Europa.

P. ater, nitidus, elytris totis pedibusque teſtaceis. *fuluipennis*
Ent. ſyſt. 2. 537. 4. *
Panz. Fn. Germ. 27. tab. 23.
Habitat Kiliae.

P. niger, pedibus teſtaceis. Ent. ſyſt. 2. 537. 5. * *brunnipes.*
Habitat in Holſatia.

P. ater, nitidus, elytris plantisque teſtaceis, capi- *fulgidus.*
te punctulato. Ent. ſyſt. 2. 537. 6. *
Panz. Fn. Germ. 34. tab. 20.
Habitat Kiliae. Dom. Daldorff.

P. ater, antennis thoraceque rufis, elytris pedi- *tricolor.*
busque teſtaceis. Ent. ſyſt. 2. 537. 7. *
Payk. Monogr. 23. 15.
Fn. Sv. 3. 378. 15.
Habitat in Daniae Boletis.

P. filiformis, niger, pedibus piceis. Ent. ſyſt. 2. *filiformis.*
538. 8. *
*Staphylinus quadratus.* Payk. Monogr. 29. 21.
*Paederus filiformis.* Payk. Fn. Sv. 3. 429. 4.
Habitat in Germania. D. Smidt.

P. filiformis, niger, capite orbiculato. Ent. ſyſt. *orbiculatus*
2. 538. 9.
*Staphylinus orbiculatus.* Payk. Monogr. 35. 26.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google

*Paederus orbiculatus.* Payk. Fn. Sv. 3. 431.
Oliv. Inf. 42. tab. 1. fig. 7.
Panz. Fn. Germ. 34. tab. 21.
Habitat in Suecia, varius.

*melanocephalus.* 10. P. filiformis, niger, thorace pedibusque fulv
Ent. syst. 2. 538. 10. *
Panz. Fn. Germ. 36. tab. 24.
Habitat in Germania. Dom. Schneider.



Digitized by Google
Original from NEW YORK PUBLIC LIBRARY
Generated for Joseph Cora (The Ohio State University) on 2013-11-29 16:27 GMT / http://hdl.handle.net/2027/nyp.33433007345717
Public Domain, Google-digitized / http://www.hathitrust.org/access_use#pd-google