

ILLUSTRATIONS
OF
BRITISH ENTOMOLOGY ;
OR, A
Synopsis of Indigenous Insects:
CONTAINING
THEIR GENERIC AND SPECIFIC DISTINCTIONS ;
WITH
AN ACCOUNT OF THEIR METAMORPHOSSES, TIMES OF
APPEARANCE, LOCALITIES, FOOD, AND ECONOMY,
AS FAR AS PRACTICABLE.

BY JAMES FRANCIS STEPHENS, F.L.S.,
MEMBER OF THE ENTOMOLOGICAL SOCIETIES OF LONDON AND PARIS, ETC.

EMBELLISHED WITH COLOURED FIGURES OF THE RARER AND
MORE INTERESTING SPECIES.

S U P P L E M E N T.

"In his tam parvis tamque ferè nullis quæ ratio ! quanta vis ! quam inextricabilis
perfectio!"—*Plin.*
"Finis Creationis Telluris est gloria DEI, ex opera Naturæ, per Hominem
solum."—*Linné.*

LONDON :
HENRY G. BOHN, YORK STREET, COVENT GARDEN.

—
1867.

C. C. BABINGTON, F.L.S., F.G.S.,	JNO. E. GRAY, F.R.S., &c.
Sec. Camb. Phil. S.	
W. BAIRD, M.D., Brit. Mus.	J. S. HENSLOW, M.A., F.L.S.,
JNO. BARLOW, Sec. R.S., F.R.S.,	Prof. Botany, Cambridge.
&c.	
EDW. BARNARD, F.L.S.	W. J. HOOKER, F.R.S., &c.
THOS. BELL, F.R.S., F.L.S., &c.	F. W. HOPE, F.R.S., &c.
Prof. Zoology, King's Coll.	TUOS. HORSFIELD, M.D., F.R.S.,
J. J. BENNETT, F.R.S., F.L.S., &c.	&c.
FRAS. BOOT, M.D., F.L.S., &c.	WM. KIRBY, F.R.S., &c.
J. BOSTOCK, M.D., F.R.S., &c.	CH. KONIG, F.R.S., &c.
J. S. BOWERBANK, F.R.S., &c.	WM. NORTON LLOYD, F.R.S.,
W. J. BRODERIP, F.R.S., &c.	F.L.S., &c.
RT. BROWN, V.P.L.S., &c.	EDW. NEWMAN, F.L.S.
WALTER BUCHANAN, F.L.S.,	RICHD. OWEN, Hunterian Pro-
F.R.S., &c.	fessor Royal Coll. of Surgeons,
WM. BUCKLAND, F.R.S., &c.	F.R.S., &c.
RD. CHAMBERS, F.L.S.	J. J. S. PARRY, F.L.S.
JNO. GEO. CHILDREN, F.R.S., &c.	J. F. ROYLE, M.D., F.R.S., Prof.
BRACEY CLARK, F.L.S., &c.	Mat. Med. &c., King's Coll.
CH. R. DARWIN, F.R.S., &c.	W. W. SAUNDERS, F.L.S., S.A.,
L. W. DILLWYN, F.R.S., &c.	&c.
EDW. DOUBLEDAY, F.L.S., &c.	R. H. SOLLY, F.R.S., F.L.S., &c.
HY. DOWNES, Com. R.N., F.L.S.,	G. B. SOWERBY, F.L.S., &c.
&c.	WM. SPENCE, F.R.S., &c.
W. F. EVANS, M.E.S.	GEO. T. STAUNTON, Bart., M.P.,
F. J. FARRE, M.D., F.L.S., &c.	F.R.S., &c.
EDW. FORSTER, V.P.L.S., &c.	CH. STOKES, F.R.S., &c.
J. D. E. FORTNUM, M.E.S.	W. H. SYKES, Lt.-Col., F.R.S.,
JNO. GOULD, F.R.S., &c.	&c.
GEO. GRANT, M.D.	A. TULK, M.R.C.S.
ROBT. E. GRANT, M.D., F.R.S.,	GEO. R. WATERHOUSE, Brit. Mus.
L. & E. Prof. Zool. University	
Coll.	J. O. WESTWOOD, F.L.S., Sec.
GEO. R. GRAY, F.L.S., &c.	Ent. S., &c.
	ADAM WHITE, Brit. Mus.
	WM. WOOD, F.R.S., &c.
	WM. YARRELL, F.L. & Z.S.

Eltham Cottage, Foxley Road, Kennington,

September, 1816.

ILLUSTRATIONS
OF
BRITISH ENTOMOLOGY.

MANDIBULATA.

HYMENOPTERA.

FAMILY V. ICHNEUMONIDÆ, Leach.

Plate XXXIX. fig. 3.

Phytodictus segmentator.

Pn. *Niger*, marginibus segmentorum albidis; pedibus rufis anticorum trochanteribus et coxis flavis, posticorum tarsis et tibiarum apice nigris. (Long. Corp. 2½ 3½ lin. Exp. Alar. 7 lin.)

Phy. *segmentator*, Gravenhorst, ii. 944.—Steph. Nomen. 2 edit.

Black, with the margin of the abdominal segments whitish; legs red, slender, with the trochanters of the anterior and the coxae yellow, the tarsi of the posterior, and the apex of the tibiae black:—the male has the face yellow, and the posterior coxae and trochanters varied with black and yellow: the female has the face marked with yellow, and the hinder coxae red.!

Apparently rare: taken in the vicinity of London.

Plate XXXIX. fig. 2.

Schizopyga analis. ?

Sc. *Nigra*, abdomine rufo, margine segmentorum 1—3 annoque nigris; pedibus rufis, nigro maculatis; aculeo subexerto. (Long. Corp. 3—3½ lin.)

Sch. *analisis*, Gravenhorst, iii. 130.—Steph. Nomen. 2 edit.

Black: mouth and antennæ beneath pale testaceous: abdomen with the basal segment black, with its sides and back red, 2nd and 3rd segments red, the edges black, two following wholly red, the rest black: legs red, the tips of the four hinder thighs and tibiae black: hinder tarsi black, the base of the joints testaceous: female with the ovipositor slightly exserted.

Taken at Darenth, in June.

MANDIBULATA—SUPPLEMENT, AUGUST, 1846.

PLATE I.




PLATE II.

PLATE II.

Published by J. F. Smith, Boston.

Plate XLVI. fig. 3.

Cyrositus Walkeri.

Dr. *Flavus, macula thoracis, abdominisque apice viridi-aeneis, tibiis intermediis apicibus nigris.* (Long. Corp. $1\frac{1}{4}$ lin.)

Cl. *Walkeri, Steph. Nomen. 2d edit.*

Yellow; ocelli black; eyes fuscous; an irregular blotch on the thorax, and the metathorax and scutellum brassy-green; abdomen yellow, its tip greenish-black; sides of thorax yellow, with a greenish apex: legs yellow, the apex of the middle tibiae and tips of the hinder tarsi, black.

Found in June, near London.

PROCTOTRUPIDÆ.

Plate XLVI. fig. 2.

Diapria conica. ♀

Dr. *Antennis apicem versus incrassatis, clara subsexarticulata, nigra, scapo pedibusque pedis, tibiis pallidioribus; abdome conico-acuminato.*
Mas. *antennis corpore longioribus fuscis, scapo pedibusque rufis, flagelli articulis elongato-cylindricis verticillato pilosis, secundo basi retrosum distincte emarginato; ano obtuso.* (Long. Corp. $1\frac{1}{2}$ — $1\frac{3}{4}$ lin.)

Ich. conicus, Fabricius.—Di. conica, Steph. Cat. No. 5601.

Black: Female with the antennæ thickened towards the apex, the club black, and obscurely six-jointed, the base and the legs pitchy, the tibiae paler; abdomen conic-acuminate. Male with the antennæ longer than the body, fuscous, the scape red, the joints elongate-cylindric, verticillate pilose, the second joint distinctly notched towards the base: anus obtuse.

Not uncommon in the vicinity of London in the summer.

GONATOPIDÆ.

Plate XLVII. fig. 2.

Anteon flavicorne.

An. *Nigrum nitidum, fronte albæ-sericeâ; ore albo; antennis (feminæ) thorace brevioribus apice crassioribus et pedibus testaceis: metathorace leviter cancellato.* (Long. Corp. 1½—2 lin.)

Dr. *flavicornis*, *Dalman*.—Go. *flavicornis*, *Steph. Catal.* No. 5673.

Black, and shining: forehead with a whitish pile; mouth white, antennæ of the female rather shorter than the thorax, thickened at the apex, and, with all the legs, pale testaceous; metathorax slightly cancellated.

Found near London, in woods and shady places, in July.

SCELIONIDÆ.

Plate XLVII. fig. 3.

Teleas Procris.

Te. *Ater, apterus, pedibus piceis, antennis mari filiformibus.* (Long. Corp. ¾ lin.)

Te. *Procris*, *Walker. Ent. M. iii. 866.—Steph. Nomen. 2d edit.*

Black, punctured; slightly pubescent; wingless; legs pitchy, the knees, tips of the tibiae and the tarsi reddish; abdomen with the basal segment, and the greater portion of the 2nd deeply striated.

Common in grassy places about London.

PLATE II.



PLATE III.

London Published by J. E. Suckling, Oct. 30th 1846.